

## Package leaflet: information for the user

### **Bexsero suspension for injection in pre-filled syringe** Meningococcal group B Vaccine (rDNA, component, adsorbed)

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

**Read all of this leaflet carefully before you or your child receive this medicine because it contains important information for you or your child.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or nurse.
- This vaccine has been prescribed for you or your child only.
- If you get any side effects talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

1. What Bexsero is and what it is used for
2. What you need to know before you or your child receive Bexsero
3. How to use Bexsero
4. Possible side effects
5. How to store Bexsero
6. Contents of the pack and other information

#### **1. WHAT BEXSERO IS AND WHAT IT IS USED FOR**

Bexsero is a Meningococcal group B Vaccine.

Bexsero contains four different components from the surface of the bacteria *Neisseria meningitidis* group B.

Bexsero is given to individuals from 2 months of age and older to help protect against disease caused by the *Neisseria meningitidis* group B bacteria. These bacteria can cause serious, and sometimes life-threatening, infections such as meningitis (inflammation of the covering of the brain and spinal cord) and sepsis (blood poisoning).

The vaccine works by specifically stimulating the body's natural defense system of the vaccinated person. This results in protection against the disease.

#### **2. WHAT YOU NEED TO KNOW BEFORE YOU OR YOUR CHILD RECEIVE BEXSERO**

##### **Do NOT use Bexsero:**

- If you or your child are allergic to the active substances or any of the other ingredients of this vaccine (listed in section 6).

## **Warnings and precautions**

Talk to your doctor or nurse before you or your child receive Bexsero, if you or your child have:

- a severe infection with a high temperature. If this is the case, then vaccination will be postponed. The presence of a minor infection, such as a cold, should not require postponement of the vaccination, but talk to your doctor or nurse first.
- haemophilia or any other problem that may stop your blood from clotting properly, such as treatment with blood thinners (anticoagulants). Talk to your doctor or nurse first.
- an allergy to the antibiotic kanamycin. If present, the kanamycin level in the vaccine is low. If you or your child may have allergy to kanamycin, talk to your doctor or nurse first.

Tell your doctor or nurse if you know that you or your child is allergic to latex. The tip cap of the syringe may contain natural rubber latex. The risk for developing an allergic reaction is very small, but your doctor or nurse needs to be aware of your allergy when deciding if you or your child should receive Bexsero.

There are no data on the use of Bexsero in adults above 50 years of age or in patients with chronic medical conditions or with weakened immunity. If you or your child have weakened immunity (for example, due to the use of immunosuppressive medications, or HIV infection, or hereditary defects of the body's natural defence system), it is possible that the effectiveness of Bexsero is reduced.

As with any vaccine, Bexsero may not fully protect all of those who are vaccinated.

## **Other medicines and Bexsero**

Tell your doctor or nurse if you or your child are taking, have recently taken, or might take any other medicines, or have recently received any other vaccine.

Bexsero can be given at the same time as any of the following vaccine components: diphtheria, tetanus, whooping cough (pertussis), *Haemophilus influenzae* type b, polio, hepatitis B, pneumococcus, measles, mumps, rubella, and chickenpox. Talk to your doctor or nurse for further information.

When given at the same time with other vaccines Bexsero must be given at separate injection sites.

Your doctor or nurse may ask you to give your child medicines that lower fever at the time and after Bexsero has been given. This will help to reduce some of the side effects of Bexsero.

## **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before Bexsero is given. Your doctor may still recommend that you receive Bexsero if you are at risk of exposure to meningococcal infection.

## **Driving and using machines**

Bexsero has no or negligible influence on the ability to drive and use machines. However, some of the effects mentioned under section 4 "Possible side effects" may temporarily affect the ability to drive or use machines.

## **Bexsero contains sodium chloride**

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially 'sodium-free'.

### **3. HOW TO USE BEXSERO**

Bexsero (0.5 ml) will be given to you or your child by a doctor or nurse. It will be injected into a muscle, usually the thigh for infants or the upper arm for children, adolescents and adults.

It is important to follow the instructions from the doctor or nurse so that you or your child completes the course of injections.

#### **Infants 2 months to 5 months of age**

Your child should receive an initial course of three injections of the vaccine followed by a fourth injection (booster).

- The first injection should be given at 2 months of age.
- The interval between each injection should be at least 1 month.
- A fourth injection (booster) will be given between 12 months and 15 months of age. In case of delay, the booster should not be given later than 24 months.

#### **Infants 6 months to 11 months of age**

Unvaccinated infants 6 months to 11 months of age should receive two injections followed by a third injection (booster).

- The interval between the injections should be at least 2 months.
- A third injection (booster) will be given in the second year of life after an interval of at least 2 months from the second injection.

#### **Children 12 months to 23 months of age**

Children 12 months to 23 months of age should receive two injections followed by a third injection (booster).

- The interval between each injection should be at least 2 months
- A third injection (booster) will be given after an interval of 12 to 23 months from the second injection.

#### **Children 2 years to 10 years of age**

Children 2 years to 10 years of age should receive two injections.

- The interval between each injection should be at least 2 months.

#### **Adolescents (from 11 years of age) and adults**

Adolescents (from 11 years of age) and adults should receive two injections.

- The interval between each injection should be at least 1 month.

#### **Adults above 50 years of age**

There are no data in adults above 50 years of age. Ask your doctor for advice whether it is beneficial for you to receive Bexsero.

If you have any further questions on Bexsero, ask your doctor or nurse.

#### 4. POSSIBLE SIDE EFFECTS

Like all vaccines, this vaccine can cause side effects, although not everybody gets them.

When Bexsero is given to you or your child, the very common side effects (may affect more than 1 in 10 people) that you or your child may get (reported in all age groups) are:

- pain/tenderness at the injection site, redness of the skin at the injection site, swelling of the skin at the injection site, hardness of the skin at the injection site.

The following side effects may also occur after receiving this vaccine.

##### **Infants and children (up to 10 years of age)**

**Very common** (these may affect more than 1 in 10 people)

- fever ( $\geq 38^{\circ}\text{C}$ )
- loss of appetite
- tenderness or discomfort at the injection site (including severe injection site tenderness resulting in crying when injected limb is moved)
- skin rash (children aged 12 to 23 months) (uncommon after booster)
- sleepiness
- feeling irritable
- unusual crying
- vomiting
- diarrhea

**Common** (these may affect up to 1 in 10 people)

- skin rash (infants and children 2 to 10 years of age)

**Uncommon** (these may affect up to 1 in 100 people)

- high fever ( $\geq 40^{\circ}\text{C}$ )
- seizures (including febrile seizures)
- vomiting (after booster)
- dry skin
- paleness (rare after booster)

**Rare** (these may affect up to 1 in 1,000 people)

- Kawasaki disease which may include symptoms such as fever that lasts for more than five days, associated with a skin rash on the trunk of the body, and sometimes followed by a peeling of the skin on the hands and fingers, swollen glands in the neck, red eyes, lips, throat and tongue
- Itchy rash, skin rash

##### **Adolescents (from 11 years of age) and adults**

**Very common** (these may affect more than 1 in 10 people)

- pain at the injection site resulting in inability to perform normal daily activity
- painful muscles and joints
- nausea
- generally feeling unwell
- headache

##### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Medicines and Healthcare products Regulatory Agency (MHRA) at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard). By reporting side effects you can help provide more information on the safety of this medicine.

## 5. HOW TO STORE BEXSERO

Keep this vaccine out of the sight and reach of children.

Do not use this vaccine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

Store in a refrigerator (2°C – 8°C). Do not freeze.  
Store in the original package in order to protect from light.

Do not throw away any medicines via wastewater or household waste. Ask your doctor or nurse how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. CONTENTS OF THE PACK AND OTHER INFORMATION

### What Bexsero contains

One dose (0.5 ml) contains:

Active substances:

Recombinant <i>Neisseria meningitidis</i> group B NHBA fusion protein <sup>1,2,3</sup>	50 micrograms
Recombinant <i>Neisseria meningitidis</i> group B NadA protein <sup>1,2,3</sup>	50 micrograms
Recombinant <i>Neisseria meningitidis</i> group B fHbp fusion protein <sup>1,2,3</sup>	50 micrograms
Outer membrane vesicles (OMV) from <i>Neisseria meningitidis</i> group B strain NZ98/254 measured as amount of total protein containing the PorA P1.4 <sup>2</sup>	25 micrograms

<sup>1</sup> produced in *E. coli* cells by recombinant DNA technology

<sup>2</sup> adsorbed on aluminium hydroxide (0.5 mg Al<sup>3+</sup>)

<sup>3</sup> NHBA (Neisseria Heparin Binding Antigen), NadA (Neisserial adhesin A), fHbp (factor H binding protein)

Other ingredients:

Sodium chloride, histidine, sucrose and water for injections (see section 2 for further information on sodium and latex).

### What Bexsero looks like and contents of the pack

Bexsero is a suspension for injection in pre-filled syringe (Type I glass) with a plunger stopper (Type I bromobutyl rubber) and with a protective tip cap (Type I or Type II rubber) with or without needles.

Pack sizes of 1 or 10 syringes.

The suspension is white opalescent liquid.

Not all pack sizes may be marketed.

### Marketing Authorisation Holder:

Novartis Vaccines and Diagnostics S.r.l.  
Via Fiorentina 1  
53100 Siena  
Italy.

**Manufacturer:**

Novartis Vaccines and Diagnostics S.r.l.  
Bellaria-Rosia  
53018 Sovicille (Siena)  
Italy.

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Novartis Vaccines and Diagnostics S.r.l.  
Tél/Tel: +39 0577 243638  
Italië/Italie/Italien

**България**

Novartis Vaccines and Diagnostics S.r.l.  
Тел.: +39 0577 243638  
Италия

**Česká republika**

Novartis s.r.o.  
Vaccines & Diagnostics  
Gemini B  
Na Pankráci 1724/129  
140 00 Prague 4  
Czech Republic  
+420 225 775 111

**Danmark**

Novartis Vaccines and Diagnostics S.r.l.  
Tlf: +39 0577 243638  
Italien

**Deutschland**

Novartis Vaccines Vertriebs GmbH  
Rudolf-Diesel-Ring 27  
D-83607 Holzkirchen  
Tel: + 49 (8024) 646 5777

**Eesti**

Novartis Vaccines and Diagnostics S.r.l.  
Tel: +39 0577 243638  
Itaalia

**Ελλάδα**

Novartis (Hellas) A.E.B.E.  
Τηλ: +30 210 281 17 12

**España**

Novartis Vaccines and Diagnostics, S.L.  
Gran Via de les Corts Catalanes 764  
08013 Barcelona  
tel: +34 93.306.42.00

**Lietuva**

Novartis Vaccines and Diagnostics S.r.l.  
Tel: +39 0577 243638  
Italija

**Luxembourg/Luxemburg**

Novartis Vaccines and Diagnostics S.r.l.  
Tél/Tel: +39 0577 243638  
Italie/Italien

**Magyarország**

Novartis Hungária Kft.  
Vakcina Divízió  
Bartók Béla út 43-47.  
H-1114 Budapest  
Tel.: + 36 1279 1744

**Malta**

Novartis Vaccines and Diagnostics S.r.l.  
Tel: +39 0577 243638  
L-Italja

**Nederland**

Novartis Vaccines and Diagnostics S.r.l.  
Tel: +39 0577 243638  
Italië

**Norge**

Novartis Vaccines and Diagnostics S.r.l.  
Tlf: +39 0577 243638  
Italia

**Österreich**

Novartis Pharma GmbH  
Tel: +43 1 86 6570

**Polska**

Novartis Poland Sp. z. o. o.  
Marynarska 15  
02-674 Warszawa  
Polska  
Tel: +48 22 3754888

**France**

Novartis Vaccines and Diagnostics SAS  
10 rue Chevreul  
92 150 Suresnes France  
tél: 00 33 1 55 49 00 30

**Hrvatska**

Novartis Vaccines and Diagnostics S.r.l.  
Tel: +39 0577 243638  
Italija

**Ireland**

Novartis Vaccines and Diagnostics S.r.l.  
Tel: +39 0577 243638  
Italy

**Ísland**

Novartis Vaccines and Diagnostics S.r.l.  
Sími: +39 0577 243638  
Ítalía

**Italia**

Novartis Vaccines and Diagnostics s.r.l.  
Via Fiorentina 1  
53100 Siena  
Tel: 800867121

**Κύπρος**

Novartis Vaccines and Diagnostics S.r.l.  
Τηλ: +39 0577 243638  
Ιταλία

**Latvija**

Novartis Vaccines and Diagnostics S.r.l.  
Tel: +39 0577 243638  
Itālija

**Portugal**

Novartis Farma - Produtos Farmacêuticos, S.A.  
Tel: +351 21 000 8600

**România**

Novartis Vaccines and Diagnostics S.r.l.  
Tel: +39 0577 243638  
Italia

**Slovenija**

Novartis Vaccines and Diagnostics S.r.l.  
Tel: +39 0577 243638  
Italija

**Slovenská republika**

Novartis Slovakia s.r.o.  
Galvaniho 15/A  
821 08 Bratislava  
Slovenská republika  
Tel: + 421 2 5070 6111

**Suomi/Finland**

Novartis Vaccines and Diagnostics S.r.l.  
Puh/Tel: +39 0577 243638  
Italia/Italien

**Sverige**

Novartis Vaccines and Diagnostics S.r.l.  
Tel: +39 0577 243638  
Italien

**United Kingdom**

Novartis Vaccines and Diagnostics Limited  
Gaskill Road Speke  
Liverpool L24 9GR  
Tel: +44(0) 845 745 1500

**This leaflet was last revised in June 2014**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu/>

-----

**The following information is intended for healthcare professionals only:**

Upon storage a fine off-white deposit may be observed in the pre-filled syringe containing the suspension.

Before use, the pre-filled syringe should be well shaken in order to form a homogeneous suspension.

The vaccine should be visually inspected for particulate matter and discoloration prior to administration. In the event of any foreign particulate matter and/or variation of physical aspect being observed, discard the vaccine.

Do not freeze.

Bexsero must not be mixed with other vaccines in the same syringe.

Should concomitant administration of other vaccines be necessary, vaccines must be administered at separate injection sites.

Care must be taken to ensure that the vaccine is injected intramuscularly only.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.