

Package leaflet: Information for the user

THORENS 25 000 I.U. capsules, hard

Colecalciferol (vitamin D₃)

Read all of this leaflet carefully before you start using this medicine, because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist, or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What THORENS is and what it is used for
2. What you need to know before you use THORENS
3. How to use THORENS
4. Possible side effects
5. How to store THORENS
6. Contents of the pack and other information

1. What THORENS is and what it is used for

THORENS hard capsules contains the active ingredient colecalciferol (vitamin D₃). Vitamin D₃ can be found in some foods and is also produced by the body when skin is exposed to sunlight. Vitamin D₃ helps the kidneys and intestine absorb calcium and it helps build bones. Vitamin D₃ deficiency is the predominant cause of rickets (defective mineralization of bones in children) and osteomalacia (inadequate mineralization of bones in adults).

THORENS hard capsules is used for initial treatment of clinically relevant vitamin D deficiency in adults.

2. What you need to know before you use THORENS

Do not use THORENS:

- if you are allergic to vitamin D₃ or any of the other ingredients of this medicine (listed in section 6);
- if you have high levels of calcium in your blood (hypercalcaemia) or urine (hypercalciuria);
- if you have kidney stones (renal calculi) or severe renal impairment;
- if you have high levels of vitamin D₃ in your blood (hypervitaminosis D)

Warnings and precautions

Talk to your doctor, pharmacist or nurse before using THORENS if you:

- are undergoing treatment with certain medicines used to treat heart disorders (eg, cardiac glycosides, such as digoxin);
- have sarcoidosis (an immune system disorder which may cause increased levels of vitamin D₃ in the body);

- are treated with diuretics (e.g. benzothiadiazine)
- are immobilized
- suffer from pseudohypoparathyroidism
- are taking medicines containing vitamin D₃, or eating foods or milk enriched with vitamin D₃;
- are likely to be exposed to a lot of sunshine whilst using THORENS;
- take additional supplements containing calcium. Your doctor will monitor your blood levels of calcium to make sure they are not too high whilst you are using THORENS;
- have kidney damage or disease and if you have a tendency for the formation of renal stones. Your doctor may want to measure the levels of calcium in your blood or urine.
- take a daily dose of vitamin D₃ exceeding 1,000 I.U. over a long period of time, your doctor should monitor the level of calcium in your blood by lab test.

Children and Adolescents

The use is not recommended in children and adolescents under 18 years of age.

Other medicines and THORENS

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines. This is especially important if you are taking:

- medicines that act on the heart or kidneys, such as cardiac glycosides (eg, digoxin) or diuretics (eg, bendroflumethiazide). When used at the same time as vitamin D₃ these medicines may cause a large increase in the level of calcium in the blood and urine;
- medicines containing vitamin D₃ or eating food rich in vitamin D₃, such as, some types of vitamin D₃-enriched milk;
- actinomycin (a medicine used to treat some forms of cancer) and imidazole antifungals (eg, clotrimazole and ketoconazole, medicines used to treat fungal disease). These medicines may interfere with the way your body process vitamin D₃;
- medicines to treat tuberculosis e.g rifampicin, isoniazid;
- the following medicines because they can interfere with the effect or the absorption of vitamin D₃:
 - antiepileptic medicines (anticonvulsants), barbiturates;
 - glucocorticoids (steroid hormones such as hydrocortisone or prednisolone). These can decrease the effect of vitamin D₃;
 - medicines that lower the level of cholesterol in the blood (such as cholestyramine, or colestipol);
 - certain medicines for weight loss that reduce the amount of fat your body absorbs (eg, orlistat);
 - certain laxatives (such as liquid paraffin).

THORENS with food, drink and alcohol

You should take this medicine preferably together with a meal to help your body absorb the vitamin D₃.

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

This high strength formulation is not recommended for use in pregnant and breastfeeding women.

Driving and using machines

There is limited information on the possible effects of this medicine on your ability to drive. However, it is not expected that it would affect your ability to drive or to operate machinery.

3. How to use THORENS

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The capsule should be swallowed whole (e.g. with water).
You should take THORENS preferably together with a meal.

Use in adults

The recommended dose is one capsule 25.000 IU/week during the first month. After first month, lower doses may be considered.

This initial treatment may be followed by maintenance therapy, as directed by your doctor.

Your doctor will adjust the dose for you.

Paediatric population

THORENS 25.000 I.U. is not recommended in children and adolescents under 18 years of age.

Pregnancy and breastfeeding

THORENS 25.000 I.U. is not recommended.

If you take more THORENS than you should

If you take more medicine than prescribed, stop using this medicine and contact your doctor. If it is not possible to talk to a doctor go to the nearest hospital emergency department and take the medicine package with you.

The most common symptoms of overdose are: nausea, vomiting, excessive thirst, the production of large amounts of urine over 24 hours, constipation and dehydration, high levels of calcium in the blood and in urine (hypercalcaemia and hypercalciuria) shown by lab test.

If you forget to take THORENS

If you forget to take a dose of THORENS, take the forgotten dose as soon as possible. Then take the next dose at the correct time. However, if it is almost time to take the next dose, do not take the dose you have missed; just take the next dose as normal.

Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Possible side effects may include:

Uncommon (affects less than 1 in 100 people)

- Too much calcium in your blood (hypercalcaemia)
- Too much calcium in your urine (hypercalciuria)

Rare (affects less than 1 in 1000 people)

- Skin rash
- Itching
- Hives

Reporting of side effects

If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store THORENS

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister pack after "Exp". The expiry date refers to the last day of that month.

Do not store above 30° C.

Store in the original package in order to protect from light.

Do not freeze.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What THORENS contains

The active substance is colecalciferol (vitamin D₃).

One capsule contains: 0.625 mg colecalciferol (vitamin D₃) equivalent to 25 000 IU.

The other ingredients are:

- Refined olive oil,
- Gelatine,
- Titanium dioxide (E 171),
- Iron oxide yellow (E 172),
- Iron oxide black (E 172).

What THORENS looks like and contents of pack

THORENS 25 000 I.U. is a hard capsule with transparent body and white cap, with a green band. Capsule dimensions are 15.9 mm x 5.8 mm. It contains an oily solution and it is supplied in Aluminium- PVC/PVDC blisters in cardboard boxes.

Each box may contain 3, 4, 8, 12 or 40 capsules.

Not all pack sizes may be marketed.

Marketing authorisation holder

Galen Limited
Seagoe Industrial Estate
Craigavon
BT63 5UA
UK

Manufacturer

Abiogen Pharma S.p.A.
Via Meucci, 36 – Pisa (Italy)

This medicinal product is authorised in the Member States of the EEA under the following names:

United Kingdom:	THORENS 25.000 IU capsules, hard
Ireland:	THORENS 25.000 IU capsules, hard
Portugal:	Thorens 25,000 IE cápsulas, duras
Netherlands:	THORENS 25.000 IE harde capsules
Germany:	THORENS 25.000 I.E. Hartkapseln
Spain:	Thorens 25.000 UI cápsulas, duras
Norway:	Deltius 25 000 IE harde kapsler
Sweden:	Deltius 25 000 IE hårda kapslar
Poland:	THORENS 25.000 IU kapsułki, twarde
Belgium:	Thorens 25.000 IE capsules, hard Thorens 25 000 UI gélules Thorens 25.000 I.E. Hartkapseln

This leaflet was last revised in September 2021