

Package leaflet: Information for the user

Eslicarbazepine Zentiva 800 mg tablets

eslicarbazepine acetate

Read all of this leaflet carefully before you or your child start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Eslicarbazepine Zentiva is and what it is used for
2. What you need to know before you use Eslicarbazepine Zentiva
3. How to use Eslicarbazepine Zentiva
4. Possible side effects
5. How to store Eslicarbazepine Zentiva
6. Contents of the pack and other information

1. What Eslicarbazepine Zentiva is and what it is used for

The name of your medicine is Eslicarbazepine Zentiva 800mg Tablets (called Eslicarbazepine Zentiva throughout this leaflet). Eslicarbazepine Zentiva contains the active substance eslicarbazepine acetate. Eslicarbazepine Zentiva belongs to a group of medicines called antiepileptics used to treat epilepsy, a condition where someone has repeated seizures or fits.

Eslicarbazepine Zentiva is used:

- on its own (monotherapy) in adult patients with newly diagnosed epilepsy
- with other antiepileptic medicines (adjunctive therapy), in adult, adolescents and children patients above 6 years of age, who are experiencing seizures that affect one part of the brain (partial seizure). These seizures may or may not be followed by a seizure affecting all of the brain (secondary generalisation).

Eslicarbazepine Zentiva has been given to you by your doctor to reduce your number of seizures.

2. What you need to know before you take Eslicarbazepine Zentiva

Do not take Eslicarbazepine Zentiva

- if you are allergic to eslicarbazepine acetate, to other carboxamide derivatives (e.g. carbamazepine or oxcarbazepine, medicines used to treat epilepsy) or to any of the other ingredients of this medicine (listed in section 6);
- if you suffer from a certain type of heart rhythm disorder (second or third degree atrioventricular (AV) block).

Warnings and precautions

Talk to your doctor or pharmacist before taking Eslicarbazepine Zentiva.

Contact your doctor immediately:

- if you have blistering or peeling of the skin and/or mucous membranes, rash, swallowing or breathing problems, swelling of your lips, face, eyelids, throat or tongue. These could be signs of an allergic reaction.
- if you suffer from confusion, worsening of seizures or decreased consciousness which can be signs of low blood salt levels.

Please tell your doctor:

- if you have kidney problems. Your doctor may need to adjust the dose. Eslicarbazepine Zentiva is not recommended in patients with severe renal disease.
- if you have liver problems. Eslicarbazepine Zentiva is not recommended in patients with severe liver problems.
- if you are taking any medicine which can cause an abnormality on the ECG (electrocardiogram) called increased PR interval. If you are not sure if the medicines you are taking could have this effect, discuss with your doctor.
- if you suffer from a heart disease such as heart failure or heart attack, or have any heart rhythm disorder.
- if you suffer from seizures that begin with a widespread electric discharge that involves both sides of the brain.

A small number of people being treated with antiepileptics have had thoughts of harming or killing themselves. If at any time you have these thoughts, when taking Eslicarbazepine Zentiva, contact your doctor immediately.

Eslicarbazepine Zentiva may make you feel dizzy and/or drowsy, particularly at the beginning of treatment. Take special care when taking Eslicarbazepine Zentiva to avoid accidental injury, such as fall.

Take special care with Eslicarbazepine Zentiva: Serious and potentially life-threatening skin reactions including Stevens-Johnson syndrome/toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported in post-marketing experience in patients treated with eslicarbazepine acetate.

If you develop a serious rash or another skin symptoms (see section 4), stop taking Eslicarbazepine Zentiva and contact your doctor or seek medical attention immediately.

In patients of Han Chinese or Thai origin the risk of serious skin reactions associated with carbamazepine or chemically-related compounds may be predicted by testing a blood sample of these patients. Your doctor should be able to advise if a blood test is necessary before taking Eslicarbazepine Zentiva.

Children

Eslicarbazepine Zentiva is not to be given to children aged 6 years and below.

Other medicines and Eslicarbazepine Zentiva

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This is just in case any of them interfere with how Eslicarbazepine Zentiva works or how Eslicarbazepine Zentiva interferes with their effect. Tell your doctor if you are taking:

- phenytoin (a medicine used to treat epilepsy) since your dose may need to be adjusted;
- carbamazepine (a medicine used to treat epilepsy) since your dose may have to be adjusted and the following side effects of Eslicarbazepine Zentiva may occur in higher frequency: seeing double, abnormal coordination and dizziness;
- hormonal contraceptives (such as the contraceptive pill) since Eslicarbazepine Zentiva may make these less effective;
- simvastatin (a medicine used to lower cholesterol levels) since your dose may have to be adjusted;
- rosuvastatin, a medicine used to lower cholesterol level;
- the blood thinner - warfarin;
- tricyclic antidepressants e.g. amitriptyline;
- do not take oxcarbazepine (a medicine used to treat epilepsy) with Eslicarbazepine Zentiva, as it is not known whether it is safe to take these medicines together.

See 'Pregnancy and breast-feeding' section for advice about contraception.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

There are no data from the use of eslicarbazepine acetate in pregnant women. Research has shown an increased risk of birth defects in children of women taking antiepileptic medicines. On the other hand effective antiepileptic therapy must not be interrupted since the worsening of the disease is harmful to both the mother and the unborn child.

Do not breast-feed while you are taking Eslicarbazepine Zentiva. It is not known whether it passes into breast milk.

Eslicarbazepine Zentiva may make hormonal contraceptives such as the contraceptive pill less effective. Therefore, it is recommended that you use other forms of safe and effective contraception, when taking Eslicarbazepine Zentiva up to the end of the current menstrual cycle after stopping treatment.

Driving and using machines

Eslicarbazepine Zentiva may make you feel dizzy, drowsy and affect your vision, particularly at the beginning of treatment. If this happens to you, do not drive or use any tools or machines.

Eslicarbazepine Zentiva contains sodium:

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Eslicarbazepine Zentiva

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Adults

Dose when you start treatment

400 mg once daily for one or two weeks, before increasing to the maintenance dose. Your doctor will decide whether you will be given this dose for one or two weeks.

Maintenance dose

The usual maintenance dose is 800 mg once daily. Depending on how you respond to Eslicarbazepine Zentiva, your dose may be increased to 1,200 mg once daily. If you are taking Eslicarbazepine Zentiva on its own, your doctor may consider you can benefit of a dose of 1,600 mg once daily.

Patients with kidney problems

If you have kidney problems you will usually be given a lower dose of Eslicarbazepine Zentiva. Your doctor will work out the correct dose for you. Eslicarbazepine Zentiva is not recommended if you have severe kidney problems.

Elderly (over 65 years of age)

If you are elderly and taking Eslicarbazepine Zentiva on its own the dose of 1,600 mg is not a suitable dose for you

Children above 6 years of age

Dose when you start treatment

The starting dose is 10 mg per kg body weight taken once a day for one or two weeks, before increasing to the maintenance dose.

Maintenance dose

Depending on the response to Eslicarbazepine Zentiva, the dose may be increased by 10 mg per kg body weight, at intervals of one or two weeks, up to 30 mg per kg body weight. The maximum dose is 1,200 mg once daily.

Children with ≥ 60 kg

Children with 60 kg or more body weight should take the same dose as adults.

Other form of the medicine, like oral suspension, maybe more suitable for children. Ask your doctor or pharmacist.

Method and route of administration

Eslicarbazepine Zentiva is for oral use. Swallow the tablet with a glass of water. Eslicarbazepine Zentiva tablets may be taken with or without food. The tablet can be divided into equal doses.

If you take more Eslicarbazepine Zentiva than you should

If you accidentally take more Eslicarbazepine Zentiva than you should, you are potentially at risk of having more seizures; or you may feel like your heart beat is irregular or faster. Contact a doctor or go to a hospital immediately if you experience any of the above symptoms. Take the medicine pack with you. This is so the doctor knows what you have taken.

If you forget to take Eslicarbazepine Zentiva

If you forget to take a tablet, take it as soon as you remember and carry on as usual. Do not take a double dose to make up for a forgotten dose.

If you stop taking Eslicarbazepine Zentiva

Do not stop taking your tablets suddenly. If you do, you are at risk of having more seizures. Your doctor will decide how long you should take Eslicarbazepine Zentiva. Should your doctor decide to stop your treatment with Eslicarbazepine Zentiva your dose will usually be reduced gradually. It is important that your treatment is completed as advised by your doctor or your symptoms may get worse.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The following side effects can be very serious. If they happen to you stop taking Eslicarbazepine Zentiva and tell a doctor or go to a hospital immediately, as you may need urgent medical treatment:

- blistering or peeling of the skin and/or mucous membranes, rash, swallowing or breathing problems, swelling of your lips, face, eyelids, throat or tongue. These could be signs of an allergic reaction.

Very common (may affect more than 1 in 10 people) side effects are:

- Feeling dizzy or sleepy.

Common (may affect up to 1 in 10 people) side effects are:

- Feeling unsteady or having a sensation of spinning or floating;
- Feeling sick or vomiting;
- Headache;
- Diarrhoea;
- Seeing double or blurred vision;
- Difficulty in concentration;
- Feeling low in energy or tired;
- Shaking;
- Skin rash;
- Blood tests showing that you have low levels of sodium in your blood;
- Decrease of appetite;
- Difficulty in sleeping;
- Difficulty in coordinating movements (ataxia).
- Weight increase.

Uncommon (may affect up to 1 in 100 people) side effects are:

- Clumsiness;
- Allergy;
- Constipation;
- Seizures;
- Underactive thyroid gland. Symptoms include decreased level of thyroid hormone levels (seen in blood tests), cold intolerance, large tongue, thin and brittle fingernails or hair and low body temperature;
- Liver problems;
- High blood pressure or severe increase in blood pressure;
- Low blood pressure or a fall in blood pressure on standing up;
- Blood tests showing that you have low levels of salts (including chloride) in your blood or a reduction in red blood cells;
- Dehydration;
- Eye movement changes, fuzzy vision or red eye;
- Having falls;
- Thermal burn;
- Poor memory or forgetfulness;
- Crying, feeling depressed, nervous or confused, lack of interest or emotion;
- Inability to speak or write or understand spoken or written language;
- Agitation;
- Attention deficit/ hyperactivity disorder;
- Irritability;
- Mood changes or hallucinations;
- Difficulty in speaking;
- Nosebleed;
- Chest pain;
- Tingling and/or feeling numb in any part of your body;
- Migraine;
- Burning sensation;
- Abnormal sense of touch;
- Disturbances in the sense of smell;
- Ringing in the ears;
- Hearing difficulty;
- Swelling in your legs and arms;
- Heart burn, stomach upset, abdominal pain, abdominal bloating and discomfort or dry mouth;
- Charcoal (dark) stool;
- Inflamed gums or toothache;
- Sweating or having dry skin;
- Itching;
- Skin changes (e.g. red skin);
- Hair loss;
- Urinary tract infection;
- Feeling generally weak, unwell or having chills;

- Weight loss;
- Muscle pain, pain in limbs, muscular weakness;
- Bone metabolism disorder;
- Increased bone proteins;
- Flushing, cold limbs;
- Slower or irregular heart beat;
- Feeling extremely sleepy;
- Sedation;
- Neurological movement disorder where your muscles contract causing twisting and repetitive movements or abnormal postures. Symptoms include tremors, pain, cramping;
- Medicine toxicity;
- Anxiety.

Not known (frequency cannot be estimated from available data) side effects are:

- Reduction in blood platelets which increases risk of bleeding or bruising;
- Severe pain in the back and stomach (caused by inflammation of the pancreas);
- Reduction in white blood cells which makes infections more likely.
- Reddish target-like macules or circular patches often with central blisters on the trunk, skin peeling, ulcers of mouth, throat, nose, genitals and eyes, red and swollen eyes and can be preceded by fever and/or flu-like symptoms (Stevens-Johnson syndrome/toxic epidermal necrolysis);
- Initially flu-like symptoms, rash on the face then widespread rash, high body temperature, liver enzyme elevations, blood abnormalities (eosinophilia), enlarged lymph nodes and other body organs involvement (Drug Reaction with Eosinophilia and Systemic Symptoms which is also known as DRESS or drug hypersensitivity syndrome);
- Serious allergic reaction which causes swelling of the face, throat, hand, feet, ankles, or lower legs;
- Urticaria (skin rash with itching).

The use of Eslicarbazepine Zentiva is associated with an abnormality in ECG (electrocardiogram) called increase in PR interval. Side effects associated with this ECG abnormality (e.g. fainting and slowing of heart beat) may occur.

There have been reports of bone disorders including osteopenia and osteoporosis (thinning of the bone) and fractures with structurally related antiepileptics medicines like carbamazepine and oxcarbazepine. Check with your doctor or pharmacist, if you are on long-term antiepileptic treatment, have a history of osteoporosis, or take steroids.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Eslicarbazepine Zentiva

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date, which is stated on the blister, bottle and the carton after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment

6. Contents of the pack and other information

What Eslicarbazepine Zentiva contains

- Each tablet contains 800 mg of eslicarbazepine acetate
- The other ingredients are croscarmellose sodium, povidone K29/32 and magnesium stearate.

What Eslicarbazepine Zentiva looks like and contents of the pack

White to off-white tablets, oval and biconvex 19 mm in length with IC debossed on one side and a break score on one side. The tablet can be divided into equal doses.

The tablets are packaged in transparent blisters in cardboard boxes containing 10 tablets per blister, with pack sizes of 20, 30, 60 and 90 tablets, or HDPE containers with child resistant closure in cardboard boxes containing 90 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Zentiva Pharma UK Limited
12 New Fetter Lane
London EC4A 1JP
United Kingdom

Manufacturer

Coripharma ehf.
Reykjavikurvegur 78
IS-220 Hafnarfjordur
Iceland

This leaflet was last revised in March 2022

ZENTIVA

1065032271