

Package leaflet: Information for the patient
Sitagliptin/Metformin hydrochloride 50 mg/850 mg, 50 mg/1000 mg
Film-Coated Tablets
 sitagliptin/metformin hydrochloride

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist, or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. **What Sitagliptin/Metformin hydrochloride is and what it is used for**
2. **What you need to know before you take Sitagliptin/Metformin hydrochloride**
3. **How to take Sitagliptin/Metformin hydrochloride**
4. **Possible side effects**
5. **How to store Sitagliptin/Metformin hydrochloride**
6. **Contents of the pack and other information**

1. What Sitagliptin/Metformin hydrochloride is and what it is used for

Sitagliptin/Metformin hydrochloride tablet contains two different medicines called sitagliptin and metformin.

- Sitagliptin belongs to a class of medicines called DPP-4 inhibitors (dipeptidyl peptidase-4 inhibitors).
- Metformin belongs to a class of medicines called biguanides.

They work together to control blood sugar levels in adult patients with a form of diabetes called 'type 2 diabetes mellitus'. This medicine helps to increase the levels of insulin produced after a meal and lowers the amount of sugar made by your body.

Along with diet and exercise, this medicine helps lower your blood sugar. This medicine can be used alone or with certain other medicines for diabetes (insulin, sulphonylureas, or glitazones).

What is type 2 diabetes?

Type 2 diabetes is a condition in which your body does not make enough insulin, and the insulin that your body produces does not work as well as it should. Your body can also make too much sugar. When this happens, sugar (glucose) builds up in the blood. This can lead to serious medical problems like heart disease, kidney disease, blindness, and amputation.

2. What you need to know before you take Sitagliptin/Metformin hydrochloride

Do not take Sitagliptin/Metformin hydrochloride

- if you are **allergic** to sitagliptin or metformin or any of the other ingredients of this medicine (listed in section 6)
- if you have **severely reduced kidney function**
- if you have **uncontrolled diabetes**, with e.g. severe hyperglycaemia (high blood glucose), nausea, vomiting, diarrhoea, rapid weight loss, lactic acidosis (see "Risk of lactic acidosis" below) or ketoacidosis. Ketoacidosis is a condition in which substances called 'ketone bodies' accumulate in the blood and which can lead to diabetic pre-coma. Symptoms include stomach pain, fast and deep breathing, sleepiness or your breath developing an unusual fruity smell
- if you have a **severe infection** or are dehydrated
- if you are going to have an X-ray where you will be injected with a dye. You will need to stop taking Sitagliptin/Metformin hydrochloride at the time of the X-ray and for 2 or more days after as directed by your doctor, depending on how your kidneys are working
- if you have recently had a **heart attack** or have **severe circulatory problems**, such as 'shock' or breathing difficulties
- if you have **liver problems**
- if you drink alcohol to excess (either every day or only from time to time)
- if you are **breast-feeding**

Do not take Sitagliptin/Metformin hydrochloride if any of the above apply to you and talk with your doctor about other ways of managing your diabetes. If you are not sure, talk to your doctor, pharmacist or nurse before taking Sitagliptin/Metformin hydrochloride.

Warnings and precautions

Cases of inflammation of the pancreas (pancreatitis) have been reported in patients receiving Sitagliptin/Metformin hydrochloride (see section 4).

If you encounter blistering of the skin it may be a sign for a condition called bullous pemphigoid. Your doctor may ask you to stop Sitagliptin/Metformin hydrochloride.

Risk of lactic acidosis

Sitagliptin/Metformin hydrochloride may cause a very rare, but very serious side effect called lactic acidosis, particularly if your kidneys are not working properly. The risk of developing lactic acidosis is also increased with uncontrolled diabetes, serious infections, prolonged fasting or alcohol intake, dehydration (see further information below), liver problems and any medical conditions in which a part of the body has a reduced supply of oxygen (such as acute severe heart disease). If any of the above apply to you, talk to your doctor for further instructions.

Stop taking Sitagliptin/Metformin hydrochloride for a short time if you have a condition that may be associated with dehydration (significant loss of body fluids) such as severe vomiting, diarrhoea, fever, exposure to heat or if you drink less fluid than normal. Talk to your doctor for further instructions.

Stop taking Sitagliptin/Metformin hydrochloride and contact a doctor or the nearest hospital immediately if you experience some of the symptoms of lactic acidosis, as this condition may lead to coma.

Symptoms of lactic acidosis include:

- vomiting
- stomach ache (abdominal pain)
- muscle cramps
- a general feeling of not being well with severe tiredness
- difficulty in breathing
- reduced body temperature and heartbeat

Lactic acidosis is a medical emergency and must be treated in a hospital.

Talk to your doctor or pharmacist before taking Sitagliptin/Metformin hydrochloride:

- if you have or have had a **disease of the pancreas** (such as pancreatitis)
- if you have or have had **gallstones, alcohol dependence** or very **high levels of triglycerides** (a form of fat) in your blood. These medical conditions can increase your chance of getting pancreatitis (see section 4)

- if you have **type 1 diabetes**. This is sometimes called insulin-dependent diabetes
- if you have or have had an **allergic reaction** to sitagliptin, metformin, or Sitagliptin/Metformin hydrochloride (see section 4)
- if you are taking a sulphonylurea or insulin, diabetes medicines, together with Sitagliptin/Metformin hydrochloride, as you may experience low blood sugar levels (hypoglycaemia). Your doctor may reduce the dose of your sulphonylurea or insulin.

If you need to have major surgery you must stop taking Sitagliptin/Metformin hydrochloride during and for some time after the procedure.

Your doctor will decide when you must stop and when to restart your treatment with Sitagliptin/Metformin hydrochloride.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Sitagliptin/Metformin hydrochloride.

During treatment with Sitagliptin/Metformin hydrochloride, your doctor will check your kidney function at least once a year or more frequently if you are elderly and/or if you have worsening kidney function.

Children and adolescents

Children and adolescents below 18 years should not use this medicine. It is not effective in children and adolescents between the ages of 10 and 17 years. It is not known if this medicine is safe and effective when used in children younger than 10 years.

Other medicines and Sitagliptin/Metformin hydrochloride

If you need to have an injection of a contrast medium that contains iodine into your bloodstream, for example, in the context of an X-ray or scan, you must stop taking Sitagliptin/Metformin hydrochloride before or at the time of the injection. Your doctor will decide when you must stop and when to restart your treatment with Sitagliptin/Metformin hydrochloride.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. You may need more frequent blood glucose and kidney function tests, or your doctor may need to adjust the dosage of Sitagliptin/Metformin hydrochloride. It is especially important to mention the following:

- medicines (taken by mouth, inhalation, or injection) used to treat diseases that involve inflammation, like asthma and arthritis (corticosteroids)
- medicines which increase **urine production** (diuretics)
- medicines used to treat **pain and inflammation** (NSAID and COX-2-inhibitors, such as ibuprofen and celecoxib)
- certain medicines for the treatment of **high blood pressure** (ACE inhibitors and angiotensin II receptor antagonists)
- specific medicines for the treatment of **bronchial asthma** (β -sympathomimetics)
- **iodinated contrast agents or alcohol-containing medicines**
- certain medicines used to treat **stomach problems** such as cimetidine
- ranolazine, a medicine used to treat **angina**
- dolutegravir, a medicine used to treat **HIV infection**
- **vandetanib**, a medicine used to treat a specific type of thyroid cancer (medullary thyroid cancer)
- **digoxin** (to treat irregular heart beat and other heart problems). The level of digoxin in your blood may need to be checked if taking with Sitagliptin/Metformin hydrochloride.

Sitagliptin/Metformin hydrochloride with alcohol

Avoid excessive alcohol intake while taking Sitagliptin/Metformin hydrochloride since this may increase the risk of lactic acidosis (see section "Warnings and precautions").

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, speak to your doctor in case any changes will be needed to your treatment or monitoring of your blood glucose levels. This medicine is not recommended if you are breast-feeding or if you are planning to breast-feed your baby. See section 2, **Do not take Sitagliptin/Metformin hydrochloride**.

Driving and using machines

This medicine has no or negligible influence on the ability to drive and use machines. However, dizziness and drowsiness have been reported with sitagliptin, which may affect your ability to drive or use machines.

Taking this medicine in combination with medicines called sulphonylureas or with insulin can cause hypoglycaemia, which may affect your ability to drive and use machines or work without safe foothold.

Sitagliptin/Metformin hydrochloride Tablets contain sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Sitagliptin/Metformin hydrochloride

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The usual recommended dose is:

- Take one tablet:
 - twice daily by mouth
 - with meals to lower your chance of an upset stomach.
- Your doctor may need to increase your dose to control your blood sugar.
- If you have reduced kidney function, your doctor may prescribe a lower dose.

You should continue the diet recommended by your doctor during treatment with this medicine and take care that your carbohydrate intake is equally distributed over the day.

This medicine alone is unlikely to cause abnormally low blood sugar (hypoglycaemia). When this medicine is used with a sulphonylurea medicine or with insulin, low blood sugar can occur and your doctor may reduce the dose of your sulphonylurea or insulin.

Artwork	Dr.Reddy's
Version No:	v2.3
Submission	
Brand:	DR Reddy's
Country:	UK
Product Name:	Sitagliptin Metformin
Strength:	Various
Form:	Film-coated Tablets
Component:	Leaflet
Pack Size:	Various
Date Created:	17 FEB 2021
Date Modified:	23 June 2023
Project:	CC1000101915
Previous DR No:	DR000701P-3C
Previous Material Number:	150090726
Commercial	
DR No:	DR000XXX
DRUGS Code:	N/A1500XXXXX
Material Code:	1500XXXXX
Third Party Material Code:	N/A
Barcode Type:	N/A
Barcode Number:	N/A
Magnification:	
Pharmacode No:	XXX
Implementation Date:	N/A
Technical Information	
Die Cut	Guides
Min Font Size:	9 pt
Printer:	India
Dimensions:	150 x 600 mm
Colours	
Process Black	
<p align="center">Dr.Reddy's Good Health Can't Wait Dr. Reddy's Laboratories (UK) Ltd, 410 Cambridge Science Park, Milton Road, Cambridge, CB4 0PE, United Kingdom</p>	
<p align="center">UNDER NO CIRCUMSTANCES SHOULD THIS ARTWORK BE ALTERED WITHOUT PRIOR PERMISSION FROM DR.REDDY'S ARTWORK EU.</p>	
<p align="center"><small>Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.</small></p>	

150 mm

37.5 mm

37.5 mm

37.5 mm

37.5 mm

If you take more Sitagliptin/Metformin hydrochloride than you should

If you take more than the prescribed dosage of this medicine, contact your doctor immediately. Go to the hospital if you have symptoms of lactic acidosis such as feeling cold or uncomfortable, severe nausea or vomiting, stomach ache, unexplained weight loss, muscular cramps, or rapid breathing (see section "Warnings and precautions").

If you forget to take Sitagliptin/Metformin hydrochloride

If you miss a dose, take it as soon as you remember. If you do not remember until it is time for your next dose, skip the missed dose and go back to your regular schedule. Do not take a double dose of this medicine.

If you stop taking Sitagliptin/Metformin hydrochloride

Continue to take this medicine as long as your doctor prescribes it so you can continue to help control your blood sugar. You should not stop taking this medicine without talking to your doctor first. If you stop taking Sitagliptin/Metformin hydrochloride, your blood sugar may rise again.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

STOP taking Sitagliptin/Metformin hydrochloride and contact a doctor immediately if you notice any of the following serious side effects:

- severe and persistent pain in the abdomen (stomach area) which might reach through to your back with or without nausea and vomiting, as these could be signs of an inflamed pancreas (pancreatitis)

Sitagliptin/Metformin hydrochloride may cause a very rare (may affect up to 1 in 10,000 people), but very serious side effect called lactic acidosis (see section "Warnings and precautions"). If this happens, you must stop taking Sitagliptin/Metformin hydrochloride and contact a doctor or the nearest hospital immediately, as lactic acidosis may lead to coma.

If you have a serious allergic reaction (frequency not known), including rash, hives, blisters on the skin/peeling skin and swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing, stop taking this medicine and call your doctor right away. Your doctor may prescribe a medicine to treat your allergic reaction and a different medicine for your diabetes.

Some patients taking metformin have experienced the following side effects after starting sitagliptin: Common (may affect up to 1 in 10 people): low blood sugar, nausea, flatulence, vomiting Uncommon (may affect up to 1 in 100 people): stomach ache, diarrhoea, constipation, drowsiness

Some patients have experienced diarrhoea, nausea, flatulence, constipation, stomach ache or vomiting when starting the combination of sitagliptin and metformin together (frequency is common).

Some patients have experienced the following side effects while taking this medicine with a sulphonylurea such as gliclazide: Very common (may affect more than 1 in 10 people): low blood sugar Common: constipation

Some patients have experienced the following side effects while taking this medicine in combination with pioglitazone: Common: swelling of the hands or legs

Some patients have experienced the following side effects while taking this medicine in combination with insulin: Very common: low blood sugar Uncommon: dry mouth, headache

Some patients have experienced the following side effects during clinical studies while taking sitagliptin alone (one of the medicines in Sitagliptin/Metformin hydrochloride) or during post-approval use of Sitagliptin/Metformin hydrochloride or sitagliptin alone or with other diabetes medicines: Common: low blood sugar, headache, upper respiratory infection, stuffy or runny nose and sore throat, osteoarthritis, arm or leg pain Uncommon: dizziness, constipation, itching Rare: reduced number of platelets Frequency not known: kidney problems (sometimes requiring dialysis), vomiting, joint pain, muscle pain, back pain, interstitial lung disease, bullous pemphigoid (a type of skin blister)

Some patients have experienced the following side effects while taking metformin alone: Very common: nausea, vomiting, diarrhoea, stomach ache and loss of appetite. These symptoms may happen when you start taking metformin and usually go away Common: a metallic taste, decreased or low vitamin B12 levels in the blood (symptoms may include extreme tiredness (fatigue), a sore and red tongue (glossitis), pins and needles (paraesthesia) or pale or yellow skin). Your doctor may arrange some tests to find out the cause of your symptoms because some of these may also be caused by diabetes or due to other unrelated health problems.

Very rare: hepatitis (a problem with your liver), hives, redness of the skin (rash) or itching

Reporting of side effects If you get any side effects, talk to your doctor, pharmacist, or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme website www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Sitagliptin/Metformin hydrochloride Keep this medicine out of the sight and reach of children. This medicinal product does not require any special storage conditions Do not use this medicine after the expiry date which is stated on the blister and the carton after 'EXP'. The expiry date refers to the last day of the month. This medicinal product does not require any special storage conditions. Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Sitagliptin/Metformin hydrochloride Tablet contains

- The active substances are sitagliptin and metformin.
- For Sitagliptin/Metformin hydrochloride 50 mg/850 mg Tablets: Each film-coated tablet contains sitagliptin hydrochloride monohydrate equivalent to 50 mg of sitagliptin and 850 mg of metformin hydrochloride.
- For Sitagliptin/Metformin hydrochloride 50 mg/1000 mg Tablets: Each film-coated tablet contains sitagliptin hydrochloride monohydrate equivalent to 50 mg of sitagliptin and 1000 mg of metformin hydrochloride.
- The other ingredients are: In the tablet core: povidone K 30; sodium lauryl sulfate; microcrystalline cellulose, magnesium stearate. In the film coating: poly(vinyl alcohol), macrogol 3350, talc, titanium dioxide (E171). For 50mg/1000 mg only: iron oxide red (E172) and iron oxide yellow (E172).

What Sitagliptin/Metformin hydrochloride Tablets look like and contents of the pack

Sitagliptin/Metformin hydrochloride 50 mg/850 mg film-coated Tablets are white coloured, biconvex, oval tablets with "1" debossed on one side and plain on the other side. The approximate tablet dimensions: 20.1 mm x 9.8 mm x 6.4 mm

Sitagliptin/Metformin hydrochloride 50 mg/1000 mg film-coated Tablets are brown colored, biconvex, oval tablets with "7" debossed on one side and plain on other side. The approximate tablet dimensions: 21.1 mm x 10.3 mm x 6.7 mm

OPA/Alu/PVC-Alu blister. Pack of 28, 56, 112 or 196 film-coated tablets. Not all pack sizes may be marketed.

6. Contents of the pack and other information

What Sitagliptin/Metformin hydrochloride Tablet contains

- The active substances are sitagliptin and metformin.
- For Sitagliptin/Metformin hydrochloride 50 mg/850 mg Tablets: Each film-coated tablet contains sitagliptin hydrochloride monohydrate equivalent to 50 mg of sitagliptin and 850 mg of metformin hydrochloride.
- For Sitagliptin/Metformin hydrochloride 50 mg/1000 mg Tablets: Each film-coated tablet contains sitagliptin hydrochloride monohydrate equivalent to 50 mg of sitagliptin and 1000 mg of metformin hydrochloride.
- The other ingredients are: In the tablet core: povidone K 30; sodium lauryl sulfate; microcrystalline cellulose, magnesium stearate. In the film coating: poly(vinyl alcohol), macrogol 3350, talc, titanium dioxide (E171). For 50mg/1000 mg only: iron oxide red (E172) and iron oxide yellow (E172).

What Sitagliptin/Metformin hydrochloride Tablets look like and contents of the pack

Sitagliptin/Metformin hydrochloride 50 mg/850 mg film-coated Tablets are white coloured, biconvex, oval tablets with "1" debossed on one side and plain on the other side. The approximate tablet dimensions: 20.1 mm x 9.8 mm x 6.4 mm

Sitagliptin/Metformin hydrochloride 50 mg/1000 mg film-coated Tablets are brown colored, biconvex, oval tablets with "7" debossed on one side and plain on other side. The approximate tablet dimensions: 21.1 mm x 10.3 mm x 6.7 mm

OPA/Alu/PVC-Alu blister. Pack of 28, 56, 112 or 196 film-coated tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Dr. Reddy's Laboratories (UK) Ltd, 410 Cambridge Science Park, Milton Road, Cambridge, CB4 0PE, United Kingdom

This leaflet was last revised in March 2023.

600 mm