

Track & Tie Printers

02-03-2022

09-06-2022

13-06-2022

16-06-2022

24-06-2022

19-08-2022

145.00 mm

297.50 mm

PATIENT INFORMATION LEAFLET

ETHAMBUTOL 100 mg TABLETS

(Ethambutol Hydrochloride)

Read all of this leaflet carefully before you start taking this medicine.

Keep this leaflet. You may need to read it again. If you have any further questions, ask your doctor or pharmacist. This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours. If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

In this leaflet:

1. What Ethambutol Tablets are and what they are used for
2. Before you take Ethambutol Tablets
3. How to take Ethambutol Tablets
4. Possible side effects
5. How to store Ethambutol Tablets
6. Further information

1. WHAT ETHAMBUTOL TABLETS ARE AND WHAT THEY ARE USED FOR

The name of your medicine is Ethambutol Tablets. Ethambutol belongs to a group of medicines called antituberculosis drugs and is used for the treatment and prevention of tuberculosis, an infectious disease mainly affecting the lungs.

2. BEFORE YOU TAKE ETHAMBUTOL TABLETS

Do not take Ethambutol Tablets:

- if you have sight problems.
- if you are allergic (hypersensitive) to ethambutol hydrochloride or any of the other ingredients (see list under 'What Ethambutol Tablets contain' in section 6).

Problems with your sight

You should have your eyesight checked before taking these tablets. If you notice any changes to your vision whilst taking these tablets, **you should stop taking them and notify your doctor immediately.** In the majority of cases, any changes to vision will return to normal within a few weeks/months after stopping treatment, however, in rare cases, it may take longer for eyesight to recover or the changes may become permanent.

Take special care with Ethambutol if you have a history of kidney problems. Before treatment with Ethambutol you should have your kidneys checked.

If you develop symptoms suggestive of hepatitis (e.g. abdominal pain, fever, loss of appetite) or feel generally unwell, you may need your liver functions tested.

Taking other medicines

You should not take ethambutol tablets at the same time as antacid medicine containing aluminium hydroxide (used to treat heartburn and indigestion) as this can make ethambutol less effective. Please tell your doctor or pharmacist if you are taking or have recently take any other medicines, including medicines obtained without a prescription.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

Driving and using machines

Ethambutol occasionally causes sight problems and may present symptoms which affect the ability to drive or use machinery (e.g. numbness or paraesthesia (abnormal sensation like pins and needles) of the furthest parts of your body, disorientation). After taking your medicine you should not drive or use machinery until you know how ethambutol affects you.

3. HOW TO TAKE ETHAMBUTOL TABLETS

Always take Ethambutol Tablets exactly as your doctor has told you to. You should check with your doctor or pharmacist if you are not sure.

The dosage of ethambutol varies from person to person depending on your age, how much you weigh and whether it is being used for the treatment or for the prevention of tuberculosis.

Patients who have kidney problems may need blood tests so that their doctor can reduce the dose in some cases.

Ethambutol Tablets should be taken once per day. They should be swallowed whole with a drink of water; do not chew or crush the tablets.

If you are not sure how many tablets to take, or when to take them, ask your pharmacist.

Keep taking your tablets for as long as your doctor directed, even if you feel better.

Adults:

Prevention and first time treatment: The usual dose of ethambutol for adults for prevention of tuberculosis or for first time treatment of tuberculosis is 15mg per kg of body weight per day.

Second time (or subsequent) treatment: The usual dose of ethambutol for adults for second time (or subsequent) treatment of tuberculosis is 25mg per kg of body weight per day for the first 60 days, reducing to 15mg per kg of body weight per day for as long as the doctor considers necessary.

Children:

Prevention: The usual dose of ethambutol for children for prevention of tuberculosis is 15mg per kg of body weight per day.

First, second time (or subsequent) treatment: The usual dose of ethambutol for children for treatment of tuberculosis is 25mg per kg of body weight per day for the first 60 days, reducing to 15mg per kg of body weight per day for as long as the doctor considers necessary.

Track & Tie Printers

02-03-2022

09-06-2022

13-06-2022

16-06-2022

24-06-2022

19-08-2022

If you take more Ethambutol Tablets than you should

If you have taken an overdose of Ethambutol Tablets (that is more than the doctor has told you to) get medical help immediately, either by calling your doctor or going to the nearest hospital casualty department. Remember to take the labelled medicine bottle with you, whether there are any Ethambutol Tablets left or not.

If you forget to take Ethambutol Tablets

If you forget to take a tablet you should take it as soon as you remember. However, if this is within 2 hours of your next dose you should skip the missed tablet and carry on taking the rest of your tablets as usual. Do not take a double dose to make up for a forgotten tablet.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If you get any of the following symptoms, tell your doctor immediately:

- Severe allergic reaction (e.g. difficulty in breathing, flushed appearance, agitation or an irregular heartbeat.) **If this occurs, stop taking this medicine and contact your doctor immediately.**
- Eye problems (e.g. colour blindness, loss of vision or visual acuity). **If you notice any changes or problems with your sight, contact your doctor immediately.**
- Stevens-Johnson syndrome (a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and/or genitals)
- Kidney problems (lower back pain, pain or burning when passing urine, difficulty passing urine)
- Liver problems (which may result in pale stools, blood in your urine or yellowing of the skin and/or eyes).

Other possible side effects include:

Uncommon side effects (may affect up to 1 in 100 people)

- High levels of uric acid in the blood (hyperuricaemia)

Rare side effects (may affect up to 1 in 1,000 people)

- Thrombocytopenia (which may result in bruising, nosebleeds, pinpoint red spots, prolonged bleeding after injury)
- Minor allergic reactions such as rashes, itching and hives
- Muscle weakness and wasting, pain, numbness, pins and needles sensations

Very rare side effects (may affect up to 1 in 10,000 people)

- Blood disorders such as leucopenia and neutropenia
- Gout (a symptom of these conditions is joint inflammation), joint pains
- Headache, dizziness, disorientation
- Mental confusion, hallucinations
- Shortness of breath, chest pains, generally feeling unwell, fever
- Liver failure
- Blisters or peeling skin (inc. following exposure to sunlight)

Not known side effects (frequency cannot be estimated from the available data)

- Loss of appetite, feeling sick, being sick, stomach pain/upsets, diarrhoea

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any side effects not listed in this leaflet.

You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for 'MHRA Yellow Card' in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE ETHAMBUTOL TABLETS

Keep out of the sight and reach of children.

Do not use Ethambutol Tablets after the expiry date which is stated on the packaging after EXP.

Store below 30°C in the original packaging to protect from moisture.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.

6. FURTHER INFORMATION

What Ethambutol Tablets Contain:

The active ingredient is ethambutol (as ethambutol hydrochloride). Each tablet contains 100mg of ethambutol.

Other ingredients are:

Tablet core: Microcrystalline cellulose, sodium starch glycolate, maize starch, povidone, colloidal anhydrous silica and magnesium stearate.

Tablet coating: Polydextrose, hydroxypropylmethylcellulose, polyethylene glycol 4000, purified water, titanium dioxide (E171), iron oxide yellow (E172).

What Ethambutol Tablets look like and contents of the pack:

Ethambutol 100mg Tablets are round yellow coated tablets, plain on both sides. The tablets are available in bottles of 56 tablets.

Marketing Authorisation Holder

Esteve Pharmaceuticals Ltd, The Courtyard Barns, Choke Lane, Maidenhead, Berks SL6 6PT, UK.

Manufacturer:

Esteve Pharmaceuticals Ltd, The Courtyard Barns, Choke Lane, Maidenhead, Berks SL6 6PT, UK.

This leaflet was last revised in August 2022.

PMS-1211/01

ARTWORK APPROVAL
HISTORY APPROVAL
NUMBER:

REGULATORY

PACKAGING

QUALITY

DATE OF FINAL APPROVAL:

Artwork No. PMS-1211/01
Version No. 01
Version Tracks. 01
Product. Ethambutol 100mg PIL
Artwork Type: 56 Tablets
Dimension: 145x297.5mm

DATE: 10-06-2022
COLOURS: Process Black 