

<div>Other information</div> <div>Patient's Name:</div> <div>Doctor's Name:</div> <div>Doctor's Phone:</div>	<div><div>This patient alert card contains important safety information that you need to be aware of before and during treatment with Simponi.</div><div><div><div></div><div>32640B</div></div><div><div>Simponi<sup>®</sup></div><div>golimumab</div></div><div></div></div><div>Simponi Patient Alert Card</div></div> <div><div>3</div><div>Dates of Simponi treatment:</div><div>1st administration:</div><div><div>/</div><div>/</div><div>/</div></div><div>Subsequent administrations:</div><div><div>/</div><div>/</div><div>/</div></div><div><div>/</div><div>/</div><div>/</div></div><div><div>/</div><div>/</div><div>/</div></div></div> <div><div>4</div><div>Other information</div><div><div><div><div></div><div>Please make sure you also have a list of all other medicines that you are using with you at any visit to a health care professional.</div></div><div><div></div><div>Keep this card with you for 6 months after the last Simponi dose, since side effects may occur a long time after your last dose of Simponi.</div></div><div><div></div><div>Read the Simponi package leaflet carefully before you start using this medicine.</div></div></div></div></div> <div><div>1</div><div>Infections</div><div><div>When you are treated with Simponi, you might get infections more easily. Infections may progress more rapidly and may be more severe. In addition, some previous infections may reappear.</div><div><div>1.1 Prior to Simponi treatment:</div><div><div></div><div>Tell your doctor if you have an infection. You must not be treated with Simponi if you have tuberculosis (TB) or any other severe infection.</div></div></div></div><div><div>2</div><div>Heart failure</div><div><div><div>2.1 Prior to Simponi treatment:</div><div><div></div><div>Tell your doctor if you have a history of heart failure. You must not be treated with Simponi if you have moderate to severe heart failure.</div></div><div><div>2.2 During and after Simponi treatment:</div><div><div></div><div>If you develop symptoms of heart failure (e.g. shortness of breath or swelling of the feet) seek medical attention immediately.</div></div></div><div><div>Ask your doctor to record the type and date of the last screening(s) for TB below:</div><div><div><div>Test</div><div>Date</div><div>/</div><div>/</div></div><div><div>Test</div><div>Date</div><div>/</div><div>/</div></div></div><div><div><div></div><div>Tell your doctor if you know or suspect you are a carrier of the hepatitis B virus.</div></div><div><div><div>1.2 During and after Simponi treatment:</div><div><div></div><div>Seek medical attention immediately if you develop symptoms of an infection, such as fever, tiredness, (persistent) cough, shortness of breath, or flu-like signs, weight loss, night sweats, diarrhoea, wounds, dental problems and a burning feeling when urinating.</div></div></div></div></div><div><div>SIDE 1/2</div><div>FRONT</div></div><div><div>SIDE 2/2</div><div>(horizontal flipped)</div><div>BACK</div></div></div></div><div data-bbox="36 1400 414 1419" data-label="Page-Footer"><p>32640B.pdf - Page 1 of 1 - March 10, 2010 - 14:32:21</p></div></div></div></div>
--	---