



Package leaflet: Information for the patient

Co-amoxiclav 250 mg /125 mg film-coated tablets

amoxicillin/clavulanic acid

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Co-amoxiclav 250/125 mg is and what it is used for
2. What you need to know before you take Co-amoxiclav 250/125 mg
3. How to take Co-amoxiclav 250/125 mg
4. Possible side effects
5. How to store Co-amoxiclav 250/125 mg
6. Contents of the pack and other information

1. What Co-amoxiclav is and what it is used for

Co-amoxiclav is an antibiotic and works by killing bacteria that cause infections. It contains two different medicines called amoxicillin and clavulanic acid. Amoxicillin belongs to a group of medicines called "penicillins" that can sometimes be stopped from working (made inactive). The other active component (clavulanic acid) stops this from happening.

Co-amoxiclav 250/125 mg is used in adults and children to treat the following infections:

- sinus infections
- urinary tract infections
- skin infections
- dental infections.

2. What you need to know before you take Co-amoxiclav 250/125 mg

Do not take Co-amoxiclav :

- if you are allergic to amoxicillin, clavulanic acid, penicillin or any of the other ingredients of this medicine (listed in section 6)
- if you have ever had a severe allergic (hypersensitive) reaction to any other antibiotic. This can include a skin rash or swelling of the face or throat
- if you have ever had liver problems or jaundice (yellowing of the skin) when taking an antibiotic.

→ **Do not take Co-amoxiclav if any of the above apply to you.** If you are not sure, talk to your doctor or pharmacist before taking Co-amoxiclav.

Warnings and precautions

Talk to your doctor or pharmacist before taking Co-amoxiclav if you:

- have glandular fever
- are being treated for liver or kidney problems
- are not passing water regularly.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Co-amoxiclav.

In some cases, your doctor may investigate the type of bacteria that is causing your infection. Depending on the results, you may be given a different strength of Co-amoxiclav or a different medicine.

Conditions you need to look out for

Co-amoxiclav can make some existing conditions worse, or cause serious side effects. These include allergic reactions, convulsions (fits) and inflammation of the large intestine. You must look out for certain symptoms while you are taking Co-amoxiclav, to reduce the risk of any problems. See 'Conditions you need to look out for' in **section 4**.

Blood and urine tests

If you are having blood tests (such as red blood cell status tests or liver function tests) or urine tests (for glucose), let the doctor or nurse know that you are taking Co-amoxiclav. This is because Co-amoxiclav can affect the results of these types of

tests.

Other medicines and Co-amoxiclav

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines.

- If you are taking allopurinol (used for gout) with Co-amoxiclav, it may be more likely that you will have an allergic skin reaction.
- If you are taking probenecid (used for gout), your doctor may decide to adjust your dose of Co-amoxiclav.
- If medicines to help stop blood clots (such as warfarin) are taken with Co-amoxiclav then extra blood tests may be needed.
- Co-amoxiclav can affect how methotrexate (a medicine used to treat cancer or rheumatic diseases) works.
- Co-amoxiclav may affect how mycophenolate mofetil (a medicine used to prevent the rejection of transplanted organs) works.

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

Co-amoxiclav can have side effects and the symptoms may make you unfit to drive. Do not drive or operate machinery unless you are feeling well.

Co-amoxiclav contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Co-amoxiclav 250/125 mg tablets

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Adults and children weighing 40 kg and over

The usual dose is:

- 1 tablet three times a day

Children weighing less than 40 kg

Children aged 6 years or less should preferably be treated with Co-amoxiclav oral suspension or sachets. Co-amoxiclav tablets are not recommended.

Patients with kidney and liver problems

- If you have kidney problems the dose might be changed. A different strength or different medicine may be chosen by your doctor.
- If you have liver problems you may have more frequent blood tests to check your liver is working.

How to take Co-amoxiclav 250/125 mg tablets

- Take with a meal.
- Swallow the tablets whole with a glass of water.
- Space the doses evenly during the day, at least 4 hours apart. Do not take 2 doses in 1 hour.
- Do not take Co-amoxiclav for more than 2 weeks. If you still feel unwell you should go back to see the doctor.

If you take more Co-amoxiclav than you should

If you take too much Co-amoxiclav, signs might include an upset stomach (feeling sick, being sick or diarrhoea) or convulsions. Talk to your doctor as soon as possible. Take the medicine carton to show the doctor.

If you forget to take Co-amoxiclav

- If you forget to take a dose take it as soon as you remember.
- You should not take the next dose too soon, but wait around 4 hours before taking the next dose. Do not take a double dose to make up for a forgotten dose.

If you stop taking Co-amoxiclav

Keep taking Co-amoxiclav until the treatment is finished, even if you feel better. You need every dose to help fight the infection. If some bacteria survive they can cause the infection



to come back.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Conditions you need to look out for

Allergic reactions:

- skin rash
- inflammation of blood vessels (*vasculitis*) which may be visible as red or purple raised spots on the skin, but can affect other parts of the body
- fever, joint pain, swollen glands in the neck, armpit or groin
- swelling, sometimes of the face or throat (*angioedema*), causing difficulty in breathing
- collapse
- chest pain in the context of allergic reactions, which may be a symptom of allergy triggered cardiac infarction (Kounis syndrome)

→ **Contact a doctor immediately if you get any of these symptoms. Stop taking Co-amoxiclav.**

Inflammation of large intestine

Inflammation of the large intestine, causing watery diarrhoea usually with blood and mucus, stomach pain and/or fever.

Acute inflammation of the pancreas (acute pancreatitis)

If you have severe and on-going pain in the stomach area this could be a sign of acute pancreatitis.

Drug-induced enterocolitis syndrome (DIES):

DIES has been reported mainly in children receiving amoxicillin/clavulanate. It is a certain kind of allergic reaction with the leading symptom of repetitive vomiting (1-4 hours after drug intake). Further symptoms could comprise abdominal pain, lethargy, diarrhoea and low blood pressure.

→ **Contact your doctor as soon as possible for advice if you get these symptoms.**

Very common side effects

These may affect more than 1 in 10 people

- diarrhoea (in adults).

Common side effects

These may affect up to 1 in 10 people

- thrush (*candida* - a yeast infection of the vagina, mouth or skin folds)
- feeling sick (nausea), especially when taking high doses
 - if affected take Co-amoxiclav before food
- vomiting
- diarrhoea (in children).

Uncommon side effects

These may affect up to 1 in 100 people

- skin rash, itching
- raised itchy rash (*hives*)
- indigestion
- dizziness
- headache.

Uncommon side effects that may show up in your blood tests:

- increase in some substances (*enzymes*) produced by the liver.

Rare side effects

These may affect up to 1 in 1000 people

- skin rash, which may blister, and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge - *erythema multiforme*)

→ if you notice any of these symptoms contact a doctor urgently.

Rare side effects that may show up in your blood tests:

- low number of cells involved in blood clotting
- low number of white blood cells.

Frequency not known

Frequency cannot be estimated from the available data.

- allergic reactions (see above)
- inflammation of the large intestine (see above)
- inflammation of the protective membrane surrounding the brain (*aseptic meningitis*)
- serious skin reactions:
 - a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (*Stevens-Johnson syndrome*), and a more severe form, causing extensive peeling of the skin (more than 30% of the body surface - *toxic epidermal necrolysis*)
 - widespread red skin rash with small pus-containing blisters (*bullous exfoliative dermatitis*)

- a red, scaly rash with bumps under the skin and blisters (*exanthemous pustulosis*).
- flu-like symptoms with a rash, fever, swollen glands, and abnormal blood test results (including increased white blood cells (eosinophilia) and liver enzymes) (Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS))
- rash with blisters arranged in a circle with central crusting or like a string of pearls (linear IgA disease)
- inflammation of the membranes that surround the brain and spinal cord (aseptic meningitis)

Contact a doctor immediately if you get any of these symptoms.

- inflammation of the liver (hepatitis)
- jaundice, caused by increases in the blood of bilirubin (a substance produced in the liver) which may make your skin and whites of the eyes appear yellow
- inflammation of tubes in the kidney
- blood takes longer to clot
- hyperactivity
- convulsions (in people taking high doses of Coamoxiclav or who have kidney problems)
- black tongue which looks hairy

Side effects that may show up in your blood or urine tests:

- severe reduction in the number of white blood cells
- low number of red blood cells (*haemolytic anaemia*)
- crystals in urine leading to acute kidney injury.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via Yellow Card Scheme, Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Co-amoxiclav

Keep this medicine out of the sight and reach of children.

Store in the original blister in order to protect from light and moisture.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month. Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Co-amoxiclav contains

- The active substances are amoxicillin trihydrate and potassium Clavulanate. Each film-coated tablet contains amoxicillin trihydrate equivalent to 250 mg amoxicillin and potassium clavulanate equivalent to 125 mg clavulanic acid.
- The other ingredients are:
 - Tablet core:* microcrystalline cellulose, crospovidone (Type B), sodium starch glycolate (Type A), ethyl cellulose dispersion (Type B), anhydrous colloidal silica, magnesium stearate.
 - Tablet coat:* hypromellose 2910 (5cps), hypromellose 2910 (15cps), macrogol 4000, macrogol 6000, titanium dioxide (E171).

What Co-amoxiclav looks like and contents of the pack

Co-amoxiclav 250 mg /125 mg tablets are white to off-white, oval shaped, film coated tablets, debossed with 'A' on one side and '63' on the other side. The size is 18.1 mm × 8.6 mm.

The tablets are packaged in blisters. Each pack contains 21 film-coated tablets.

Marketing Authorisation Holder

Milpharm Limited
Ares Block, Odyssey Business Park
West End Road
Ruislip HA4 6QD
United Kingdom

Manufacturer

APL Swift Services (Malta) Limited
HF26, Hal Far Industrial Estate,
Hal Far Birzebbugia, BBG 3000 Malta
or
Milpharm Limited
Ares Block, Odyssey Business Park
West End Road
Ruislip HA4 6QD
United Kingdom

This leaflet was last revised in 02/2023.

The package information leaflet in formats such as Braille, audio, CD-rom or large print is made available on request from patients to the organisations.