
PRODUCT TITLE



IMPORTANT: Peel this label at corner and read BEFORE use. Do not tear off, re-fix for future use.

For symptomatic relief of dry or irritating coughs.

Take the solution by mouth.

Adults, the elderly and children over 12 years: Take 1 or 2 x 5ml spoonfuls. Repeat after 4 hours if required, but not more than 4 doses in 24 hours.

Do not give to children under 12 years old. Do not take with other cough and cold medicines

Warning: Do not take more medicine than the label tells you to.

If symptoms persist consult your doctor. Peel where shown for further precautions.

Active ingredient per 5ml:

pholcodine monohydrate 5mg. Also contains amongst other

ingredients: ethanol, colours (E104, E110, E123), sodium benzoate (E211), sucrose 4g per 5ml, alcohol 6.5 vol%

Peel where shown for full list of ingredients.

Do not store above 25°C.

Keep out of the sight and reach of children.

Manufactured by the licence holder L.C.M. Ltd., Huddersfield, HD7 5QH, UK for Thornton and Ross Ltd., Huddersfield, HD7 5QH, UK.

PL 12965/0030 92774872002

200ml **e** Р





Peel

This medicine contains pholcodine monohydrate, which reduces the desire to cough.

1. Before you use this medicine ☑ Do not use the medicine if you have....

- An allergy to any of the ingredients listed in section 3
- Liver disease.
- Shallow breath or breathing difficulties.
 Chest problems or chest infections for a long time.
- An asthma attack.
- Taken MAOIs (monoamine oxidase inhibitors) for depression within the last two weeks.
- Or in children under 12 years old.

▲ Talk to your doctor or pharmacist if you have..

- Kidney or liver problems.
- · A history of asthma or other breathing problems including bronchitis.
- A persistent cough or one that produces phlegm.
- A history of drug abuse; pholcodine is an opioid and addiction is observed with opioids as a class.

92774872002

▲ Talk to your doctor or pharmacist if you are taking.

- Medicine for high blood pressure, e.g. ACE inhibitors, beta blockers.
- . Diuretics, which help remove water from the body, e.g. furosemide, hydrochlorothiazide.
- · Medicines used to treat sleeplessness, anxiety and schizophrenia.
- · Alcohol or tranquillisers.
- · Pain killers, e.g. morphine.
- Or are having surgery, as muscle relaxants used may react with this medicine.

Concomitant use of this product and sedative medicines such as benzodiazepines or related drugs increases the risk of drowsiness, difficulties in breathing (respiratory depression), coma and may be life-threatening. Because of this, concomitant use should only be considered when other treatment options are not possible.

However if your doctor does prescribe this product together with sedative medicines the dose and duration of concomitant treatment should be limited by your doctor.

Please tell your doctor about all sedative medicines you are taking, and follow your doctor's dose recommendation closely. It could be helpful to inform friends or relatives to be aware of the signs and symptoms stated above. Contact your doctor when experiencing such symptoms.

♠Pregnant or breastfeeding....

Ask your doctor or pharmacist for advice before using this medicine if you are pregnant, might be pregnant or are breastfeeding. Pholoodine Linctus should not be used in pregnancy unless the doctor has told you to do so.

♠Driving and using machines....
This medicine may cause drowsiness. If you are affected do not drive or operate machinery.

A Important ingredient information

This medicine contains:

• 270mg of alcohol (ethanol) in each 5ml dose. The amount in 5ml of this medicine is equivalent to less than 7ml of beer or 3ml of wine. The small amount of alcohol in this medicine will not have any noticeable effects

- 4g of sucrose per 5ml dose. To be taken into account in people with diabetes mellitus. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.
 • amaranth (E123) and sunset yellow
- (E110) which may cause allergic reactions.
 2.5mg sodium benzoate in each 5ml dose.
- less than 1mmol sodium (23mg) per 5ml dose, that is to say essentially 'sodium-free'.

2. Possible side effects

Like all medicines, Pholcodine Linctus can have side effects, although these don't affect everyone. Important side effects: If you think you have any of the following side effects, stop using this medicine immediately and see a doctor as soon as possible.

· Allergic reactions which cause difficulty in breathing, fever, swelling of the mouth, lips or skin, severe rashes and itching.

Other possible side effects are: • Feeling or being sick • Constipation • Dizziness • Drowsiness • Feeling excited or confused . Rash . Failure to cough up phleam.

If you notice these or any other side effects, stop use and tell your doctor or pharmacist. They will tell you what to do.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

3. Further information

Alf you accidentally take too much see a doctor straight away. Take the pack with you to show which medicine you have swallowed.

- Do not use after the expiry date. The expiry date refers to the last day of that month.
- Return any unused medicine to the pharmacy for

The active ingredient per 5ml is: pholcodine monohydrate 5mg

The other ingredients are: citric acid (E330), ethanol, quinoline yellow (E104), sunset yellow (E110), amaranth (E123), sodium benzoate (E211), syrup (sucrose) and purified water.
What the medicine looks like

Pholoodine Linctus is an orange syrupy liquid. It is supplied in 200ml bottles.

This label was last revised in February 2020

92774872002