


140mm
125mm
3mm
5mm
150mm
140mm
5mm

PharmaCode 578 -->

1mm
10mm
40mm lead-in

Package leaflet: Information for the user



One-a-Night 50_{mg} Tablets
Diphenhydramine Hydrochloride

Read all of this leaflet before you start taking this medicine because it contains important information for you. Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.

- Keep this leaflet, you may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse.

In this leaflet:

1. What Nytol One-A-Night is and what it is used for
2. What you need to know before you take Nytol One-A-Night
3. How to take Nytol One-A-Night
4. Possible side effects
5. How to store Nytol One-A-Night
6. Contents of the pack and other information

1. What Nytol One-A-Night is and what it is used for

Nytol One-A-Night is used to relieve temporary sleeping difficulties.

The active ingredient is diphenhydramine hydrochloride, an antihistamine that causes sleepiness or drowsiness. Temporary difficulties in getting a good night's sleep can be caused by various problems such as stress, jet lag or personal problems such as bereavement. In sleep problems of this type, Nytol One-A-Night has been shown to help sufferers to fall asleep faster, have a longer and deeper sleep, and to wake up feeling rested in the morning. Do not take Nytol One-A-Night for more than 2 weeks without consulting your doctor.

2. What you need to know before you take Nytol One-A-Night

Do not take Nytol One-A-Night:

- if you have ever had an allergic reaction to diphenhydramine hydrochloride or any of the other ingredients (listed in Section 6)
- if you have an obstruction in your stomach or gut (for example, because of an ulcer).

Warnings and precautions

Talk to your doctor or pharmacist before taking Nytol One-A-Night:

- if you have **asthma, bronchitis** or Chronic Obstructive Pulmonary Disease (COPD)
- if you have **narrow-angle glaucoma** (a form of raised pressure in the eye)
- if you have an **enlarged prostate** or **urinary retention disease**
- if you have **myasthenia gravis, epilepsy** or **seizure disorders**.
- If your **sleeping difficulties** persist, talk to your doctor or pharmacist.
- Nytol One-A-Night may get less effective if you use it continuously.

Other medicines and Nytol One-A-Night

Talk to your doctor or pharmacist before taking these tablets if you are taking any prescribed medicines; particularly **tricyclic antidepressants, monoamine oxidase inhibitors (MAOIs prescribed for depression)** within the last two weeks, **atropine** (used to dilate pupils of the eyes) or medicines which make you **drowsy**. Avoid the use of **other antihistamines** including those used on your skin and in cold and cough medicines, while taking this medicine.

Nytol One-A-Night with food, drink and alcohol
Do not drink alcohol while using Nytol One-A-Night.

PharmaCode 578 -->

LF03489C
H3L60FCJB1

140mm
125mm
3mm
5mm
150mm
140mm
5mm

PharmaCode 578 -->

1mm
10mm
40mm lead-in

! Pregnancy, breast feeding and fertility
Talk to your doctor before taking Nytol One-A-Night if you are pregnant, trying to become pregnant, or breast feeding.

! Driving and using machines
Do not drive or operate machinery.
Nytol One-A-Night is intended to produce sleepiness or drowsiness soon after the dose is taken.

Nytol One-A-Night contains lactose
If you have been told by your doctor that you have an **intolerance to some sugars**, ask your doctor before taking this medicine.

3. How to take Nytol One-A-Night
Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

✓ The recommended dose for adults and children aged 16 years and over: Swallow 1 tablet with water 20 minutes before you go to bed.

Not to be used by children under 16 years. Not to be used by the elderly with confusion.

! Do not take more than the recommended dose.

- Do not take extra tablets if you wake up in the night.
- Do not take for more than 2 weeks without consulting a doctor.

If you take more Nytol One-A-Night than you should
Contact your doctor or casualty department.

If you forget to take Nytol One-A-Night
Do not take a double dose to make up for a forgotten dose. Take the next dose at the usual time as described in this leaflet.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects
Like all medicines, Nytol One-A-Night can have side effects, although not everybody gets them.
Common (may affect up to 1 in 10 people)

- The most common side effects are: tiredness, drowsiness, reduced attention, unsteadiness, dizziness or a dry mouth.

Not known (frequency cannot be estimated from the available data)

- Allergic reactions (such as rash, shortness of breath and swelling), blurred vision, palpitations (feeling your heart beat) or irregular heartbeat.
- Muscle twitching, convulsions, headache or pins and needles.
- Also nausea (feeling sick), vomiting (being sick), thicker bronchial mucus and difficulty passing urine.

Reporting of side effects:
If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Nytol One-A-Night
Keep this medicine out of the sight and reach of children.
Do not use this medicine after the expiry date (EXP) stated on the pack.
Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.
Store in a dry place.

6. Contents of the pack and other information

What Nytol One-A-Night contains

- The active substance: Each tablet contains Diphenhydramine Hydrochloride 50 mg.
- The other ingredients are: Anhydrous lactose, microcrystalline cellulose, maize starch, stearic acid and silicon dioxide.

What Nytol One-A-Night looks like and contents of the pack
Packs of Nytol One-A-Night contain 4, 8, 10, 16 or 20 tablets. Not all pack sizes may be marketed.

Marketing authorisation holder is Omega Pharma Ltd., 32 Vauxhall Bridge Road, London, SW1V 2SA, United Kingdom and all enquiries should be sent to this address.

Manufacturer
Wrafton Laboratories Ltd, Wrafton, Braunton, North Devon, EX33 2DL, U.K.
This leaflet was last revised in February 2017.

LF03489C
H3L60FCJB1

100 mm
90
80
70
60
50
40
30
20
10
0

File title: 9914_LF03489C_H3L60FCJB1_Nytol OAN 50mg 20s PIL_UK_v7
Item No.: H3L60FCJB1
OPI No.: LF03489C
Last edit: 22nd May 2017
Brand: Nytol
Variant: One-A-Night 50mg
Component: Leaflet 20s
Market/s: UK
Dimensions: 140 x 150mm (F166v2)
Printer: Nampack Liverpool
Substrate: Paper
EAN Code: N/A
PharmaCode: 578
Replacement number:
Manufacturing Site: Perrigo Wrafton
No. of colours: 1 (one)

Process Black

one
agency.co

81 Rivington Street
Shoreditch
London EC2A 3AY
UK
awk contact:
Phil Hinckley
phil@oneagency.co
+44 (0)7710 521027
+44 (0)203 008 6041

Regulatory Approval:

Sign:..... Date:.....

Name (please print):.....

Contact email:.....

Marketing Approval:

Sign:.....

Name (please print):.....

Contact email:.....