



Otrivine

Extra Dual Relief

0.5mg/ml + 0.6mg/ml
Nasal Spray

Xylometazoline hydrochloride/
Ipratropium bromide

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist have told you.
- Keep this leaflet. You may need to read it again.
 - Ask your pharmacist if you need more information or advice.
 - If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
 - This medicine should not be used for more than 7 consecutive days.
 - You must talk to a doctor or pharmacist if you do not feel better or if you feel worse.
 - This medicine has been recommended for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.

What is in this leaflet:

1. What Otrivine Extra Dual Relief Nasal spray is and what it is used for
2. What you need to know before you use Otrivine Extra Dual Relief Nasal spray
3. How to use Otrivine Extra Dual Relief Nasal spray
4. Possible side effects
5. How to store Otrivine Extra Dual Relief Nasal spray
6. Contents of the pack and other information

1. WHAT OTRIVINE EXTRA DUAL RELIEF NASAL SPRAY IS AND WHAT IT IS USED FOR

Otrivine Extra Dual Relief Nasal spray is a combination medicinal product consisting of two different substances (xylometazoline hydrochloride and ipratropium bromide). One of the active ingredients (ipratropium bromide) helps with the runny nose, the other (xylometazoline hydrochloride) has a decongesting effect.

Otrivine Extra Dual Relief Nasal spray is used for the treatment of nasal congestion with runny nose (rhinorrhea) in connection with common colds.

2. WHAT YOU NEED TO KNOW BEFORE YOU USE OTRIVINE EXTRA DUAL RELIEF NASAL SPRAY

Do not use Otrivine Extra Dual Relief Nasal spray

- In children below 18 years of age, as adequate information on safety and efficacy is not available
- If you are allergic to Xylometazoline hydrochloride or ipratropium bromide or any of the other ingredients of this medicine (listed in section 6)
- If you are allergic to atropine or similar substances, e.g. hyoscyamine and scopolamine
- If you had your pituitary gland removed by surgery carried out through the nose
- If you have had brain surgery where the operation was carried out through the nose or mouth
- If you suffer from glaucoma (increased pressure in the eye)
- If you have a very dry nose (inflammatory nasal dryness, rhinitis sicca).

Warnings and precautions

Talk to your doctor or pharmacist before using Otrivine Extra Dual Relief Nasal spray if you suffer from:

- Heart Disease (e.g. long QT syndrome)
- Raised blood pressure
- Diabetes
- An overactive thyroid gland (hyperthyroidism)
- Difficulty in urinating and/or an enlarged prostate gland
- Narrow-angle glaucoma (sudden build-up of pressure in the eye)
- A tendency to have nosebleeds
- Obstruction of the gut (small intestine)
- Cystic fibrosis
- Benign tumour of the adrenal gland that produces high amounts of adrenaline and nor-adrenaline (phaeochromocytoma) or a particular sensitivity to adrenaline and nor-adrenaline as you may experience dizziness, tremor, rapid heart rate, raised blood pressure and sleeplessness.

Immediate hypersensitivity (allergic reaction) may occur in rare cases. This may be seen as itchy red rash with raised inflammation of the skin (hives), difficulty breathing or speaking, difficulty swallowing due to swelling of the lips, face or throat. These symptoms may appear individually or all combined as a severe allergic reaction. If this occurs, immediately STOP using this medicine (see section 4).

Worsening of narrow-angle glaucoma may also occur.

Prolonged or excessive use may cause stuffiness in the nose to return or worsen and swelling of the nasal mucosa.

Children, adolescents and elderly

Otrivine Extra Dual Relief Nasal spray is not recommended for use in children and adolescents below 18 years or in the elderly above 70 years of age, as adequate information on safety and efficacy is not available.

Other medicines and Otrivine Extra Dual Relief Nasal spray

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

It is particularly important that you mention the following:

- Monoamine oxidase inhibitors (used for treatment of depression). If you are taking or have taken these in the last two weeks, dangerous rises of the blood pressure may occur
- Tri-cyclic and tetra-cyclic antidepressants (if you are taking or have taken these in the last two weeks)
- Medicines used for travel sickness (medicines containing anticholinergic substances)
- Medicines used for gut disorders (particularly those for abnormal motility) (medicines containing anticholinergic substances).

If you use any of the above medicines, consult a doctor before using Otrivine Extra Dual Relief Nasal spray.

Pregnancy and breast-feeding

Otrivine Extra Dual Relief Nasal spray should not be used during pregnancy unless your doctor recommends it.

During breast-feeding Otrivine Extra Dual Relief Nasal spray should not be used unless your doctor decides the advantages outweigh the potential risks to the infant.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines:

Visual disturbances (including blurred vision and dilation of the pupil), dizziness and fatigue have been reported with Otrivine Extra Dual Relief Nasal spray. If affected, you should avoid driving, operating machinery or taking part in activities where these symptoms may put yourself or others at risk.

3. HOW TO USE OTRIVINE EXTRA DUAL RELIEF NASAL SPRAY

Always use this medicine exactly as described in this leaflet or your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is:

Adults: One puff in each nostril as needed, up to 3 times daily for maximum 7 days. Leave at least 6 hours between two doses. Do not exceed 3 applications daily into each nostril.

It is recommended that you stop treatment with Otrivine Extra Dual Relief Nasal spray as soon as your symptoms improve, even if this is sooner than 7 days, in order to minimise the risk of adverse reactions.

If you think the effect of this medicine is too strong or too weak, consult your doctor or pharmacist.

Instructions for use:

- Always blow your nose before using the nasal spray
- Remove the dust cap
- Do not cut the nozzle. The metered dose spray is ready to prime before use



- Before the first application, prime the pump by pumping 4 times. Once primed the pump will normally remain charged throughout regular daily treatment periods. Should the spray not be ejected during the full actuation stroke, or if the product has not been used for longer than 6 days, the pump will need to be reprimed with 4 pumps as initially performed
- Hold the bottle upright
- Bend your head slightly forward
- Close one nostril by placing your finger against the side of your nose and insert the spray tip into the other nostril. Press the pump quickly while inhaling through the nose
- Repeat this procedure in the other nostril

The effect occurs within 5-15 minutes.

Avoid spraying Otrivine Extra Dual Relief Nasal spray in or around the eyes. If this happens thoroughly rinse the eyes with cold water. Your vision may become temporarily blurred and the eye irritated, painful and red. If this happens contact your doctor for advice.

If you use more Otrivine Extra Dual Relief Nasal spray than you should:

If you, or someone else, have taken more medicine than you should, contact your doctor, hospital or emergency room for assessment of the risk. It is advisable to take this leaflet, bottle or pack with you. This is particularly important for children who are more likely to develop adverse events than adults.

Symptoms of overdose are severe dizziness, sweating, severely lowered body temperature, headache, slow heart beat, rapid heart beat, impaired breathing, coma, convulsion, hypertension (high blood pressure) which may be followed by hypotension (low blood pressure).

Other symptoms may be dry mouth, difficulties focusing with the eye and hallucinations.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

STOP using Otrivine Extra Dual Relief Nasal spray and seek medical help immediately if you have any of the following:

- Palpitations and increased heart rate (affects less than 1 in 100 people)
- Signs of an allergic reaction as difficulty breathing, speaking or swallowing; swelling of the face, lips, tongue or throat; severe itching of the skin, with a red rash or raised bumps (frequency not known cannot be estimated from available data)
- Disturbances of vision (including blurred vision, worsening of glaucoma or increased pressure in the eye), rainbow-coloured circles/haloes around bright lights and/or eye pain (frequency not known cannot be estimated from available data).

The most common side effects are nose bleeding and nasal dryness. Many of the side effects reported are also symptoms of common cold.

Very common side effects (may affect more than 1 in 10 people):

- Nose bleeding, nasal dryness.

Common side effects (may affect up to 1 in 10 people):

- Nasal discomfort, congestion of the nose, dry and irritated throat, pain in the nose
- Dry mouth
- Altered taste sensation, headache.

Uncommon (may affect up to 1 in 100 people):

- Nasal ulcer, sneezing, pain in the throat, cough, hoarseness
- Stomach upsets, nausea
- Altered smell sensation, dizziness, shakiness
- Discomfort, tiredness
- Sleeplessness
- Irritation of the eyes, dry eyes

Rare (may affect up to 1 in 1,000 people):

- Runny nose

Frequency not known (cannot be estimated from available data):

- Rash, hives
- Discomfort around the nose
- Discomfort in the chest, thirst
- Sudden spasm of throat muscle
- Irregular pulse
- Difficulties focusing with the eyes, dilation of the pupils
- Itching
- Difficulties emptying the bladder.

In order to minimise the risk of side effects such as nose bleeding and other effects on the nose, it is recommended that you stop treatment with Otrivine Extra Dual Relief Nasal spray when your symptoms improve even if this is sooner than 7 days.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE OTRIVINE EXTRA DUAL RELIEF NASAL SPRAY

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month.

Do not freeze.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Otrivine Extra Dual Relief Nasal spray contains

The active substances are xylometazoline hydrochloride and ipratropium bromide.

1 ml contains 0.5 mg xylometazoline hydrochloride and 0.6 mg ipratropium bromide.

1 puff contains 70 micrograms xylometazoline hydrochloride and 84 micrograms ipratropium bromide.

The other ingredients are disodium edetate, glycerol (85%), purified water, sodium hydroxide and hydrochloric acid.

What Otrivine Extra Dual Relief Nasal spray looks like and contents of the pack

Otrivine Extra Dual Relief Nasal spray is a clear solution.

The bottle contains approximately 70 puffs.

Otrivine Extra Dual Relief Nasal spray is available as a 10 ml nasal spray with a metered-dose spray pump.

Marketing Authorisation Holder and Manufacturer
GlaxoSmithKline Consumer Healthcare (UK) Trading Limited,
Brentford, TW8 9GS, U.K.

This medicinal product is authorised in the Member States of the EEA under the following names:

AT	Otrivin Duo 0,5 mg/ml + 0,6 mg/ml Nasenspray, Lösung
BE	Otrivine Duo 0,5mg/ml + 0,6mg/ml neusspray oplossing
BG	Otrivin Complex 0.5 mg/ml + 0.6 mg/ml nasal spray, solution
CY	Otrivin Advance
CZ	Otrivin Rhinostop
DE	Otriven Duo mit Xylometazolin und Ipratropium
DK	Otrivin Comp næsespray, opløsning
EE	Otrivin Total
EL	Otrivin Advance
ES	Rhinovín Duo 0,5 mg/ml + 0,6 mg/ml solución para pulverización nasal
FI	Otrivin Comp 0,5mg/ml + 0,6mg/ml nenäsümute, liuos
HU	Otrivin Komplex 0,5mg/ml + 0,6mg/ml oldatos orrspray
IE	Otrivine Extra Dual Relief Nasal spray Relief 0.5mg/ml, 0.6mg/ml Nasal Spray
IS	Otrivin Comp 0,5mg/ml + 0,6mg/ml nefuoi lausn
IT	RINAZINA DOPPIA AZIONE 0,5 mg/ml + 0,6 mg/ml spray nasale, soluzione
LT	OtriDuo 0,5 mg/ml + 0,6 mg/ml nosies purškalas, tirpalas
LU	Otrivine Duo
LV	Otrivin Total 0,5 mg/ml + 0,6mg/ml nasal spray solution
MT	Otrivine Extra Dual Relief 0.5mg/ml, 0.6mg/ml Nasal Spray
NL	Otrivin Duo Xylometazoline hydrochloride & Ipratropium bromide, 0,5/0,6 mg/ml, neusspray, oplossing
NO	Otrivin Comp 0,5mg/ml + 0,6mg/ml nesessay, opplosning
PL	Otrivin Ipra MAX
PT	Vibrocil ActilongDuo 0.5mg/ml + 0.6mg/ml solução para pulverização
RO	Vibrocil Duo 0,5mg/ml + 0,6mg/ml spray nazal, solutie
SE	Otricom 0,5mg/ml + 0,6 mg/ml nässpray lösning
SI	Otrivin Duo 0,5 mg/0,6 mg v 1 ml pršilo za nos, raztopina
SK	Otrivin Comp 0,5mg/ml + 0,6 mg/ml nässpray lösning
UK	Otrivine Extra Dual Relief 0.5 mg/ml + 0.6 mg/ml Nasal Spray

This leaflet was last revised in February 2019.

Trade marks are owned by or licensed to the GSK group of companies.

GB 944182 – 62000000037385