

Package leaflet: Information for the patient

Rivastigmine Luye 4.6 mg/24 h transdermal patch
Rivastigmine Luye 9.5 mg/24 h transdermal patch
Rivastigmine Luye 13.3 mg/24 h transdermal patch

rivastigmine

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Rivastigmine Luye is and what it is used for
2. What you need to know before you use Rivastigmine Luye
3. How to use Rivastigmine Luye
4. Possible side effects
5. How to store Rivastigmine Luye
6. Contents of the pack and other information

1. What Rivastigmine Luye is and what it is used for

The active substance of Rivastigmine Luye is rivastigmine.

Rivastigmine belongs to a class of substances called cholinesterase inhibitors. In patients with Alzheimer's dementia, certain nerve cells die in the brain, resulting in low levels of the neurotransmitter acetylcholine (a substance that allows nerve cells to communicate with each other). Rivastigmine works by blocking the enzymes that break down acetylcholine: acetylcholinesterase and butyrylcholinesterase. By blocking these enzymes, rivastigmine allows levels of acetylcholine to be increased in the brain, helping to reduce the symptoms of Alzheimer's disease.

Rivastigmine Luye is used for the treatment of adult patients with mild to moderately severe Alzheimer's dementia, a progressive brain disorder that gradually affects memory, intellectual ability and behaviour.

2. What you need to know before you use Rivastigmine Luye

Do not use Rivastigmine Luye

- if you are allergic to rivastigmine (the active substance in Rivastigmine Luye or any of the other ingredients of this medicine (listed in section 6).
- if you have ever had an allergic reaction to a similar type of medicine (carbamate derivatives).
- if you have a skin reaction spreading beyond the patch size, if there is a more intense local reaction (such as blisters, increasing skin inflammation, swelling) and if it does not improve within 48 hours after removal of the transdermal patch.

If this applies to you, tell your doctor and do not apply Rivastigmine Luye transdermal patches.

Warnings and precautions

Talk to your doctor before using Rivastigmine Luye

- if you have, or have ever had, a heart condition such as an irregular or slow heartbeat, QTc prolongation, a family history of QTc prolongation, torsade de pointes, or have a low blood level of potassium or magnesium.
- if you have, or have ever had, an active stomach ulcer.
- if you have, or have ever had, difficulties in passing urine.
- if you have, or have ever had, seizures.
- if you have, or have ever had, asthma or a severe respiratory disease.
- if you suffer from trembling.
- if you have a low body weight.
- if you have gastrointestinal reactions such as feeling sick (nausea), being sick (vomiting) and diarrhoea. You may become dehydrated (losing too much fluid) if vomiting or diarrhoea are prolonged.
- if you have impaired liver function.

If any of these apply to you, your doctor may need to monitor you more closely while you are on this medicine.

If you have not applied a patch for more than three days, do not apply the next one before you have talked to your doctor.

Children and adolescents

There is no relevant use of Rivastigmine Luye in the paediatric population in the treatment of Alzheimer's disease.

Other medicines and Rivastigmine Luye

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Rivastigmine Luye might interfere with anticholinergic medicines some of which are medicines used to relieve stomach cramps or spasms (e.g. dicyclomine), to treat Parkinson's disease (e.g. amantadine) or to prevent motion sickness (e.g. diphenhydramine, scopolamine, or meclizine).

Rivastigmine Luye patch should not be given at the same time as metoclopramide (a medicine used to relieve or prevent nausea and vomiting). Taking the two medicines together could cause problems such as stiff limbs and trembling hands.

If you have to undergo surgery whilst using Rivastigmine Luye transdermal patches, tell your doctor that you are using them because they may exaggerate the effects of some muscle relaxants during anaesthesia.

Caution when Rivastigmine Luye patch is given together with beta-blockers (medicines such as atenolol used to treat hypertension, angina, and other heart conditions). Taking the two medicines together could cause problems such as slowing of the heartbeat (bradycardia) leading to fainting or loss of consciousness.

Caution when Rivastigmine Luye is taken together with other medicines that can affect your heart rhythm or the electrical system of your heart (QT prolongation).

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

If you are pregnant, the benefits of using Rivastigmine Luye must be assessed against the possible effects on your unborn child. Rivastigmine Luye should not be used during pregnancy unless clearly necessary.

You should not breast-feed during treatment with Rivastigmine Luye transdermal patches.

Driving and using machines

Your doctor will tell you whether your illness allows you to drive vehicles and use machines safely. Rivastigmine Luye transdermal patches may cause fainting or severe confusion. If you feel faint or confused do not drive, use machines or perform any other tasks that require your attention.

3. How to use Rivastigmine Luye

Always use Rivastigmine Luye transdermal patches exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

How to start treatment

Your doctor will tell you which Rivastigmine Luye transdermal patch is most suitable for you.

- Treatment usually starts with Rivastigmine Luye 4.6 mg/24 h.
- The recommended usual daily dose is Rivastigmine Luye 9.5 mg/24 h. If well tolerated, the treating physician may consider increasing the dose to 13.3 mg/24 h
- Only wear one transdermal patch at a time and replace the patch with a new one after 24 hours.

During the course of the treatment your doctor may adjust the dose to suit your individual needs.


If you have not applied a patch for more than three days, do not apply the next one before you have talked to your doctor. Transdermal patch treatment can be resumed at the same dose if treatment is not interrupted for more than three days. Otherwise your doctor will restart your treatment on Rivastigmine Luye 4.6 mg/24 h.

Rivastigmine Luye can be used with food, drink and alcohol.

Where to apply your Rivastigmine Luye transdermal patch

- Before you apply a patch, make sure that your skin is clean, dry and hairless, free of any powder, oil, moisturiser, or lotion that could keep the patch from sticking to your skin properly, free of cuts, rashes and/or irritations.
- **Carefully remove any existing patch before putting on a new one.** Having multiple patches on your body could expose you to an excessive amount of this medicine which could be potentially dangerous.
- Apply **ONE** patch per day to **ONLY ONE** of the possible locations shown in the following diagrams:
 - left upper arm **or** right upper arm
 - left upper chest **or** right upper chest (**avoid breast**)
 - left upper back **or** right upper back
 - left lower back **or** right lower back



Every 24 hours take off the previous patch before putting ONE new patch on to ONLY ONE of the following possible locations.






When changing the patch, you must remove the previous day's patch before you apply the new one to a different location of skin each time (for example on the right side of your body one day, then on the left side the next day, and on your upper body one day, then on your lower body the next day). Do not apply a new patch to the same skin area twice within 14 days.

How to apply your Rivastigmine Luye transdermal patch

Rivastigmine Luye patches are thin, tan coloured patches that stick to the skin. Each patch is sealed in a sachet that protects it until you are ready to put it on. Do not open the sachet or remove a patch until just before you apply it.

	<p>Carefully remove the existing patch before putting on a new one.</p> <p>For patients starting treatment for the first time and for patients restarting rivastigmine after treatment interruption, please begin with the second picture.</p>
	<p>Each patch is sealed in its own protective sachet. You should only open the sachet when you are ready to apply the patch. Cut the sachet at both scissors marks, but not further than the indicated line. Tear the sachet to open. Do not cut the entire length of the sachet in order to avoid damaging of the patch.</p> <p>Remove the patch from the sachet.</p>

	<p>A protective liner covers the sticky side of the patch. Peel off one side of the protective liner and do not touch the sticky part of the patch with the fingers.</p>
	<p>Put the sticky side of the patch on the upper or lower back, upper arm or chest and then peel off the second side of the protective liner.</p>
	<p>Then press the patch firmly in place for at least 30 seconds using the palm of the hand to make sure that the edges stick well.</p>

IMPORTANT:

- **Take off the previous patch before putting ONE new patch on.**
- **Only one patch per day.**
- **Do not cut the patch into pieces.**
- **Press the patch firmly in place for at least 30 seconds using the palm of the hand.**

If it helps you, you may write, for example, the day of the week, on the patch with a thin ball point pen.

The patch should be worn continuously until it is time to replace it with a new one. You may wish to experiment with different locations when applying a new patch, to find ones that are most comfortable for you and where clothing will not rub on the patch.

How to remove your Rivastigmine Luye transdermal patch

Gently pull at one edge of the patch to remove it slowly from the skin. In case the adhesive residue is left over on your skin, gently soak the area with warm water and mild soap or use baby oil to remove it. Alcohol or other dissolving liquids (nail polish remover or other solvents) should not be used.

You should wash your hands with soap and water after removing the patch. In case of contact with eyes or if the eyes become red after handling the patch, rinse immediately with plenty of water and seek medical advice if symptoms do not resolve.

Can you wear your Rivastigmine Luye transdermal patch when you are bathing, swimming, or in the sun?

- Bathing, swimming or showering should not affect the patch. Make sure the patch does not loosen during these activities.
- Do not expose the patch to any external heat sources (e.g. excessive sunlight, saunas, solarium) for long periods of time.

What to do if a patch falls off

If a patch falls off, apply a new one for the rest of the day, then replace it at the same time as usual the next day.

When and for how long to apply your Rivastigmine Luye transdermal patch?

- To benefit from treatment, you must apply a new patch every day, preferably at the same time of day.
- Only wear one Rivastigmine Luye patch at a time and replace the patch with a new one after 24 hours.

If you use more Rivastigmine Luye than you should

If you accidentally apply more than one patch, remove all the patches from your skin, then inform your doctor that you have accidentally applied more than one patch. You may require medical attention. Some people who have accidentally taken too much rivastigmine have experienced feeling sick (nausea), being sick (vomiting), diarrhoea, high blood pressure and hallucinations. Slow heartbeat and fainting may also occur.

If you forget to use Rivastigmine Luye

If you find you have forgotten to apply a patch, apply one immediately. You may apply the next patch at the usual time the next day. Do not apply two patches to make up for the one that you missed.

If you stop using Rivastigmine Luye

Tell your doctor or pharmacist if you stop using the patch.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, Rivastigmine Luye transdermal patches can cause side effects, although not everybody gets them.

You may have side effects more often when you start your medicine or when your dose is increased. Usually, the side effects will slowly go away as your body gets used to the medicine.

Take off your patch and tell your doctor straight away, if you notice any of the following side effects which could become serious:

Common (may affect up to 1 in 10 people)

- Loss of appetite
- Feeling dizzy
- Feeling agitated or sleepy
- Urinary incontinence (inability to retain adequate urine)

Uncommon (may affect up to 1 in 100 people)

- Problems with your heartbeat such as slow heartbeat
- Seeing things that are not really there (hallucinations)
- Stomach ulcer
- Dehydration (losing too much fluid)
- Hyperactivity (high level of activity, restlessness)
- Aggression

Rare (may affect up to 1 in 1,000 people)

- Falling

Very rare (may affect up to 1 in 10,000 people)

- Stiff arms or legs
- Trembling hands

Not known (frequency cannot be estimated from the available data)

- Allergic reaction where the patch was used, such as blisters or inflamed skin
- The signs of Parkinson's disease get worse – such as tremor, stiffness and shuffling
- Pisa syndrome (a condition involving involuntary muscle contraction with abnormal bending of the body and head to one side)
- Inflammation of the pancreas – signs include serious upper stomach pain, often with feeling sick (nausea) or being sick (vomiting)
- Fast or uneven heartbeat
- High blood pressure
- Fits (seizures)
- Liver disorders (yellow skin, yellowing of the whites of the eyes, abnormal darkening of the urine or unexplained nausea, vomiting, tiredness and loss of appetite)
- Changes in tests which show how well the liver is working
- Feeling restless
- Nightmares

Take off your patch and tell your doctor straight away, if you notice any of the side effects above.

Other side effects seen with rivastigmine capsules or oral solution and which may occur with the patch:

Common (may affect up to 1 in 10 people)

- Too much saliva
- Loss of appetite
- Feeling restless
- Generally feeling unwell
- Trembling or feeling confused
- Increased sweating

Uncommon (may affect up to 1 in 100 people)

- Uneven heart rate (e.g. fast heart rate)
- Difficulty sleeping
- Accidental falls

Rare (may affect up to 1 in 1,000 people)

- Fits (seizures)
- Ulcer in the intestine
- Chest pain - this may be caused by heart spasm

Very rare (may affect up to 1 in 10,000 people)

- High blood pressure
- Inflammation of the pancreas – the signs include serious upper stomach pain, often with feeling sick (nausea) or being sick (vomiting)
- Bleeding in the gut – shows as blood in stools or when being sick
- Seeing things that are not there (hallucinations)
- Some people who have been violently sick have had tearing of the tube that connects your mouth with your stomach (oesophagus)

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Rivastigmine Luye

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date, which is stated on the carton and sachet after EXP. The expiry date refers to the last day of that month.
- Do not use any patch that is damaged or shows signs of tampering.
- After removing a patch, fold it in half with the sticky sides on the inside and press them together. Return the used patch to its sachet and dispose of it. Do not touch your eyes with your fingers and wash your hands with soap and water after removing the patch. If your community burns domestic rubbish, you can dispose of the patch with your domestic rubbish.
- Do not throw away any medicines via wastewater. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Rivastigmine Luye contains

The active substance is rivastigmine.

Rivastigmine Luye 4.6 mg/24 h transdermal patches:

Each transdermal patch releases 4.6 mg of rivastigmine per 24 hours. Each transdermal patch of 4.3 cm² contains 6.4 mg of rivastigmine.

Rivastigmine Luye 9.5 mg/24 h transdermal patches:

Each transdermal patch releases 9.5 mg of rivastigmine per 24 hours. Each transdermal patch of 8.5 cm² contains 12.8 mg of rivastigmine.

Rivastigmine Luye 13.3 mg/24 h transdermal patches:

Each transdermal patch releases 13.3 mg of rivastigmine per 24 hours. Each transdermal patch of 12.8 cm² contains 19.2 mg of rivastigmine.

The other ingredients are

Backing layer:

- polyethylene/thermoplastic resin/aluminium coated polyester film

Active layer:

- poly [(2-ethylhexyl)acrylate, vinylacetate] (1:1)

Adhesive matrix layer:

- medium molecular weight polyisobutene

- high molecular weight polyisobutene

- silica colloidal anhydrous
 - paraffin light liquid
- Release liner:
- polyester film fluoropolymer-coated
- Printing ink

What Rivastigmine Luye looks like and contents of the pack

Each transdermal patch is a thin patch of a circular shape. The outside of the backing layer is tan coloured and imprinted in orange with:

- “RIV-TDS 4.6 mg/24 h”
- “RIV-TDS 9.5 mg/24 h”
- “RIV-TDS 13.3 mg/24 h”

One transdermal patch is sealed in one sachet.

Rivastigmine Luye 4.6 and 9.5 mg/24 h transdermal patches are available in packs containing 7, 30 or 42 sachets and in multipacks containing 60 (2x30), 84 (2x42) or 90 (3x30) sachets.

Rivastigmine Luye 13.3 mg/24 h transdermal patches are available in packs containing 7 or 30 sachets and in multipacks containing 60 (2x30) or 90 (3x30) sachets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Luye Pharma Ltd.
40 Occam Road
Guildford, GU2 7YG
UK

Manufacturer

Luye Pharma AG
Am Windfeld 35
83714 Miesbach
Germany

This leaflet was last revised in: January 2025

You can order a patient diary and patient reminder card to help you use your Rivastigmine Luye transdermal patches free of charge from Luye Pharma Ltd., 40 Occam Road, Guildford, GU2 7YG, UK, at safety@luyepharma.co.uk or via phone: 0203 992 7900.