

PACKAGE LEAFLET

Package leaflet: Information for the user

ChloraPrep 2% w/v / 70% v/v impregnated cutaneous swab

chlorhexidine gluconate / isopropyl alcohol

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or nurse.
- If you get any side effects talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What ChloraPrep is and what it is used for
2. What you need to know before you use ChloraPrep
3. How to use ChloraPrep
4. Possible side effects
5. How to store ChloraPrep
6. Contents of the pack and other information

1. What ChloraPrep is and what it is used for

ChloraPrep is a cutaneous solution of chlorhexidine gluconate and isopropyl alcohol presented as a plastic stick with a foam pad impregnated with a fast-acting antiseptic solution on one end. The product is used to disinfect the skin and help prevent infections before invasive medical procedures.

2. What you need to know before you use ChloraPrep

Do not use ChloraPrep:

- if you are allergic (hypersensitive) to chlorhexidine gluconate or any of the other ingredients of ChloraPrep, especially if you have a history of possible chlorhexidine-related allergic reactions (see Section 6).
- in the ear canal as the solution may damage your hearing.

Warnings and precautions

ChloraPrep is for external use only.

ChloraPrep should not be used:

- near delicate linings (mucous membranes), as it may cause irritation. If it does get into the delicate linings to the body passages, it should be washed quickly with plenty of water.
- on open skin wounds.
- in direct contact with neural tissue (for example brain and spinal cord tissue).

ChloraPrep must not come into contact with the eye due to the risk of visual damage. If it comes into contact with the eyes, wash out immediately and thoroughly with water. In case of any irritation, redness or pain in the eye, or visual disturbance, ask for medical advice promptly.

Serious cases of persistent corneal injury (injury to the surface of the eye) potentially requiring corneal transplant have been reported when similar products have accidentally come in contact with the eye during surgical procedures, in patients under general anaesthesia (deep painless sleep).

ChloraPrep may in rare cases cause severe allergic reactions, leading to a drop in blood pressure and even to unconsciousness. Early symptoms of a severe allergic reaction may be skin rash or asthma. If you notice these

symptoms, stop using ChloroPrep and contact your doctor as soon as possible (see under section 4: “Possible side effects”).

ChloroPrep should only be applied to the skin gently. When the solution has been applied in an over-vigorous manner to very fragile or sensitive skin or after repeated use, rash, inflammation, itching, dry and/or flaky skin and pain may occur. At the first sign of any of these reactions, applications of ChloroPrep should be stopped.

Prolonged skin contact should be avoided.

Soaked materials, such as drapes or gowns should be removed before use. The solution should not be allowed to pool.

The solution is flammable. Do not use ignition sources such as cigarettes, naked flames or strong heat source until the skin is completely dry.

Children

Use with care in newborn babies, especially those born prematurely. ChloroPrep may cause chemical skin burns.

Other medicines and ChloroPrep

Tell your doctor or nurse if you have recently had a vaccine or skin test injection (patch test used to test for allergies).

Pregnancy, breast-feeding and fertility

There are no studies with ChloroPrep in pregnant or lactating women.

Pregnancy

No effects during pregnancy are anticipated, since systemic exposure to chlorhexidine gluconate is negligible. ChloroPrep can be used during pregnancy.

Breast-feeding

No effects on the breastfed newborn/infant are anticipated since the systemic exposure of the breast-feeding woman to chlorhexidine gluconate is negligible. ChloroPrep can be used during breast-feeding.

Fertility

The effects of chlorhexidine gluconate on human reproduction have not been studied.

Driving and using machines

ChloroPrep does not affect your driving or ability to use machines.

3. How to use ChloroPrep

The antiseptic solution is kept inside the aluminium pouch. Your doctor or nurse will select the ChloroPrep pouch size based on the procedure site and area to be covered. Your doctor or nurse will rub the foam pad gently over your skin, covering the skin area that needs to be prepared. Depending on your medical procedure, more than one pouch may be used.

Other ChloroPrep applicators may be more suitable, depending on the medical procedure being undertaken.

ChloroPrep is only used on the skin and each swab is only used once.

If you have any further questions on the use of this product, ask your doctor or nurse.

4. Possible side effects

Like all medicines, ChloroPrep can cause side effects, although not everybody gets them.

If you experience any of the following reactions the product application should be stopped: swelling of the face, lips, tongue or throat; a red itchy skin rash; wheezing or difficulty breathing; feeling faint and dizzy; a strange metallic taste in the mouth; collapse. You may be having an allergic reaction.

If you develop a rash or your skin becomes itchy, painful, red, blistery, dry or inflamed where you have used the product, stop using ChloroPrep and talk to your doctor or pharmacist.

Very rarely (fewer than 1 in 10,000 people), allergic or irritated skin reactions to the ingredients in ChloroPrep (chlorhexidine gluconate and isopropyl alcohol) have been reported.

Other possible side effects, for which it is not known how often they occur, are: eye irritation, pain, chemical burns, corneal injury (injury to the surface of the eye), and permanent eye damage including permanent visual impairment (following accidental ocular exposure during head, face and neck surgical procedures) in patients under general anaesthesia (deep painless sleep) and skin burns in newborns/infants.

Reporting of side effects

If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via

Yellow Card Scheme, Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side-effects you can help provide more information on the safety of this medicine.

5. How to store ChloroPrep

Flammable. Do not use with electrocautery procedures or other ignition sources until dry. Do not use while smoking or near any naked flames or strong heat source. This medicine does not require any special temperature storage conditions.

Store in the original packaging; ChloroPrep is sterile unless seal is broken.

Avoid exposure of the container and contents to naked flames during use, storage and disposal.

Do not use this medicine after the expiry date which is stated on the label or carton after EXP. The expiry date refers to the last day of that month.

Keep this medicine out of the sight and reach of children.

Do not throw away medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What ChloroPrep contains

- The active substances are chlorhexidine gluconate and isopropyl alcohol. Each swab of 1.75 ml solution contains 35 mg of chlorhexidine gluconate (20 mg/ml) and 1.23 ml isopropyl alcohol (0.70 ml/ml). Three swabs containing 5.25 ml of solution contains 105 mg of chlorhexidine gluconate (20 mg/ml) and 3.7 ml isopropyl alcohol (0.70 ml/ml).
- The other ingredient is purified water.

What ChloroPrep looks like and contents of the pack

The ChloroPrep presentation is a plastic stick connected to an impregnated foam tip in a plastic foil

containing the antiseptic solution.
The triple swabs pack contains three swabs.

Pack Size:

1.75 ml (Single swab):	Carton of 48 pouches (48 swabs)
5.25 ml (Triple swabs):	Carton of 40 pouches of 3 swabs (120 swabs)

Not all pack sizes may be marketed

Marketing Authorisation Holder and Manufacturer

The Marketing Authorisation holder of ChloraPrep is:

Becton Dickinson UK Ltd
1030 Eskdale Road, Winnersh,
Wokingham RG41 5TS,
United Kingdom
0800 0437 546

The Manufacturer of ChloraPrep is

Insight Health Limited
Unit A Bridgefields,
Welwyn Garden City, Hertfordshire AL7 1RX
United Kingdom

This medicinal product is authorised in the Member States of the EEA and in the United Kingdom (Northern Ireland) under the following names:

Finland - ChloraPrep
France - ChloraPrep
Germany -ChloraPrep
Ireland - ChloraPrep
Italy - ChloraPrep
Norway - Chloraprep
Portugal - Chloraprep
Sweden - Chloraprep
United Kingdom (Northern Ireland) - ChloraPrep

This leaflet was last revised in June 2024.

BD, the BD Logo, ChloraPrep are the property of Becton, Dickinson and Company or its affiliates. ©2024 BD. All rights reserved.

License Number:
PL 05920/0006

The following information is intended for healthcare professionals only:

ChloraPrep 2% w/v / 70% v/v impregnated cutaneous swab

chlorhexidine gluconate / isopropyl alcohol

Instructions for using ChloraPrep:

For cutaneous use. For external use only. Choose the appropriate ChloraPrep presentation based on the invasive procedure being undertaken and the size of the area to be prepped to avoid excess of solution and the risk of product pooling.

Other ChloraPrep applicators may be more suitable depending on the medical procedure being undertaken.

- Only the inside of the pouch is sterile. In order to preserve the sterility, the pouch should be opened carefully, and the foam pad should not be allowed to touch any non-sterile surface.
- Tear pouch at side notch to reveal swab handle(s). Do not touch swab foam pad.
- Place foam pad flat side down on the treatment area.
- Apply the solution for 30 seconds using a gentle back and forth motion.
- Allow the covered area to air dry completely.
- For triple swabs, use swabs sequentially for a total of 30 seconds.

ChloraPrep solution can be left on the skin post procedure.

Maximum coverage areas:

- Single swab (1.75 ml) 6 cm x 7 cm
- Triple swabs (5.25ml) 10 cm x 16 cm

Precautions for Use:

- Allow ChloraPrep solution to dry completely on the skin before starting any medical procedure. Do not use electrocautery procedures or other ignitions until the skin is completely dry. Do not use while smoking or near any naked flames or strong heat source.
- Do not use excessive quantities and do not allow the solution to pool in skin folds or under the patient or drip on sheets or other material in direct contact with the patient.
- Use with care in neonates, especially those born before 32 weeks of gestation and within the first 2 weeks of life. ChloraPrep may cause chemical skin burns.
- Do not use near mucous membranes, as it may cause irritation, pain and chemical burns. If it does get into the mucous membranes, it should be washed quickly with plenty of water.
- ChloraPrep must not come into contact with the eye due to the risk of visual damage. If it comes into contact with the eyes, wash out immediately and thoroughly with water. An ophthalmologist's advice should be sought.
- Do not use on open skin wounds, broken or damaged skin.
- ChloraPrep should not come into contact with neural tissues. To prevent exposure of the middle ear to ChloraPrep via a perforation of the eardrum, the use of ChloraPrep in the ear canal is contraindicated.
- Chlorhexidine is incompatible with soap and other anionic agents.
- Alcohol should not be brought into contact with some vaccines and skin test injections (patch tests). If in doubt, consult the vaccine manufacturer's literature.
- Do not apply the solution in an over vigorous manner to very fragile or sensitive skin. After repeated use, local skin reaction may occur including erythema or inflammation, itching, dry and/or flaky skin and local application site pain. At the first sign of local skin reaction, stop application of ChloraPrep.
- Do not use in patients with known hypersensitivity to ChloraPrep solution or any of its components, especially in those with a history of possible chlorhexidine related allergic reactions. Chlorhexidine-containing products are known causes of anaphylactic reactions during anaesthesia. If symptoms of an anaphylactic reaction are detected during anaesthesia (e.g. abrupt fall in blood pressure, hives, angioedema), chlorhexidine-related allergic reaction should be considered.
- Special precaution should be taken to avoid patient exposure to any other product containing chlorhexidine during the course of the treatment.

Special precautions for disposal

The solution is flammable. Do not use while smoking or near any naked flames or strong heat source. Avoid exposure of the container and contents to naked flames during use, storage and disposal. Discard the swab after use as per clinical waste procedures.

Please refer to the Summary of Product Characteristics of ChloraPrep for further detailed information.

Storage Procedures

ChloraPrep is for single use only and is sterile until the packaging is opened. Do not use ChloraPrep after the expiry date stated on the label or carton. The expiry date refers to the last day of that month. This medicine does not require any special temperature storage conditions. Store in the original packaging.

Active Substances

The active substances in ChloroPrep are 2% w/v chlorhexidine gluconate and 70% v/v isopropyl alcohol. The inactive ingredient in ChloroPrep is purified water.

Marketing Authorisation Holder

Becton Dickinson UK Ltd
1030 Eskdale Road, Winnersh,
Wokingham, RG41 5TS,
United Kingdom

BD, the BD Logo, ChloroPrep are the property of Becton, Dickinson and Company or its affiliates.
©2024 BD. All rights reserved.