

## **Package leaflet: Information for the user**

**Losartan potassium 25 mg film-coated tablets**  
**Losartan potassium 50 mg film-coated tablets**  
**Losartan potassium 100 mg film-coated tablets**  
losartan potassium

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist, or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist, or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

### **What is in this leaflet**

1. What Losartan potassium is and what it is used for
2. What you need to know before you take Losartan potassium
3. How to take Losartan potassium
4. Possible side effects
5. How to store Losartan potassium
6. Contents of the pack and other information

### **1. What Losartan potassium is and what it is used for**

This medicine belongs to a group of medicines known as angiotensin-II receptor antagonists. Angiotensin-II is a substance produced in the body which binds to receptors in blood vessels, causing them to tighten. This results in an increase in blood pressure. Losartan prevents the binding of angiotensin-II to these receptors, causing the blood vessels to relax which in turn lowers the blood pressure. Losartan slows the decrease of kidney function in patients with high blood pressure and type 2 diabetes.

Losartan potassium is used

- to treat patients with high blood pressure (hypertension) in adults and in children and adolescents 6 - 18 years of age.
- to protect the kidney in hypertensive type 2 diabetic patients with laboratory evidence of impaired kidney function and proteinuria  $\geq 0.5$  g per day (a condition in which urine contains an abnormal amount of protein).
- to treat patients with chronic heart failure when therapy with specific medicines called angiotensin-converting-enzyme inhibitors (ACE inhibitors, medicine used to lower high blood pressure) is not considered suitable by your doctor. If your heart failure has been stabilised with an ACE inhibitor you should not be switched to losartan.
- in patients with high blood pressure and a thickening of the left ventricle, Losartan potassium has been shown to decrease the risk of stroke ("LIFE indication").

### **2. What you need to know before you take Losartan potassium**

**Do not take Losartan potassium**

- if you are allergic to losartan or to any of the other ingredients of this medicine (listed in section 6),
- if you are more than 3 months pregnant (it is also better to avoid Losartan potassium in early pregnancy – see section 2 “Pregnancy and breast-feeding”),
- if your liver function is severely impaired,
- if you have diabetes or impaired kidney function and you are treated with a blood pressure lowering medicine containing aliskiren.

### **Warnings and precautions**

Talk to your doctor, pharmacist, or nurse before taking Losartan potassium.

You must tell your doctor if you think you are (or might become) pregnant. Losartan potassium is not recommended in early pregnancy and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see section 2 “Pregnancy and breast-feeding”).

It is important to tell your doctor before taking Losartan potassium:

- if you have had a history of angioedema (swelling of the face, lips, throat, and/or tongue) (see also section 4 ‘Possible side effects’),
- if you suffer from excessive vomiting or diarrhoea leading to an extreme loss of fluid and/or salt in your body,
- if you receive diuretics (medicines that increase the amount of water that you pass out through your kidneys) or are under dietary salt restriction leading to an extreme loss of fluid and salt in your body (see section 3 ‘Doses in special patient groups’),
- if you are known to have narrowing or blockage of the blood vessels leading to your kidneys or if you have received a kidney transplant recently,
- if your liver function is impaired (see sections 2 “Do not take Losartan potassium” and section 3 “Doses in special patient groups”),
- if you suffer from heart failure with or without kidney impairment or concomitant severe life threatening cardiac arrhythmias. Special caution is necessary when you are treated with a  $\beta$ -blocker at the same time,
- if you have problems with your heart valves or heart muscle,
- if you suffer from coronary heart disease (caused by a reduced blood flow in the blood vessels of the heart) or from cerebrovascular disease (caused by a reduced blood circulation in the brain),
- if you suffer from primary hyperaldosteronism (a syndrome associated with increased secretion of the hormone aldosterone by the adrenal gland, caused by an abnormality within the gland),
- if you are taking any of the following medicines used to treat high blood pressure:
  - an ACE-inhibitor (for example enalapril, lisinopril, ramipril), in particular if you have diabetes-related kidney problems.
  - aliskiren

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (such as potassium) in your blood at regular intervals.

See also information under the heading “Do not take Losartan potassium”.

- if you are taking other medications that may increase serum potassium (see section 2 “Other medicines and Losartan potassium”).

Talk to your doctor if you experience abdominal pain, nausea, vomiting or diarrhea after taking Losartan potassium. Your doctor will decide on further treatment. Do not stop taking Losartan Potassium on your own.

**Children and adolescents**

Losartan potassium is not recommended for use in children suffering from kidney or liver problems, as limited data are available in these patient groups. Losartan potassium is not recommended for use in children under 6 years old, as it has not been shown to work in this age group.

Losartan potassium has been studied in children. For more information, talk to your doctor.

**Other medicines and Losartan potassium**

Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines.

Tell your doctor if you are taking potassium supplements, potassium-containing salt substitutes, potassium-sparing medicines such as certain diuretics (amiloride, triamterene, spironolactone), or other medicines that may increase serum potassium (such as heparin, trimethoprim-containing medicines), as the combination with Losartan potassium is not advisable.

Take particular care if you are taking the following medicines while under treatment with Losartan potassium:

- other blood pressure lowering medicines as they may additionally reduce your blood pressure. Blood pressure may also be lowered by one of the following medicines/class of medicines: tricyclic antidepressants, antipsychotics, baclofen, amifostine,
- non-steroidal anti-inflammatory drugs such as indomethacin, including COX-2-inhibitors (medicines that reduce inflammation, and can be used to help relieve pain) as they may reduce the blood pressure lowering effect of losartan.

Your doctor may need to change your dose and/or to take other precautions:

If you are taking an ACE-inhibitor or aliskiren (see also information under the headings “Do not take Losartan potassium” and “Warnings and precautions”).

If your kidney function is impaired, the use of these medicines at the same time may lead to a worsening of the kidney function.

Lithium containing medicines should not be taken at the same time with losartan without close supervision by your doctor. Special precautionary measures (such as blood tests) may be appropriate.

**Losartan Tablets with food and drink**

Losartan Tablets may be taken with or without food.

Grapefruit juice should be avoided while taking Losartan Tablets.

**Pregnancy and breast-feeding****Pregnancy**

You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking Losartan potassium before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Losartan potassium. This medicine is not recommended in early pregnancy and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

**Breast-feeding**

Tell your doctor if you are breast-feeding or about to start breast-feeding. Losartan potassium is not recommended for mothers who are breast-feeding, and your doctor may choose another treatment for you if you wish to breast-feed, especially if your baby is a new-born, or born prematurely.

Ask your doctor or pharmacist for advice before taking this medicine.

### **Driving and using machines**

No studies on the effects on the ability to drive and use machines have been performed.

Losartan potassium is unlikely to affect your ability to drive or use machines. However, as with many other medicines used to treat high blood pressure, losartan may cause dizziness or drowsiness in some people. If you experience dizziness or drowsiness, you should consult your doctor before attempting such activities.

### **3. How to take Losartan potassium**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Your doctor will decide on the appropriate dose of Losartan potassium, depending on your condition and whether you are taking other medicines. It is important to continue taking this medicine for as long as your doctor prescribes it in order to maintain smooth control of your blood pressure.

#### **To treat high blood pressure**

##### **Use in adults**

Treatment usually starts with 50 mg losartan (one tablet Losartan potassium 50 mg) once a day. The maximal blood pressure lowering effect should be reached 3-6 weeks after beginning treatment. In some patients the dose may later be increased to 100 mg losartan (two tablets Losartan potassium 50 mg or one tablet of Losartan potassium 100 mg) once daily.

If you have the impression that the effect of losartan is too strong or too weak, please talk to your doctor or pharmacist.

##### **Children below 6 years of age**

Losartan Tablets is not recommended for use in children under 6 years old, as it has not been shown to work in this age group.

##### **Use in children and adolescents aged 6 - 18 years old**

The recommended starting dose in patients who weigh between 20 and 50 kg is 0.7 mg of losartan per kg of body weight administered once a day (up to 25 mg of Losartan potassium). The doctor may increase the dose if blood pressure is not controlled.

Other form(s) of this medicine may be more suitable for children; ask your doctor or pharmacist.

##### **Use to treat adult patients with high blood pressure and type 2 diabetes**

Treatment usually starts with 50 mg losartan (one tablet Losartan potassium 50 mg) once a day. The dose may later be increased to 100 mg losartan (two tablets Losartan potassium 50 mg or one tablet of Losartan potassium 100 mg) once daily depending on your blood pressure response.

Losartan may be administered with other blood pressure lowering medicines (e.g. diuretics, calcium channel blockers, alpha- or beta-blockers, and centrally acting agents) as well as with insulin and other commonly used medicines that decrease the level of glucose in the blood (e.g. sulfonylureas, glitazones and glucosidase inhibitors).

##### **Use to treat adult patients with heart failure**

Treatment usually starts with 12.5 mg losartan once a day. Generally, the dose should be increased weekly step-by-step (i.e., 12.5 mg daily during the first week, 25 mg daily during the second week, 50 mg daily during the third week, 100 mg daily during the fourth week, 150 mg daily during the fifth week) up to the maintenance dose as determined by your physician. A maximum dose of 150 mg losartan (for

example, three tablets of Losartan potassium 50 mg or one tablet each of Losartan potassium 100 mg and Losartan potassium 50 mg) once daily may be used.

In the treatment of heart failure, losartan is usually combined with a diuretic (medicine that increases the amount of water that you pass out through your kidneys) and/or digitalis (medicine that helps to make the heart stronger and more efficient) and/or a beta-blocker.

### **Doses in special patient groups**

The doctor may advise a lower dose, especially when starting treatment in certain patients such as those treated with diuretics in high doses, in patients with liver impairment, or in patients over the age of 75 years. The use of losartan is not recommended in patients with severe liver impairment (see section "Do not take Losartan potassium").

### **Administration**

The tablets should be swallowed whole with a glass of water.

You should try to take your daily dose at about the same time each day. It is important that you continue to take Losartan potassium until your doctor tells you otherwise.

### **If you take more Losartan potassium than you should**

If you accidentally take too many tablets, contact your doctor immediately. Symptoms of overdose are low blood pressure, increased heartbeat, possibly decreased heartbeat.

### **If you forget to take Losartan potassium**

If you accidentally miss a daily dose, just take the next dose as normal. Do not take a double dose to make up for a forgotten tablet.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist, or nurse.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If you experience the following, **stop taking this medicine and tell your doctor immediately or go to the casualty department of your nearest hospital:**

- A severe allergic reaction (rash, itching, swelling of the face, lips, mouth or throat that may cause difficulty in swallowing or breathing).

This is a serious but rare side effect, which affects more than 1 out of 10,000 patients but fewer than 1 out of 1,000 patients. You may need urgent medical attention or hospitalisation.

The following side effects have been reported with Losartan potassium:

**Common** (may affect up to 1 in 10 people):

- dizziness,
- low blood pressure (especially after excessive loss of water from the body within blood vessels e.g. in patients with severe heart failure or under treatment with high dose diuretics),
- dose-related orthostatic effects such as lowering of blood pressure appearing when rising from a lying or sitting position,
- debility (weakness),
- fatigue (tiredness),
- too little sugar in the blood (hypoglycaemia),
- too much potassium in the blood (hyperkalaemia),

- changes in kidney function including kidney failure,
- reduced number of red blood cells (anaemia),
- increase in blood urea, serum creatinine and serum potassium in patients with heart failure.

**Uncommon** (may affect up to 1 in 100 people):

- somnolence,
- headache,
- sleep disorders,
- feeling of increased heart rate (palpitations),
- severe chest pain (angina pectoris),
- shortness of breath (dyspnoea),
- abdominal pain,
- obstipation,
- diarrhoea,
- nausea (feeling sick),
- vomiting (being sick),
- hives (urticaria),
- itching (pruritus),
- rash,
- localised swelling (oedema),
- cough.

**Rare** (may affect up to 1 in 1,000 people):

- hypersensitivity,
- angioedema (swelling),
- inflammation of blood vessels (vasculitis including Henoch-Schönlein purpura),
- numbness or tingling sensation (paraesthesia),
- fainting (syncope),
- very rapid and irregular heartbeat (atrial fibrillation),
- brain attack (stroke),
- inflammation of the liver (hepatitis),
- elevated blood alanine aminotransferase (ALT) levels, usually resolved upon discontinuation of treatment.
- Intestinal angioedema: a swelling in the gut presenting with symptoms like abdominal pain, nausea, vomiting and diarrhoea.

**Not known** (frequency cannot be estimated from the available data):

- reduced number of thrombocytes,
- migraine,
- liver function abnormalities,
- muscle and joint pain,
- flu-like symptoms,
- back pain and urinary tract infection,
- increased sensitivity to the sun (photosensitivity),
- unexplained muscle pain with dark (tea-coloured) urine (rhabdomyolysis),
- impotence,
- inflammation of the pancreas (pancreatitis),
- low levels of sodium in the blood (hyponatraemia),

- depression,
- generally feeling unwell (malaise),
- ringing, buzzing, roaring, or clicking in the ears (tinnitus),
- disturbed taste (dysgeusia).

Side effects in children are similar to those seen in adults.

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist, or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Losartan potassium**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton. The expiry date refers to the last day of that month.

Do not store above 25°C. Store in the original package in order to protect from light and moisture. Do not open the blister pack until you are ready to take the medicine.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Losartan potassium contains**

- The active substance is losartan potassium.  
Each Losartan potassium 25 mg film-coated tablet contains 25 mg of losartan potassium.  
Each Losartan potassium 50 mg film-coated tablet contains 50 mg of losartan potassium.  
Each Losartan potassium 100 mg film-coated tablet contains 100 mg of losartan potassium.
- The other ingredients are:  
Tablet core: microcrystalline cellulose (E460), maize starch, sodium starch glycolate (Type A), colloidal anhydrous silica, magnesium stearate (E572), opadry white 04F580012, purified water.  
Film-coat: hydroxypropylmethyl cellulose (E464), talc, titanium dioxide (E171), polyethylene glycol.

Losartan potassium 25 mg, 50 mg and 100 mg film-coated tablets contain potassium in the following amounts: 2.12 mg (0.054 mEq), 4.24 mg (0.108 mEq) and 8.48 mg (0.216 mEq) respectively.

### **What Losartan potassium looks like and contents of the pack**

Losartan potassium 25 mg are white to off-white, round shaped, approx. 5.5 mm in diameter, biconvex, film-coated tablets debossed with "J" on one side and "I" on other side.

Losartan potassium 50 mg are white to off white, round shaped, approx. 7.5 mm in diameter, biconvex, film-coated tablets debossed with "J" on one side and on other side "I" on both side of score notch. The score line is not intended for breaking the tablet.

Losartan potassium 100 mg are white to off white, almond shaped, approx. 11.5 mm x 7.0 mm in dimension, biconvex, film-coated tablets debossed with "J" on one side and "12" on the other side.

Losartan potassium 25 mg, 50 mg and 100 mg film-coated tablets are supplied in white opaque PVC/Alu blister pack.

Pack size: 7, 14, 28, 56 and 84 tablets.

Not all pack sizes may be marketed.

**Marketing Authorisation Holder and Manufacturer**

**Ipca Laboratories UK Ltd.**

Unit 97-98, Silverbriar, Sunderland Enterprise Park East,

Sunderland, SR5 2TQ

United Kingdom

**This leaflet was last revised in 06/2025**