

## **Package Leaflet : Information for the User**

### **Apraclonidine 1% w/v Eye Drops, Solution (Apraclonidine Hydrochloride)**

**Read all of this leaflet carefully before you start using this medicine.**

Keep this leaflet. You may need to read it again.

If you have any further questions, ask your doctor or pharmacist.

If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet

1. What Apraclonidine 1% w/v Eye Drops, Solution is and what it is used for
2. Before you are given Apraclonidine 1% w/v Eye Drops, Solution
3. How Apraclonidine 1% w/v Eye Drops, Solution is given to you
4. Possible side effects
5. How Apraclonidine 1% w/v Eye Drops, Solution is stored
6. Further information

Apraclonidine 1% w/v Eye Drops, Solution **will be put in by your doctor. This leaflet is provided to give you information on your medicine.**

#### **1. What Apraclonidine 1% w/v Eye Drops, Solution is and what it is used for**

Apraclonidine 1% w/v Eye Drops, Solution eye drops contain apraclonidine which is used to control or prevent the pressure in the eye increasing after laser surgery.

#### **2. Before you are given Apraclonidine 1% w/v Eye Drops, Solution**

This medicine may not be suitable for you. If it is not suitable your doctor may want to give you another medicine so it is important you check all of the points below:

Apraclonidine 1% w/v Eye Drops, Solution **will not be used**

- \* in children (this medicine is not for anyone under the age of 18);
- \* If you are **allergic** to clonidine, apraclonidine or any of the ingredients listed in the “**Other ingredients**” section of this leaflet (section 6)
- \* if you have a history of any severe or unstable and uncontrolled heart disease (including chest pain, angina, heart attacks or heart failure);
- \* if you are currently taking monoamine oxidase inhibitor antidepressants

If you think any of the above conditions apply to you or if you are not sure, talk to your doctor first.

#### **Special care will be taken using Apraclonidine 1% w/v Eye Drops, Solution**

**Ask your doctor for advice if:**

- \* **you suffer from, or are taking medication for:**
  - heart disease (including chest pain, angina, heart attacks or heart failure);
  - high blood pressure;
  - circulatory problems (including stroke, Raynaud’s disease or Buerger’s disease);
  - vasovagal attacks (fainting spells);
  - reduced kidney or liver function;

If any of the above apply to you, you may be at greater risk for effects on your heart or circulatory system so your blood pressure and heart rate will be closely monitored.

- \* **you suffer from depression**, as medicines like clonidine have been associated with depression;
- \* **you suffer from diabetes** or low blood sugar, as this medicine may hide the signs and symptoms of a sudden reduction in blood sugar such as a fast heartbeat or trembling;
- \* **you regularly drink alcohol**, as Apraclonidine 1% w/v Eye Drops, Solution may increase its effect;
- \* **you have had an extreme response to other medicines** that lower the pressure inside your eye, so your eye pressure should be monitored closely;

If any of the above applies to you, or if you are not sure, talk to your doctor first.

#### **Using other medicines *Apraclonidine 1% w/v Eye Drops, Solution will not be used***

- \* **if you are taking monoamine oxidase inhibitors (used to treat depression)** e.g. phenelzine, isocarboxazid, tranylcypromine, moclobemide
- \* **if you are taking tricyclic antidepressants (often used to treat depression)** e.g. amitriptyline, imipramine, doxepin, mianserin, trazodone, dosulepin, lofepramine;
- \* **if you are currently using medicines containing sympathomimetic agents.**

These agents are used to treat many diseases, including asthma e.g. salbutamol, terbutaline, bambuterol, Parkinson's disease and heart disease and also may be present in cough and cold remedies.

#### ***Special care will be taken using Apraclonidine 1% w/v Eye Drops, Solution***

- \* **if you are using eye drops to treat glaucoma** which contain sympathomimetics e.g. phenylephrine, brimonidine, dipivefrine as your blood pressure may increase; or
- \* **if you are taking sleeping tablets, sedatives or painkillers**, as Apraclonidine 1% w/v Eye Drops, Solution may increase their effects;
- \* **if you are taking or have taken neuroleptics** (major tranquillisers used to treat some forms of mental illness e.g. chlorpromazine, flupentixol, haloperidol) as your blood pressure could become too low;

When there might be effects on your blood pressure then your blood pressure and heart rate will be closely monitored.

Discuss any medicines you are taking with your doctor before you are given Apraclonidine 1% w/v Eye Drops, Solution, including medicines obtained without a prescription.

#### **Using Apraclonidine 1% w/v Eye Drops, Solution with food and drink**

Apraclonidine 1% w/v Eye Drops, Solution should be used with caution if you regularly drink alcohol or have recently drunk alcohol.

#### **Use during pregnancy and breastfeeding**

Apraclonidine 1% w/v Eye Drops, Solution should not be used if you are pregnant, planning to become pregnant or are breastfeeding,

Your doctor should have discussed this with you. If not, ask your doctor for advice before taking any medicine.

#### **Driving and using machines**

This type of medicine may cause you to feel sleepy. If you are affected, do not drive or use any tools or machines.

### **3. How Apraclonidine 1% w/v Eye Drops, Solution is given to you**

Adults and elderly

Apraclonidine 1% w/v Eye Drops, Solution comes in a specially designed container which can be used only once. Your doctor will put in one drop 1 hour before your laser surgery and another drop immediately after surgery.

After being given Apraclonidine 1% w/v Eye Drops, Solution, you will be instructed to press a finger to the corner of your eye, by your nose. This helps to stop Apraclonidine 1% w/v Eye Drops, Solution getting into the rest of the body.

If more than one ophthalmic medicinal product is being used, the medicines must be administered at least 5 minutes apart. Eye ointments should be administered last.

Children

This medicine is not for use in children.

#### **4. Possible side effects**

Like all medicines, Apraclonidine 1% w/v Eye Drops, Solution can have side effects, although not everybody gets them.

**When used before and after laser surgery**, you may experience some or all of the following reactions **in your eyes**.

##### **Common**

(may affect up to one in 10 people treated)

- redness in the eye or eyes.
- one pupil getting bigger
- inability to close eye
- an eye disorder (whitening of the surface of the eye)
- an unusual or uncomfortable feeling in your eyes
- your eye feeling dry

##### **Uncommon**

(may affect up to one in 100 people treated)

- bleeding on the surface of your eye (conjunctival haemorrhage);
- eye inflammation (which may include pain, heat, redness or swelling of the eye); or
- your upper eyelid lifting up.
- extremely low pressure inside the eye (your doctor will see this during eye tests);
- reduced or dim vision;
- bleeding on the surface of your eye;
- blurred vision;
- eye allergy;
- your upper eyelid lifting;
- your eye itching;
- an eye disorder (whitening of the surface of the eye);
- eye irritation;

Or, you may suffer reaction in **other areas of your body**.

##### **Common**

(may affect up to one in 10 people treated)

- a dry feeling in your nose
- a bad or unusual taste in your mouth;

##### **Uncommon**

(may affect up to one in 100 people treated)

- an irregular heartbeat;
- difficulty sleeping;
- disturbed dreams;
- fainting spells;
- burning or tingling sensations;
- a lower sex drive;
- feeling irritated;
- a slow heartbeat or palpitations;
- dizziness or feeling faint when you sit up or stand up;
- diarrhoea;
- being sick or vomiting;
- pain or discomfort in your stomach;
- headaches;
- tiredness;

### **Rare**

(may affect up to one in 1000 people treated)

- reduced sensation or numbness;
- shortness of breath;
- a runny nose or too much saliva;
- a dry feeling, or burning in your nose;
- a dry mouth;
- increased sweating;
- itching skin;
- pain in your arms or legs;
- heaviness or pain in your chest;
- feeling hot or cold.

Not known (frequency cannot be estimated from available data);

- decreased blood pressure
- increased blood pressure

If you experience an allergic reaction, feel unusually dizzy or lightheaded, or experience irregular heartbeats, you should tell your doctor immediately.

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system (see details below). By reporting side effects you can help provide more information on the safety of this medicine.

United Kingdom

Yellow Card Scheme

Website : [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store

## **5. How Apraclonidine 1% w/v Eye Drops, Solution is stored**

- This medicine should not be used after the expiry date shown as 'EXP' on the package and plastic container. The expiry date refers to the last day of that month.
- This medicine should not be stored above 25°C and should be kept in the outer carton to protect it from the light.
- Keep out of the reach and sight of children.
- The medicine should be used immediately after opening the container.

## 6. Further information

**The active substance is** apraclonidine 1% w/v (10mg/ml) (as hydrochloride).

**Other ingredients are** sodium acetate, sodium chloride and purified water. Very small amounts of hydrochloric acid or sodium hydroxide may have been added to make the drops comfortable for your eyes.

**Apraclonidine 1% w/v Eye Drops, Solution eye drops solution** is pale yellow and comes in a container of 0.25 ml.

### **Marketing authorisation holder:**

Essential Pharma Limited,  
Vision Exchange Building  
Triq it-Territorjals, Zone 1,  
Central Business District,  
Birkirkara, CBD 1070,  
Malta

### **Manufacturer:**

Kaysersberg Pharmaceuticals S.A.S  
23 Avenue Georges Ferrenbach  
68240 Kaysersberg  
France

This leaflet was last revised in 05/2025