

Package leaflet: Information for the patient


Apremilast Krka 10 mg
film-coated tablets**Apremilast Krka 20 mg**
film-coated tablets**Apremilast Krka 30 mg**
film-coated tablets

apremilast

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

From here onwards the product name will be referred to as Apremilast.



What is in this leaflet

1. What Apremilast is and what it is used for
2. What you need to know before you take Apremilast
3. How to take Apremilast
4. Possible side effects
5. How to store Apremilast
6. Contents of the pack and other information

1. What Apremilast is and what it is used for

What Apremilast is

Apremilast contains the active substance 'apremilast'. This belongs to a group of medicines called phosphodiesterase 4 inhibitors, which help to reduce inflammation.

What Apremilast is used for

Apremilast is used to treat adults with the following conditions:

- **Active psoriatic arthritis** - if you cannot use another type of medicine called 'Disease-Modifying Antirheumatic Drugs' (DMARDs) or when you have tried one of these medicines and it did not work.
- **Moderate to severe chronic plaque psoriasis** - if you cannot use one of the following treatments or when you have tried one of these treatments and it did not work:
 - phototherapy - a treatment where certain areas of skin are exposed to ultraviolet light
 - systemic therapy - a treatment that affects the entire body rather than just one local area, such as 'ciclosporin', 'methotrexate' or 'psoralen'.
- **Behcet's disease (BD)** - to treat the mouth ulcers which is a common problem for people with this illness.

What psoriatic arthritis is

Psoriatic arthritis is an inflammatory disease of the joints, usually accompanied by psoriasis, an inflammatory disease of the skin.

What plaque psoriasis is

Psoriasis is an inflammatory disease of the skin, which can cause red, scaly, thick, itchy, painful patches on your skin and can also affect your scalp and nails.

2. What you need to know before you take Apremilast

Do not take Apremilast

- if you are allergic to apremilast or any of the other ingredients of this medicine (listed in section 6).
- if you are pregnant or think you may be pregnant.

How Apremilast works

Talk to your doctor or pharmacist before taking Apremilast.

Depression and suicidal thoughts

Tell your doctor before starting Apremilast if you have depression which is getting worse with thoughts of suicide.

You or your caregiver should also tell your doctor straight away of any changes in behaviour or mood, feelings of depression and of any suicidal thoughts you may have after taking Apremilast.

In psoriatic arthritis, treatment with Apremilast results in an improvement in swollen and painful joints, and can improve your general physical function.

In psoriasis, treatment with Apremilast results in a reduction in psoriatic skin plaques and other signs and symptoms of the disease.

In Behcet's disease, treatment with Apremilast reduces the number of mouth ulcers and can stop them completely. It can also reduce the associated pain.

Apremilast has also been shown to improve the quality of life in patients with psoriasis, psoriatic arthritis or Behcet's disease. This means that the impact of your condition on daily activities, relationships and other factors should be less than it was before.

Gut problems

If you experience severe diarrhoea, nausea, or vomiting, you should talk to your doctor.

Children and adolescents

Apremilast has not been studied in children and adolescents, therefore it is not recommended for use in children and adolescents aged 17 years and under.

Apremilast contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

3. How to take Apremilast

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

How much to take

- When you first start taking Apremilast, you will receive a 'treatment initiation pack' which contains all the doses as listed in the table below.
- The 'treatment initiation pack' is clearly labelled to make sure you take the correct tablet at the correct time.
- Your treatment will start at a lower dose and will gradually be increased over the first 6 days of treatment.
- The 'treatment initiation pack' will also contain enough tablets for another 8 days at the recommended dose (days 7 to 14).

Severe kidney problems

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

If you are underweight

Talk to your doctor while taking Apremilast if you lose weight without meaning to.

In pregnancy. You should not become pregnant while taking this medicine and should use an effective method of contraception during treatment with Apremilast.

It is not known if this medicine passes into human milk. You should not use Apremilast while breast-feeding.

Driving and using machines

Apremilast has no effect on the ability to drive and use machines.

Other medicines and Apremilast

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes medicines obtained without a prescription and herbal medicines. This is because Apremilast can affect the way some other medicines work. Also some other medicines can affect the way Apremilast works.

Warnings and precautions

In particular, tell your doctor or pharmacist before taking Apremilast if you are taking any of the following medicines:

- rifampicin – an antibiotic used for tuberculosis
- phenytoin, phenobarbital and carbamazepine - medicines used in the treatment of seizures or epilepsy
- St John's Wort – a herbal medicine for mild anxiety and depression.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

- The recommended dose of Apremilast is 30 mg twice a day after the titration phase is complete - one 30 mg dose in the morning and one 30 mg dose in the evening, approximately 12 hours apart, with or without food.
- This is a total daily dose of 60 mg. By the end of day 6 you will have reached this recommended dose.

- Once the recommended dose has been reached, you will only get the 30 mg tablet strength in your prescribed packs. You will only ever need to go through this stage of gradually increasing your dose once even if you restart treatment.

Day	Morning Dose	Evening Dose	Total Daily Dose
Day 1	10 mg (pink)	Do not take a dose	10 mg
Day 2	10 mg (pink)	10 mg (pink)	20 mg
Day 3	10 mg (pink)	20 mg (orange brown)	30 mg
Day 4	20 mg (orange brown)	20 mg (orange brown)	40 mg
Day 5	20 mg (orange brown)	30 mg (light brownish violet)	50 mg
Day 6 onwards	30 mg (light brownish violet)	30 mg (light brownish violet)	60 mg

People with severe kidney problems

If you have severe kidney problems then the

(morning dose).

Your doctor will talk to you about how

to increase your dose when you first start taking Apremilast.

Black U (20%)

Black U (40%)

Black U





If your condition has not improved after six months of treatment, you should talk to your doctor.

If you take more Apremilast than you should

If you take more Apremilast than you should, talk to a doctor or go to a hospital straight away. Take the medicine pack and this leaflet with you.

If you forget to take Apremilast

- If you miss a dose of Apremilast, take it as soon as you remember. If it is close to the time for your next dose, just skip the missed dose. Take the next dose at your regular time.
- Do not take a double dose to make up for a forgotten dose.

If you stop taking Apremilast

- You should continue taking Apremilast until your doctor tells you to stop.
- Do not stop taking Apremilast without talking to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Serious side effects – depression and suicidal thoughts

Tell your doctor straight away about any changes in behaviour or mood, feelings of depression, thoughts of suicide or suicidal behaviour (this is uncommon).

Very common side effects (may affect more than 1 in 10 people)

- diarrhoea
- nausea
- headaches
- upper respiratory tract infections such as cold, runny nose, sinus infection

Common side effects (may affect up to 1 in 10 people)

- cough
- back pain
- vomiting
- feeling tired
- stomach pain
- loss of appetite
- frequent bowel movements
- difficulty sleeping (insomnia)
- indigestion or heartburn
- inflammation and swelling of the tubes in your lungs (bronchitis)
- common cold (nasopharyngitis)
- depression
- migraine
- tension headache

Uncommon side effects (may affect up to 1 in 100 people)

- rash
- hives (urticaria)
- weight loss
- allergic reaction
- bleeding in the bowel or in the stomach
- suicidal ideation or behaviour

Not known side effects (frequency cannot be estimated from the available data):

- severe allergic reaction (may include swelling of the face, lips, mouth, tongue, or throat that may lead to difficulty breathing or swallowing)

If you are 65 years of age or older, you might have a higher risk of complications of severe diarrhoea, nausea and vomiting. If your gut problems become severe, you should talk to your doctor.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via Yellow Card Scheme

Website: yellowcard.mhra.gov.uk or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Apremilast

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the packaging after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Apremilast contains

- The active substance is apremilast. Each film-coated tablet contains 10 mg, 20 mg or 30 mg apremilast.
- The other ingredients (excipients) are:

Tablet core: mannitol (E421), microcrystalline cellulose, croscarmellose sodium (See section 2 "Apremilast contains sodium") and magnesium stearate (E470b).

Film coating: poly(vinyl alcohol), macrogol 3350, titanium dioxide (E171), talc, red iron oxide (E172), yellow iron oxide (E172) - only for 20 mg and 30 mg and black iron oxide (E172) - only for 30 mg.

What Apremilast looks like and contents of the pack

Apremilast 10 mg film-coated tablets

Film-coated tablets (tablets) are pink, round, biconvex, marked with 10 on one side of the tablet. Tablet dimension: diameter approx. 6 mm.

Apremilast 20 mg film-coated tablets

Film-coated tablets (tablets) are orange brown, round, biconvex marked with 20 on one side of the tablet. Tablet dimension: diameter approx. 8 mm.

Apremilast 30 mg film-coated tablets

Film-coated tablets (tablets) are light brownish violet, round, biconvex, marked with 30 on one side of the tablet. Tablet dimension: diameter approx. 10 mm.

Apremilast 30 mg film-coated tablets is available in:

- packs containing 14, 56 or 168 film-coated tablets, in a blister.

• Treatment initiation pack:

Each pack of 27 film-coated tablets contains:

- 4 film-coated tablets of Apremilast 10 mg

- 4 film-coated tablets of Apremilast 20 mg

- 19 film-coated tablets of Apremilast 30 mg

Not all pack sizes may be marketed.

Marketing Authorisation Holder

KRKA, d.d., Novo mesto, Šmarješka cesta 6,

8501 Novo mesto, Slovenia

Manufacturer

KRKA, d.d., Novo mesto, Šmarješka cesta 6,

8501 Novo mesto, Slovenia

TAD Pharma GmbH, Heinz-Lohmann-Straße 5,

27472 Cuxhaven, Germany

This leaflet was last revised in October 2024.

