

Apremilast 10 mg film-coated tablets

Apremilast 20 mg film-coated tablets

Apremilast 30 mg film-coated tablets

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Apremilast film-coated tablets are and what they are used for
2. What you need to know before you take Apremilast film-coated tablets
3. How to take Apremilast film-coated tablets
4. Possible side effects
5. How to store Apremilast film-coated tablets
6. Contents of the pack and other information

1. What Apremilast film-coated tablets are and what they are used for

What apremilast is

Apremilast film-coated tablets contain the active substance 'apremilast'. This belongs to a group of medicines called phosphodiesterase 4 inhibitors, which help to reduce inflammation.

What apremilast is used for

Apremilast is used to treat adults with the following conditions:

- **Active psoriatic arthritis** - if you cannot use another type of medicine called 'Disease-Modifying Antirheumatic Drugs' (DMARDs) or when you have tried one of these medicines and it did not work.
- **Moderate to severe chronic plaque psoriasis** - if you cannot use one of the following treatments or when you have tried one of these treatments and it did not work:
 - o phototherapy - a treatment where certain areas of skin are exposed to ultraviolet light
 - o systemic therapy - a treatment that affects the entire body rather than just one local area, such as 'ciclosporin', 'methotrexate' or 'psoralen'.
- **Behçet's disease (BD)** - to treat the mouth ulcers which is a common problem for people with this illness.

What psoriatic arthritis is

Psoriatic arthritis is an inflammatory disease of the joints, usually accompanied by psoriasis, an inflammatory disease of the skin.

What plaque psoriasis is

Psoriasis is an inflammatory disease of the skin, which can cause red, scaly, thick, itchy, painful patches on your skin and can also affect your scalp and nails.

What Behçet's disease is

Behçet's disease is a rare type of inflammatory disease which affects many parts of the body. The most common problem is mouth ulcers.

How apremilast works

Psoriatic arthritis, psoriasis and Behçet's disease are usually lifelong conditions and there is currently no cure. Apremilast works by reducing the activity of an enzyme in the body called 'phosphodiesterase 4', which is involved in the process of inflammation. By reducing the activity of this enzyme, Apremilast can help to control the inflammation associated with psoriatic arthritis, psoriasis and Behçet's disease, and thereby reduce the signs and symptoms of these conditions.

In psoriatic arthritis, treatment with apremilast results in an improvement in swollen and painful joints, and can improve your general physical function.

In psoriasis, treatment with apremilast results in a reduction in psoriatic skin plaques and other signs and symptoms of the disease.

In Behçet's disease, treatment with apremilast reduces the number of mouth ulcers and can stop them completely. It can also reduce the associated pain.

Apremilast has also been shown to improve the quality of life in patients with psoriasis, psoriatic arthritis or Behçet's disease. This means that the impact of your condition on daily activities, relationships and other factors should be less than it was before.

2. What you need to know before you take Apremilast film-coated tablets

DO NOT take Apremilast film-coated tablets:

- if you are allergic to apremilast or any of the other ingredients of this medicine (listed in section 6).
- if you are pregnant or think you may be pregnant.

Warnings and precautions

Talk to your doctor or pharmacist before taking this medicine.

Depression and suicidal thoughts

Tell your doctor before starting apremilast treatment if you have depression which is getting worse with thoughts of suicide.

You or your caregiver should also tell your doctor straight away of any changes in behaviour or mood, feelings of depression and of any suicidal thoughts you may have after taking this medicine.

Severe kidney problems

If you have severe kidney problems, your dose will be different – see section 3.

If you are underweight

Talk to your doctor while taking this medicine if you lose weight without meaning to.

Gut problems

If you experience severe diarrhoea, nausea, or vomiting, you should talk to your doctor.

Children and adolescents

Apremilast has not been studied in children and adolescents, therefore it is not recommended for use in children and adolescents aged 17 years and under.

Other medicines and apremilast

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes medicines obtained without a prescription and herbal medicines. This is because apremilast can affect the way some other medicines work. Also some other medicines can affect the way apremilast works.

In particular, tell your doctor or pharmacist before taking apremilast if you are taking any of the following medicines:

- rifampicin – an antibiotic used for tuberculosis
- phenytoin, phenobarbital and carbamazepine - medicines used in the treatment of seizures or epilepsy
- St John's Wort – a herbal medicine for mild anxiety and depression.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

There is little information about the effects of apremilast in pregnancy. You should not become pregnant while taking this medicine and should use an effective method of contraception during treatment with apremilast.

It is not known if this medicine passes into human milk. You should not use this medicine while breast-feeding.

Driving and using machines

Apremilast has no effect on the ability to drive and use machines.

This medicine contains lactose

Apremilast film-coated tablets contain lactose (a type of sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Apremilast film-coated tablets

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

How much to take

- When you first start taking this medicine, you will receive a 'treatment initiation pack' which contains all the doses as listed in the table below.
- The 'treatment initiation pack' is clearly labelled to make sure you take the correct tablet at the correct time.
- Your treatment will start at a lower dose and will gradually be increased over the first 6 days of treatment.

- The 'treatment initiation pack' will also contain enough tablets for another 8 days at the recommended dose (days 7 to 14).
- The recommended dose of apremilast is 30 mg twice a day after the titration phase is complete - one 30 mg dose in the morning and one 30 mg dose in the evening, approximately 12 hours apart, with or without food.
- This is a total daily dose of 60 mg. By the end of day 6 you will have reached this recommended dose.
- Once the recommended dose has been reached, you will only get the 30 mg tablet strength in your prescribed packs. You will only ever need to go through this stage of gradually increasing your dose once even if you re-start treatment.

Day	Morning Dose	Evening Dose	Total Daily Dose
Day 1	10 mg (pink)	Do not take a dose	10 mg
Day 2	10 mg (pink)	10 mg (pink)	20 mg
Day 3	10 mg (pink)	20 mg (brown)	30 mg
Day 4	20 mg (brown)	20 mg (brown)	40 mg
Day 5	20 mg (brown)	30 mg (beige)	50 mg
Day 6 onwards	30 mg (beige)	30 mg (beige)	60 mg

People with severe kidney problems

If you have severe kidney problems then the recommended dose of apremilast is 30 mg **once a day (morning dose)**. Your doctor will talk to you about how to increase your dose when you first start taking this medicine.

How and when to take Apremilast film-coated tablets

- Apremilast film-coated tablets are for oral use.
- Swallow the tablets whole, preferably with water.
- You can take the tablets either with or without food.
- Take this medicine at about the same time each day, one tablet in the morning and one tablet in the evening.

If your condition has not improved after six months of treatment, you should talk to your doctor.

If you take more of this medicine than you should

If you take more Apremilast film-coated tablets than you should, talk to a doctor or go to a hospital straight away. Take the medicine pack and this leaflet with you.

If you forget to take this medicine

- If you miss a dose of Apremilast film-coated tablets, take it as soon as you remember. If it is close to the time for your next dose, just skip the missed dose. Take the next dose at your regular time.
- Do not take a double dose to make up for a forgotten dose.

If you stop taking this medicine

- You should continue taking this medicine until your doctor tells you to stop.
- Do not stop taking this medicine without talking to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Serious side effects – depression and suicidal thoughts

Tell your doctor straight away about any changes in behaviour or mood, feelings of depression, thoughts of suicide or suicidal behaviour (this is uncommon).

Very common side effects (may affect more than 1 in 10 people)

- diarrhoea
- nausea
- headaches
- upper respiratory tract infections such as cold, runny nose, sinus infection

Common side effects (may affect up to 1 in 10 people)

- cough
- back pain
- vomiting
- feeling tired
- stomach pain
- loss of appetite
- frequent bowel movements
- difficulty sleeping (insomnia)
- indigestion or heartburn
- inflammation and swelling of the tubes in your lungs (bronchitis)
- common cold (nasopharyngitis)
- depression
- migraine
- tension headache

Uncommon side effects (may affect up to 1 in 100 people)

- rash
- hives (urticaria)
- weight loss
- allergic reaction
- bleeding in the bowel or in the stomach
- suicidal ideation or behaviour

Not known side effects (frequency cannot be estimated from the available data):

- severe allergic reaction (may include swelling of the face, lips, mouth, tongue, or throat that may lead to difficulty breathing or swallowing)

If you are 65 years of age or older, you might have a higher risk of complications of severe diarrhoea, nausea and vomiting. If your gut problems become severe, you should talk to your doctor.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Apremilast film-coated tablets

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister or on the carton after EXP. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Apremilast film-coated tablets contain

The active substance is apremilast.
Each tablet contains 10 mg, 20 mg, 30 mg apremilast.

The other ingredients are:

Tablet core: cellactose (consisting of cellulose, powdered and lactose monohydrate), calcium carbonate (consisting of calcium carbonate and starch, pregelatinised (maize starch), crospovidone type B, sodium stearyl fumarate

Film-coating 10 mg tablets: hypromellose E464, macrogol 4000 E1521, titanium dioxide E171, iron oxide yellow E172

Film-coating 20 mg tablets: hypromellose E464, macrogol 4000 E1521, titanium dioxide E171, iron oxide yellow E172, iron oxide red E172

Film-coating 30 mg tablets: hypromellose E464, titanium dioxide E171, macrogol 4000 E1521, iron oxide red E172, iron oxide yellow E172, iron oxide black E172

What Apremilast film-coated tablets look like and contents of the pack

Apremilast 10 mg film-coated tablets are oval, biconvex, pink film-coated tablets

Apremilast 20 mg film-coated tablets are oval, biconvex, brown film-coated tablets

Apremilast 30 mg film-coated tablets are oval, biconvex, beige film-coated tablets

Pack sizes

Apremilast 30 mg is available in packs containing 56 or 168 film coated tablets.

The treatment initiation pack is available in packs containing 27 film-coated tablets: 4 x 10 mg tablets, 4 x 20 mg tablets and 19 x 30 mg tablets.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

STADA, Linthwaite, Huddersfield, HD7 5QH, UK

Manufacturer

STADA Arzneimittel AG, Stadastrasse 2-18, 61118, Bad Vilbel, Germany

This leaflet was last revised in 03/2025