

Difflam 3 mg Lozenges, lemon flavour
Benzydamine Hydrochloride

Always take this medicine exactly as described in this leaflet or as your doctor, or pharmacist have told you

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must talk to a doctor if you do not feel better or if you feel worse after 3 days.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet, see section 4.

1. What Diffiam Lozenges is and what it is used for
2. What you need to know before you take Diffiam Lozenges
3. How to take Diffiam Lozenges
4. Possible side effects
5. How to store Diffiam Lozenges
6. Contents of the pack and other information

You must talk to a doctor if you do not feel better or if you feel worse after 3 days.

- If you are allergic to benzydamine or any of the other ingredients of this medicine (listed in section 6).
- If you suffer from a condition known as phenylketonuria.
- If you are allergic to salicylic acid or to other anti-inflammatory painkillers called NSAIDs (e.g. ibuprofen).

- If you have a history of Asthma.
- If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.
- If after using the lozenges the pain in your mouth or throat gets worse or does not improve within 3 days, or you appear feverish or have other symptoms, you should speak to your doctor or pharmacist.

Food and drink have no influence.

Allergic reactions are rare, but can be serious. Tell your doctor immediately if you get any sudden swelling in



