




Sodium

Dazublys contains less than 1 mmol of sodium (23 mg) per dose, that is to say essentially sodium-free.

Polysorbate 20

This medicine contains 0.6 mg of polysorbate 20 in each 150 mg/vial (21 mg/mL) which is equivalent to 0.6 mg/60 kg/7.4 mL. Polysorbates may cause allergic reactions. Tell your doctor if you have any known allergies.

3. How Dazublys is given

Before starting the treatment, your doctor will determine the amount of HER2 in your tumor. Only patients with a large amount of HER2 will be treated with Dazublys. Dazublys should only be given by a doctor or nurse. Your doctor will prescribe a dose and treatment regimen that is right for you. The dose of Dazublys depends on your body weight.

The first dose of your treatment is given over 90 minutes, and you will be observed by a health professional while it is being given in case you have any side effects. If the first dose is well tolerated, the next doses may be given over 30 minutes (see section 2 under "Warnings and precautions"). The number of infusions you receive will depend on how you respond to the treatment. Your doctor will discuss this with you.

Dazublys is given as an infusion into a vein (intravenous infusion, drip); this intravenous formulation is not for subcutaneous use and should be given as an intravenous infusion only.

In order to prevent medication errors, it is important to check the vial labels to ensure that the medicine is prepared and given is Dazublys (trastuzumab) and not another trastuzumab-containing product (e.g., trastuzumab emtansine or trastuzumab deruxtecan).

For early breast cancer, metastatic breast cancer, and metastatic gastric cancer, Dazublys is given every three weeks. Dazublys may also be given once a week for metastatic breast cancer.

If you stop using Dazublys

Do not stop using this medicine without talking to your doctor first. All doses should be taken at the right time every week or every three weeks (depending on your dosing schedule). This helps your medicine work as well as it can.

It may take up to 7 months for Dazublys to be removed from your body. Therefore, your doctor may decide to continue to check your heart functions even after you finish treatment.

If you have any further questions about the use of this medicine, ask your doctor, pharmacist, or nurse.

4. Possible side effects

Like all medicines, trastuzumab can cause side effects, although not everybody gets them. Some of these side effects may be serious and may lead to hospitalization.

During a Dazublys infusion, chills, fever, and other flu-like symptoms may occur. These are very common (may affect more than 1 in 10 people). Other infusion-related symptoms are feeling sick (nausea), vomiting, pain, increased muscle tension and shaking, headache, dizziness, breathing difficulties, high or low blood pressure, heart rhythm disturbances (palpitations, heart fluttering, or irregular heartbeat), swelling of the face and lips, rash and feeling tired. Some of these symptoms can be serious, and some patients have died (see section 2 under "Warnings and precautions").

These effects mainly occur with the first intravenous infusion ("drip" into your vein) and during the first few hours after the start of the infusion. They are usually temporary. You will be observed by a healthcare professional during the infusion and for at least six hours after the start of the first infusion and for two hours after the start of other infusions. If you develop a reaction, they will slow down or stop the infusion and may give you treatment to counteract the side effects. The infusion may be continued after the symptoms improve.

Occasionally, symptoms start later than six hours after the infusion begins. If this happens to you, contact your doctor immediately. Sometimes, symptoms may improve and then get worse later.

Serious side effects

Other side effects can occur at any time during treatment with Dazublys, not just related to an infusion.

Tell a doctor or nurse straight away if you notice any of the following side effects:

- Heart problems can sometimes occur during treatment and occasionally after treatment has stopped and can be serious. They include weakening of the heart muscle, possibly leading to heart failure, inflammation of the lining around the heart, and heart rhythm disturbances. This can lead to symptoms such as breathlessness (including breathlessness at night), cough, fluid retention (swelling) in the legs or arms, and palpitations (heart fluttering or irregular heartbeat) (see section 2. Heart checks).

Your doctor will monitor your heart regularly during and after treatment, but you should tell your doctor immediately if you notice any of the above symptoms.

- Tumour lysis syndrome (a group of metabolic complications occurring after cancer treatment characterized by high blood levels of potassium and phosphate and low blood levels of calcium). Symptoms may include kidney problems (weakness, shortness of breath, fatigue, and confusion), heart problems (fluttering of the heart or a faster or slower heartbeat), seizures, vomiting or diarrhoea, and tingling in the mouth, hands, or feet.

If you experience any of the above symptoms when your treatment with Dazublys has finished, you should see your doctor and tell them that you have previously been treated with Dazublys.

Very common side effects of Trastuzumab: may affect more than 1 in 10 people

- infections
- diarrhoea
- constipation
- heartburn (dyspepsia)
- fatigue
- skin rashes
- chest pain
- abdominal pain
- joint pain
- low counts of red blood cells and white blood cells (which help fight infection), sometimes with fever
- muscle pain
- conjunctivitis
- watery eyes
- nose bleeds
- runny nose
- hair loss
- tremor
- hot flush
- dizziness
- nail disorders
- weight loss
- loss of appetite
- inability to sleep (insomnia)
- altered taste
- low platelet count
- bruising
- numbness or tingling of the fingers and toes, which occasionally may extend to the rest of the limb
- redness, swelling, or sores in your mouth and/or throat
- pain, swelling, redness, or tingling of hands and/or feet
- breathlessness
- headache
- cough
- vomiting
- nausea

Common side effects of Trastuzumab: may affect up to 1 in 10 people

- allergic reactions
- throat infections
- bladder and skin infections
- inflammation of the breast
- inflammation of the liver
- kidney disorders
- increased muscle tone or tension (hypertonia)
- pain in the arms and/or legs
- itchy rash
- sleepiness (somnolence)
- haemorrhoids
- itchiness
- dry mouth and skin
- dry eyes
- sweating
- feeling weak and unwell
- anxiety
- depression
- asthma
- infection of the lungs
- lung disorders
- back pain
- neck pain
- bone pain
- acne
- leg cramps

Uncommon side effects of Trastuzumab: may affect up to 1 in 100 people:

- deafness
- bumpy rash
- wheezing
- inflammation or scarring of the lungs

Rare side effects of Trastuzumab: may affect up to 1 in 1000 people

- jaundice
- anaphylactic reactions

Other side effects that have been reported with Trastuzumab use:

- frequency cannot be estimated from the available data
- abnormal or impaired blood clotting
- high potassium levels
- swelling or bleeding at the back of the eyes
- shock
- abnormal heart rhythm
- respiratory distress
- respiratory failure
- acute accumulation of fluid in the lungs
- acute narrowing of the airways
- abnormally low oxygen levels in the blood
- difficulty in breathing when lying flat
- liver damage
- swelling of the face, lips and throat
- kidney failure
- abnormally low levels of fluid around the baby in the womb
- failure of the lungs of the baby to develop in the womb
- abnormal development of the kidneys of the baby in the womb

Some of the side effects you experience may be due to your underlying cancer. If you receive Dazublys in combination with chemotherapy, some of them may also be due to the chemotherapy.

If you get any side effects, talk to your doctor, pharmacist, or nurse.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist, or nurse. This includes any possible side effects not listed in this leaflet. Healthcare professionals are asked to report any suspected adverse reactions (see details below). By reporting side effects you can help provide more information on the safety of this medicine.

United Kingdom

Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store

5. How to store Dazublys

Dazublys will be stored by the health professionals at the hospital or clinic.

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiration date, which is stated on the outer carton and on the vial label after EXP. The expiry date refers to the last day of that month.
- The unopened vial should be stored in a refrigerator (2 °C - 8 °C).
- Do not freeze the reconstituted solution.
- Infusion solutions should be used immediately after dilution. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would not normally be longer than 48 hours at 2 °C - 8 °C.
- Do not use Dazublys if you notice any particulate matter or discoloration prior to administration.
- Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines that are no longer required. These measures will help to protect the environment. Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

6. Contents of the pack and other information What Dazublys contains.

- The active substance is trastuzumab. Each vial contains 150 mg trastuzumab that has to be dissolved in 7.2 mL of water for injection. The resulting solution contains approximately 21 mg/mL of trastuzumab.
- The other ingredient(s) are L-histidine hydrochloride monohydrate, L-histidine, α,α-trehalose dihydrate, and polysorbate 20 (E 432).
- This medicine contains 0.6 mg of polysorbate 20 in each 150 mg /vial (21 mg/mL) which is equivalent to 0.6 mg/60 kg/7.4 mL. Polysorbates may cause allergic reactions. Tell your doctor if you have any known allergies.

What Trastuzumab looks like and the contents of the pack

Trastuzumab is a powder for concentration solution for intravenous infusion, which is supplied in a glass vial with a rubber stopper containing 150 mg of trastuzumab. The powder is a white to pale yellow pellet. Each carton contains one vial of powder.

Marketing Authorization Holder

CuraTeQ Biologics s.r.o
Trtinova 260/1, Čakovice,
19600, Prague,
Czech Republic

Manufacturer

APL Swift Services Malta Ltd.
HF26, Hal Far Industrial Estate,
Qasam Industrijali Hal Far,
Birzebbugia, BBG 3000,
Malta.

Distributed By:

Milpharm Limited
Ares.Odyssey Business Park,
West End Road,Ruislip,Middlesex,
HA4 6QD,United Kingdom.

This leaflet was last revised in 10/2025

Detailed information on this medicine is available on the Medicines and Healthcare products Regulatory Agency website:
<https://products.mhra.gov.uk>

The following information is intended for medical or healthcare professionals only

Dazublys intravenous is provided in sterile, preservative-free, non-pyrogenic, single use vials.

In order to prevent medication errors, it is important to check the vial labels to ensure that the medicine being prepared and given is Dazublys and not another trastuzumab-containing product (e.g. trastuzumab emtansine or trastuzumab deruxtecan).

Always keep this medicine in the closed original pack at a temperature of 2 °C –8 °C in a refrigerator.

Appropriate aseptic technique should be used for reconstitution and dilution procedures. Care must be taken to ensure the sterility of prepared solutions. Since the medicinal product does not contain any anti-microbial preservative or bacteriostatic agents, aseptic technique must be observed.

A vial of Dazublys aseptically reconstituted with sterile water for injections (not supplied) is chemically and physically stable for 48 hours at 2 °C - 8 °C after reconstitution and must not be frozen.

After aseptic dilution in polyvinylchloride, polyethylene, or polypropylene bags containing sodium chloride 9 mg/mL (0.9%) solution for injection, chemical and physical stability of Dazublys has been demonstrated for up to 48 hours at 5 °C ± 3 °C and 30 °C ± 2 °C.

From a microbiological point of view, the reconstituted solution and trastuzumab infusion solution should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would not normally be longer than 24 hours at 2 °C to 8 °C, unless reconstitution and dilution have taken place under controlled and validated aseptic conditions.

Aseptic preparation, handling, and storage:

Aseptic handling must be ensured when preparing the infusion. Preparation should be:

- performed under aseptic conditions by trained personnel in accordance with good practice rules especially with respect to the aseptic preparation of parenteral products.
- prepared in a laminar flow hood or biological safety cabinet using standard precautions for the safe handling of intravenous agents.
- followed by adequate storage of the prepared solution for intravenous infusion to ensure maintenance of the aseptic conditions.

Each vial of Dazublys is reconstituted with 7.2 mL of water for injections (not supplied). Use of other reconstitution solvents should be avoided. This yields a 7.4 mL solution for single-dose use, containing approximately 21 mg/mL trastuzumab. A volume overage of 4 % ensures that the labelled dose of 150 mg can be withdrawn from each vial.

Dazublys should be carefully handled during reconstitution. Causing excessive foaming during reconstitution or shaking the reconstituted trastuzumab may result in problems with the amount of trastuzumab that can be withdrawn from the vial.

Instructions for reconstitution:

- Using a sterile syringe, slowly inject 7.2 mL of water for injections in the vial containing the lyophilized Dazublys, directing the stream into the lyophilized cake.
- Swirl vial gently to aid reconstitution. DO NOT SHAKE!

Slight foaming of the product upon reconstitution is not unusual. Allow the vial to stand undisturbed for approximately 5 minutes. The reconstituted Dazublys results in a colourless to pale yellow transparent solution and should be essentially free of visible particulates.

Determine the volume of the solution required:

- based on a loading dose of 8 mg trastuzumab/kg body weight, or a subsequent weekly dose of 2 mg trastuzumab/kg body weight:

$$\text{Volume (mL)} = \frac{\text{Body weight (kg)} \times \text{dose (4 mg/kg for loading or 2 mg/kg for maintenance)}}{21 \text{ (mg/mL, concentration of reconstituted solution)}}$$

- based on a loading dose of 8 mg trastuzumab/kg body weight, or a subsequent 3-weekly dose of 6 mg trastuzumab/kg body weight:

$$\text{Volume (mL)} = \frac{\text{Body weight (kg)} \times \text{dose (8 mg/kg for loading or 6 mg/kg for maintenance)}}{21 \text{ (mg/mL, concentration of reconstituted solution)}}$$

The appropriate amount of solution should be withdrawn from the vial using a sterile needle and syringe and added to a polyvinylchloride, polyethylene, or polypropylene bags containing 250 mL of 0.9% sodium chloride solution. Do not use with glucose-containing solutions. The bag should be gently inverted to mix the solution to avoid foaming. Parenteral solutions should be inspected visually for particulates and discoloration prior to administration. Once the infusion is prepared it should be administered immediately. If diluted aseptically, it may be stored for 48 hours at 5 °C ± 3 °C and 30 °C ± 2 °C.