



Package Leaflet: Information for the user
STRIVIT-D3 3,200 IU capsules, soft
Colecalciferol

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What STRIVIT-D3 is and what it is used for
2. What you need to know before you take STRIVIT-D3
3. How to take STRIVIT-D3
4. Possible side effects
5. How to store STRIVIT-D3
6. Contents of the pack and other information

1. What STRIVIT-D3 is and what it is used for

STRIVIT-D3 contains the active substance Colecalciferol. Colecalciferol is a form of vitamin D.

Vitamin D is found in the diet and is also produced in the skin after exposure to the sun. Often vitamin D is given in combination with calcium. STRIVIT-D3 is recommended for use when you have a normal intake of dietary calcium.

STRIVIT-D3 may be prescribed by your doctor to treat or prevent vitamin D deficiency. Deficiency of vitamin D may occur when your diet or lifestyle does not provide you enough vitamin D or when your body requires more vitamin D (for instance when you are pregnant).

2. What you need to know before you take STRIVIT-D3

Do not take STRIVIT-D3:

- if you are allergic (hypersensitive) to vitamin D or any of the other ingredients of this medicine (listed in section 6);
- if you have high levels of vitamin D in your blood (hypervitaminosis D);
- if you have high blood levels of calcium (hypercalcaemia) or high urine levels of calcium (hypercalciuria);
- if you have kidney stones or serious kidney problems.

If any of the above applies to you, do not take STRIVIT-D3.

Warnings and precautions

Talk to your doctor or pharmacist before taking STRIVIT-D3:

- if you have kidney damage or disease and long term treatment with this medicine. Your doctor may want to measure the levels of calcium in your blood or urine;
- if you are being treated for heart disease;
- you have sarcoidosis (an immune system disorder which may affect your liver, lungs, skin or lymph nodes);
- if you are already taking additional doses of calcium or vitamin D. Whilst you are taking STRIVIT-D3 your doctor will monitor your blood levels of calcium to make sure they are not too high.

Children and adolescents

STRIVIT-D3 is not suitable for children under 12 years.

Other medicines and STRIVIT-D3

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

In particular the following medicines may interact with STRIVIT-D3:

- **Heart medicines** (cardiac glycosides such as digoxin). Your doctor may monitor your heart with an electrocardiogram (ECG) and measure the levels of calcium in your blood.
- **Thiazide diuretics** (used to treat high blood pressure) reduce the urinary excretion of calcium and can increase risk of hypercalcaemia.
- **Medicines used to treat epilepsy** (such as phenytoin) or medicines to make you sleep (barbiturates such as phenobarbitone) can decrease the effects of vitamin D.
- **Glucocorticoids** (steroid hormones such as hydrocortisone or prednisolone) can decrease the effects of vitamin D.
- **Laxatives** (such as paraffin oil), or a cholesterol lowering drug (colestyramine, colestipol), orlistat may reduce the absorption of vitamin D.
- **Actinomycin** (a medicine used to treat some forms of cancer) and imidazole antifungals (medicines such as clotrimazole and ketoconazole used to treat fungal diseases) may interfere with the metabolism of vitamin D.

STRIVIT-D3 with food and drink

You can take STRIVIT-D3 with or without food and drink.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. If you require extra vitamin D, do not take it without the supervision of your doctor. Too much vitamin D may harm your baby.

Driving and using machines

STRIVIT-D3 should not affect your ability to drive or operate machinery.



Dimension - 140 x 240 mm

Font: Times New Roman 9 pt

Black

Front Side

3. How to take STRIVIT-D3

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The capsules should be swallowed whole with water and can be taken with or without food.

The recommended dose is:

In severe vitamin D deficiency (for adults and the elderly)

Your doctor will usually prescribe you a dose of 1 capsule daily for up to 12 weeks. The amount will depend on how low your vitamin D levels are and how you respond to treatment.

Use in children

STRIVIT-D3 is not suitable for children under 12 years.

If you take more STRIVIT-D3 than you should

If you accidentally take one capsule too many, nothing is likely to happen. If you accidentally take several capsules too many, tell your doctor or get other medical advice immediately. If possible, take the capsules, the box and this leaflet with you to show the doctor. If you take too many capsules, you may feel or be sick, become constipated or have stomach pains, weak muscles, tiredness, lack of appetite, kidney problems and in severe cases irregular heartbeats.

If you forget to take STRIVIT-D3

If you forget to take your capsules, take them as soon as you can. After that, take the next capsule in accordance with the instructions given to you by your doctor.

Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Side effects with STRIVIT-D3 may include:

Uncommon (may affect up to 1 in 100 people)

- too much calcium in your blood (hypercalcaemia). You may feel or be sick, lose your appetite, have constipation, stomach ache, feel very thirsty, have muscle weakness, drowsiness or confusion;
- too much calcium in your urine (hypercalciuria).

Rare (may affect up to 1 in 1,000 people)

- skin rash;
- itching;
- hives.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can

also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store STRIVIT-D3

Keep this medicine out of the sight and reach of children.

This medicine does not require any special temperature storage conditions. Keep the blister in the outer carton in order to protect from light.

Do not use this medicine after the expiry date which is stated on the carton after "EXP". The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What STRIVIT-D3 contains

- The active ingredient is colecalciferol. Each capsule contains 3,200 IU colecalciferol (equivalent to 80 micrograms Vitamin D3).
- The other ingredients are:

Capsule Content: maize oil, refined

Capsule Shell: gelatin, glycerol (E 422), chlorophyllin copper complex sodium (E141), purified water

What STRIVIT-D3 looks like and contents of the pack


STRIVIT-D3 3,200 IU Capsule is a translucent green to greenish yellow coloured and has no markings.

It is available in blister packs containing 7, 10, 14, 20, 28, 30, 56, 60, 84 or 90 capsules. Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Strides Pharma UK Ltd.
Unit 4, Metro Centre, Tolpits Lane,
Watford, Hertfordshire WD18 9SS,
United Kingdom

This leaflet was last revised in 03/2019.



1XXXXXX

