

PACKAGE LEAFLET: INFORMATION FOR THE USER

Bortezomib 1mg powder for solution for injection

Bortezomib

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Bortezomib is and what it is used for
2. What you need to know before you use Bortezomib
3. How to use Bortezomib
4. Possible side effects
5. How to store Bortezomib
6. Contents of the pack and other information

1. WHAT BORTEZOMIB IS AND WHAT IT IS USED FOR

This medicine contains the active substance bortezomib, a so-called 'proteasome inhibitor'. Proteasomes play an important role in controlling cell function and growth. By interfering with their function, bortezomib can kill cancer cells.

Bortezomib is used for the treatment of multiple myeloma (a cancer of the bone marrow) in patients older than 18 years:

- alone or together with the medicines pegylated liposomal doxorubicin or dexamethasone, for patients whose disease is worsening (progressive) after receiving at least one prior treatment and for whom blood stem cell transplantation was not successful or is unsuitable.
- in combination with the medicines melphalan and prednisone, for patients whose disease has not been previously treated and are unsuitable for high-dose chemotherapy with blood stem cell transplantation.
- in combination with the medicines dexamethasone or dexamethasone together with thalidomide, for patients whose disease has not been previously treated and before receiving high-dose chemotherapy with blood stem cell transplantation (induction treatment).

Bortezomib is used for the treatment of mantle cell lymphoma (a type of cancer affecting the lymph nodes) in patients 18 years or older in combination with the medicines rituximab, cyclophosphamide, doxorubicin and prednisone, for patients whose disease has not been previously treated and for whom blood stem cell transplantation is unsuitable.

2. WHAT YOU NEED TO KNOW BEFORE YOU USE BORTEZOMIB

Do not use this medicine

- if you are allergic to bortezomib, boron or to any of the other ingredients of this medicine (listed in section 6)
- if you have certain severe lung or heart problems.

Warnings and precautions

Talk to your doctor or pharmacist before using Bortezomib, if you have any of the following:

- low numbers of red or white blood cells
- bleeding problems and/or low number of platelets in your blood
- diarrhoea, constipation, nausea or vomiting
- fainting, dizziness or light-headedness in the past
- kidney problems
- moderate to severe liver problems
- numbness, tingling, or pain in the hands or feet (neuropathy) in the past
- heart or blood pressure problems
- shortness of breath or cough
- seizures
- shingles (localised including around the eyes or spread across the body)
- symptoms of tumour lysis syndrome such as muscle cramping, muscle weakness, confusion, visual loss or disturbances and shortness of breath
- memory loss, trouble thinking, difficulty with walking or loss of vision. These may be signs of a serious brain infection and your doctor may suggest further testing and follow-up.

You will have to take regular blood tests before and during your treatment with Bortezomib, to check your blood cell counts regularly.

If you have mantle cell lymphoma and are given the medicine rituximab with Bortezomib you should tell your doctor:

- if you think you have a hepatitis infection now or have had one in the past. In a few cases, patients who have had hepatitis B might have a repeated attack of hepatitis, which can be fatal. If you have a history of hepatitis B infection you will be carefully checked by your doctor for signs of active hepatitis B.

You must read the package leaflets of all medicines to be taken in combination with Bortezomib for information related to these medicines before starting treatment with Bortezomib. When thalidomide is used, particular attention to pregnancy testing and prevention requirements is needed (see Pregnancy and breast-feeding in this section).

Children and adolescents

Bortezomib should not be used in children and adolescents because it is not known how the medicine will affect them.

Other medicines and Bortezomib

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines, including medicines obtained without a prescription.

In particular, tell your doctor if you are using medicines containing any of the following active substances:

- ketoconazole, used to treat fungal infections
- ritonavir, used to treat HIV infection
- rifampicin, an antibiotic used to treat bacterial infections
- carbamazepine, phenytoin or phenobarbital used to treat epilepsy
- St. John's Wort (*Hypericum perforatum*), used for depression or other conditions
- oral antidiabetics

Pregnancy and breast-feeding

You should not use Bortezomib if you are pregnant, unless clearly necessary.

Both men and women receiving Bortezomib must use effective contraception during and for up to 3 months after treatment. If, despite these measures, pregnancy occurs, tell your doctor immediately.

You should not breast-feed while using Bortezomib. Discuss with your doctor when it is safe to restart breast-feeding after finishing your treatment.

Thalidomide causes birth defects and foetal death. When Bortezomib is given in combination with thalidomide you must follow the pregnancy prevention programme for thalidomide (see package leaflet for thalidomide).

Driving and using machines

Bortezomib might cause tiredness, dizziness, fainting, or blurred vision. Do not drive or operate tools or machines if you experience such side effects; even if you do not, you should still be cautious.

3. HOW TO USE BORTEZOMIB

Your doctor will work out your dose of bortezomib according to your height and weight (body surface area). The usual starting dose of Bortezomib is 1.3mg/m² body surface area twice a week. Your doctor may change the dose and total number of treatment cycles, depending on your response to the treatment, on the occurrence of certain side effects and on your underlying conditions (e.g. liver problems).

Progressive multiple myeloma

When Bortezomib is given alone, you will receive 4 doses of Bortezomib intravenously on days 1, 4, 8 and 11, followed by a 10-day 'rest period' without treatment. This 21-day period (3 weeks) corresponds to one treatment cycle. You might receive up to 8 cycles (24 weeks).

You may also be given Bortezomib together with the medicines pegylated liposomal doxorubicin or dexamethasone.

When Bortezomib is given together with pegylated liposomal doxorubicin, you will receive Bortezomib intravenously as a 21-day treatment cycle and pegylated liposomal doxorubicin 30mg/m² is given on day 4 of the Bortezomib 21-day treatment cycle as an intravenous infusion after the Bortezomib injection.

You might receive up to 8 cycles (24 weeks).

When Bortezomib is given together with dexamethasone, you will receive Bortezomib intravenously as a 21-day treatment cycle and dexamethasone 20mg is given orally on days 1, 2, 4, 5, 8, 9, 11, and 12 of the Bortezomib 21-day treatment cycle.

You might receive up to 8 cycles (24 weeks).



The following information is intended for healthcare professionals only:

1 RECONSTITUTION FOR INTRAVENOUS INJECTION

Note: Bortezomib is a cytotoxic agent. Therefore, caution should be used during handling and preparation. Use of gloves, eye protection and other protective clothing to prevent skin contact is recommended.

Pregnant personnel should not handle this medicine.

ASEPTIC TECHNIQUE MUST BE STRICTLY OBSERVED THROUGHOUT HANDLING OF BORTEZOMIB SINCE NO PRESERVATIVE IS PRESENT.

1.1 **Preparation of the 1.0mg vial: carefully add 1.0ml** of sterile, 9mg/ml (0.9%) sodium chloride solution for injection to the vial containing the Bortezomib powder by using a 1 ml syringe without removing the vial stopper. Dissolution of the lyophilised powder is completed in less than 2 minutes.

The concentration of the resulting solution will be 1mg/ml. The solution will be clear and colourless, with a final pH of 4 to 7. You do not need to check the pH of the solution.

1.2 Before administration, visually inspect the solution for particulate matter and discoloration. If any discoloration or particulate matter is observed, the solution should be discarded. Confirm the concentration on the vial to ensure that the correct dose is being given for the **intravenous route** of administration (1mg/ml).

1.3 The reconstituted solution is preservative free and should be used immediately after preparation. However, the chemical and physical in-use stability has been demonstrated for 8 days at 25°C or 15 days at 5 ± 3°C, in the dark, when stored in the original vial and/or a syringe. The total storage time for the reconstituted medicinal product should not exceed 8 or 15 days, depending on storage temperature, prior to administration. If the reconstituted solution is not used immediately, in-use storage times and conditions prior to use are the responsibility of the user.

2 ADMINISTRATION

- Once dissolved, withdraw the appropriate amount of the reconstituted solution according to calculated dose based upon the patient's Body Surface Area.
- Confirm the dose and concentration in the syringe prior to use (check that the syringe is marked as intravenous administration).
- Inject the solution as a three to five second bolus intravenous injection through a peripheral or central intravenous catheter into a vein.
- Flush the peripheral or intravenous catheter with sterile, 9mg/ml (0.9%) sodium chloride solution.

Bortezomib 1mg powder for solution for injection IS FOR INTRAVENOUS USE ONLY. Do not give by other routes. Intrathecal administration has resulted in death.

3 DISPOSAL

A vial is for single use only and the remaining solution must be discarded.

Any unused product or waste material should be disposed of in accordance with local requirements.

Previously untreated multiple myeloma

If you have not been treated before for multiple myeloma, and **you are not** suitable for blood stem cell transplantation you will receive Bortezomib together with two other medicines, melphalan and prednisone.

In this case, the duration of a treatment cycle is 42 days (6 weeks). You will receive 9 cycles (54 weeks).

- In cycles 1 to 4, Bortezomib is administered twice weekly on days 1, 4, 8, 11, 22, 25, 29 and 32.

- In cycles 5 to 9, Bortezomib is administered once weekly on days 1, 8, 22 and 29.

Melphalan (9mg/m²) and prednisone (60mg/m²) are both given orally on days 1, 2, 3 and 4 of the first week of each cycle.

If you have not been treated before for multiple myeloma, and **you are** suitable for blood stem cell transplantation you will receive Bortezomib intravenously together with the medicines dexamethasone, or dexamethasone and thalidomide, as induction treatment.

When Bortezomib is given together with dexamethasone, you will receive Bortezomib intravenously as a 21-day treatment cycle and dexamethasone 40mg is given orally on days 1, 2, 3, 4, 8, 9, 10 and 11 of the Bortezomib 21-day treatment cycle.

You will receive 4 cycles (12 weeks).

When Bortezomib is given together with thalidomide and dexamethasone, the duration of a treatment cycle is 28 days (4 weeks).

Dexamethasone 40mg is given orally on days 1, 2, 3, 4, 8, 9, 10 and 11 of the Bortezomib 28-day treatment cycle and thalidomide is given orally daily at 50mg up to day 14 of the first cycle, and if tolerated, the thalidomide dose is increased to 100mg on days 15-28 and may be further increased to 200mg daily from the second cycle onwards.

You might receive up to 6 cycles (24 weeks).

Previously untreated mantle cell lymphoma

If you have not been treated before for mantle cell lymphoma you will receive Bortezomib intravenously together with the medicines rituximab, cyclophosphamide, doxorubicin and prednisone.

Bortezomib is given intravenously on days 1, 4, 8 and 11, followed by a 'rest period' without treatment. The duration of a treatment cycle is 21 days (3 weeks). You might receive up to 8 cycles (24 weeks).

The following medicines are given on day 1 of each Bortezomib 21-day treatment cycle as intravenous infusions:

Rituximab at 375mg/m², cyclophosphamide at 750mg/m² and doxorubicin at 50mg/m².

Prednisone is given orally at 100mg/m² on days 1, 2, 3, 4 and 5 of the Bortezomib treatment cycle.

How Bortezomib is given

This medicine is for intravenous use. Bortezomib will be administered by a health care professional experienced in the use of cytotoxic medicines.

Bortezomib powder has to be dissolved before administration. This will be done by a healthcare professional. The resulting solution is then injected into a vein rapidly, over 3 to 5 seconds.

If you are given too much Bortezomib

As this medicine is being given by your doctor or nurse, it is unlikely that you will be given too much. In the unlikely event of an overdose, your doctor will monitor you for side effects.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them. Some of these effects may be serious.

If you are given Bortezomib for multiple myeloma or mantle cell lymphoma, tell your doctor straight away if you notice any of the following symptoms:

- muscle cramping, muscle weakness
- confusion, visual loss or disturbances, blindness, seizures, headaches
- shortness of breath, swelling of your feet or changes in your heartbeat, high blood pressure, tiredness, fainting
- coughing and breathing difficulties or tightness in the chest.

Treatment with Bortezomib can very commonly cause a decrease in the numbers of red and white blood cells and platelets in your blood. Therefore, you will have to take regular blood tests before and during your treatment with Bortezomib, to check your blood cell counts regularly. You may experience a reduction in the number of:

- platelets, which may make you be more prone to bruising, or to bleeding without obvious injury (e.g. bleeding from your bowels, stomach, mouth and gums, bleeding in the brain or bleeding from the liver)
- red blood cells, which can cause anaemia, with symptoms such as tiredness and paleness
- white blood cells, which may make you more prone to infections or flu-like symptoms

If you are given Bortezomib for the treatment of multiple myeloma the side effects you may get are listed below:

Very common side effects

(may affect more than 1 in 10 people)

- Sensitivity, numbness, tingling or burning sensation of the skin or pain in the hands or feet, due to nerve damage
- Reduction in the number of red blood cells and/or white blood cells (see above)
- Fever
- Feeling sick (nausea) or vomiting, loss of appetite
- Constipation, with or without bloating (can be severe)
- Diarrhoea: if this happens, it is important that you drink more water than usual. Your doctor may give you another medicine to control diarrhoea
- Tiredness (fatigue), feeling weak
- Muscle pain, bone pain

Common side effects

(may affect up to 1 in 10 people)

- Low blood pressure, sudden fall of blood pressure on standing which may lead to fainting
- High blood pressure
- Reduced functioning of your kidneys
- Headache
- General ill feeling, pain, vertigo, light-headedness, a feeling of weakness or loss of consciousness
- Shivering
- Infections, including pneumonia, respiratory infections, bronchitis, fungal infections, coughing with phlegm, flu like illness
- Shingles (localised including around the eyes or spread across the body)
- Chest pains or shortness of breath with exercise
- Different types of rash
- Itching of the skin, lumps on the skin or dry skin
- Facial flushing or tiny broken capillaries
- Redness of the skin
- Dehydration
- Heartburn, bloating, belching, wind, stomach pain, bleeding from your bowels or stomach
- Alteration of liver functioning
- A sore mouth or lips, dry mouth, mouth ulcers or throat pain
- Weight loss, loss of taste
- Muscle cramps, muscle spasms, muscle weakness, pain in your limbs
- Blurred vision
- Infection of the outermost layer of the eye and the inner surface of the eyelids (conjunctivitis)
- Nose bleeds
- Difficulty or problems in sleeping, sweating, anxiety, mood swings, depressed mood, restlessness or agitation, changes in your mental status, disorientation
- Swelling of body, including around the eyes and other parts of the body

Uncommon side effects

(may affect up to 1 in 100 people)

- Heart failure, heart attack, chest pain, chest discomfort, increased or reduced heart rate
- Failing of your kidneys
- Inflammation of a vein, blood clots in your veins and lungs
- Problems with blood clotting
- Insufficient circulation
- Inflammation of the lining around your heart or fluid around your heart

- Infections including urinary tract infections, the flu, herpes virus infections, ear infection and cellulitis
- Bloody stools, or bleeding from mucosal membranes, e.g. mouth, vagina
- Cerebrovascular disorders
- Paralysis, seizures, falling, movement disorders, abnormal or change in, or reduced sensation (feeling, hearing, tasting, smelling), attention disturbance, trembling, twitching
- Arthritis, including inflammation of the joints in the fingers, toes, and the jaw
- Disorders that affect your lungs, preventing your body from getting enough oxygen. Some of these include: difficulty breathing; shortness of breath; shortness of breath without exercise; breathing that becomes shallow, difficult or stops; wheezing
- Hiccups, speech disorders
- Increased or decreased urine production (due to kidney damage), painful passing of urine or blood/proteins in the urine, fluid retention
- Altered levels of consciousness, confusion, memory impairment or loss
- Hypersensitivity
- Hearing loss, deafness or ringing in the ears, ear discomfort
- Hormone abnormality which may affect salt and water absorption
- Overactive thyroid gland
- Inability to produce enough insulin or resistance to normal levels of insulin
- Irritated or inflamed eyes, excessively wet eyes, painful eyes, dry eyes, eye infections, lump in the eyelid (chalazion) and red and swollen eyelids, discharge from the eyes, abnormal vision, bleeding of the eye
- Swelling of your lymph glands
- Joint or muscle stiffness, sense of heaviness, pain in your groin
- Hair loss and abnormal hair texture
- Allergic reactions
- Redness or pain at the injection site
- Mouth pain
- Infections or inflammation of the mouth, mouth ulcers, oesophagus, stomach and intestines, sometimes associated with pain or bleeding, poor movement of the intestines (including blockage), abdominal or oesophageal discomfort, difficulty swallowing, vomiting of blood
- Skin infections
- Bacterial and viral infections
- Tooth infection
- Inflammation of the pancreas, obstruction of the bile duct
- Genital pain, problem having an erection
- Weight increase
- Thirst
- Hepatitis
- Injection site or injection device related disorders
- Skin reactions and disorders (which may be severe and life threatening), skin ulcers
- Bruises, falls and injuries
- Inflammation or haemorrhage of the blood vessels that can appear as small red or purple dots (usually on the legs) to large bruise-like patches under the skin or tissue
- Benign cysts
- A severe reversible brain condition which includes seizures, high blood pressure, headaches, tiredness, confusion, blindness or other vision problems.

Rare side effects

(may affect up to 1 in 1,000 people)

- Heart problems to include heart attack, angina
- Flushing
- Discoloration of the veins
- Inflammation of the spinal nerve
- Problems with your ear, bleeding from your ear
- Underactivity of your thyroid gland
- Budd-Chiari syndrome (the clinical symptoms caused by blockage of the hepatic veins)
- Changes in or abnormal bowel function
- Bleeding in the brain
- Yellow discolouration of eyes and skin (jaundice)
- Serious allergic reaction (anaphylactic shock) signs of which may include difficulty breathing, chest pain or chest tightness, and/or feeling dizzy/faint, severe itching of the skin or raised lumps on the skin, swelling of the face, lips, tongue and /or throat, which may cause difficulty in swallowing, collapse
- Breast disorders
- Vaginal tears
- Genital swelling
- Inability to tolerate alcohol consumption
- Wasting, or loss of body mass
- Increased appetite
- Fistula
- Joint effusion
- Cysts in the lining of joints (synovial cysts)
- Fracture
- Breakdown of muscle fibres leading to other complications
- Swelling of the liver, bleeding from the liver
- Cancer of the kidney
- Psoriasis like skin condition
- Cancer of the skin
- Paleness of the skin
- Increase of platelets or plasma cells (a type of white cell) in the blood
- Abnormal reaction to blood transfusions
- Blood clot in small blood vessels (thrombotic microangiopathy)
- Partial or total loss of vision
- Decreased sex drive
- Drooling
- Bulging eyes
- Sensitivity to light
- Rapid breathing
- Rectal pain
- Gallstones
- Hernia
- Injuries
- Brittle or weak nails
- Abnormal protein deposits in your vital organs
- Coma
- Intestinal ulcers
- Multi-organ failure
- Death

If you are given Bortezomib together with other medicines for the treatment of mantle cell lymphoma the side effects you may get are listed below:

Very common side effects

(may affect more than 1 in 10 people)

- Pneumonia
- Loss of appetite
- Sensitivity, numbness, tingling or burning sensation of the skin or pain in the hands or feet, due to nerve damage
- Nausea and vomiting
- Diarrhoea
- Mouth ulcers
- Constipation
- Muscle pain, bone pain
- Hair loss and abnormal hair texture
- Tiredness, feeling weak
- Fever

Common side effects

(may affect up to 1 in 10 people)

- Shingles (localised including around the eyes or spread across the body)
- Herpes virus infections
- Bacterial and viral infections
- Respiratory infections, bronchitis, coughing with phlegm, flu like illness
- Fungal infections
- Hypersensitivity (allergic reaction)
- Inability to produce enough insulin or resistance to normal levels of insulin
- Fluid retention
- Difficulty or problems in sleeping
- Loss of consciousness
- Altered level of consciousness, confusion
- Feeling dizzy
- Increased heartbeat, high blood pressure, sweating
- Abnormal vision, blurred vision
- Heart failure, heart attack, chest pain, chest discomfort, increased or reduced heart rate
- High or low blood pressure
- Sudden fall of blood pressure upon standing which may lead to fainting
- Shortness of breath with exercise
- Cough
- Hiccups
- Ringing in the ears, ear discomfort
- Bleeding from your bowels or stomach
- Heartburn
- Stomach pain, bloating
- Difficulty swallowing
- Infection or inflammation of the stomach and intestines
- Stomach pain
- Sore mouth or lips, throat pain
- Alteration of liver function
- Itching of skin
- Redness of skin
- Rash
- Muscle spasms
- Infection of the urinary tract
- Pain in limbs

- Swelling of body, including eyes and other parts of the body
- Shivering
- Redness and pain at injection site
- General ill feeling
- Weight loss
- Weight increase

Uncommon side effects

(may affect up to 1 in 100 people)

- Hepatitis
- Severe allergic reaction (anaphylactic reaction) signs of which may include difficulty breathing, chest pain or chest tightness, and/or feeling dizzy/faint, severe itching of the skin or raised lumps on the skin, swelling of the face, lips, tongue and /or throat, which may cause difficulty in swallowing, collapse
- Movement disorders, paralysis, twitching
- Lump in the eyelid (chalazion), red and swollen eyelids
- Vertigo
- Hearing loss, deafness
- Disorders that affect your lungs, preventing your body from getting enough oxygen. Some of these include: difficulty breathing; shortness of breath; shortness of breath without exercise; breathing that becomes shallow, difficult or stops; wheezing
- Blood clots in your lungs
- Yellow discoloration of the eyes and skin (jaundice)

Rare side effects

(may affect up to 1 in 1,000 people)

- Blood clot in small blood vessels (thrombotic microangiopathy)
- Serious nerve inflammation, which may cause paralysis and difficulty breathing (Guillain-Barré syndrome)

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE BORTEZOMIB

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date stated on the vial and the carton after EXP.

Keep the vial in the outer carton in order to protect from light.

This medicine does not require any special temperature storage conditions.

Reconstituted solution

The reconstituted solution should be used immediately after preparation. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user. However, the chemical and physical stability of the reconstituted solution has been demonstrated for 8 days at 25°C or 15 days at 5 ± 3°C, in the dark, when stored in a vial or in a polypropylene syringe. The total storage time for the reconstituted medicinal product should not exceed 8 or 15 days, depending on storage temperature, prior to administration.

Bortezomib is for single use only. Any unused product or waste material should be disposed of in accordance with local requirements.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What this medicine contains

- The active substance is bortezomib. Each vial contains 1mg of bortezomib (as a mannitol boronic ester). After reconstitution, 1ml of solution for injection contains 1mg bortezomib.
- The other ingredient is mannitol (E421).

What this medicine looks like and contents of the pack

Bortezomib powder for solution for injection is a white to off-white cake or powder.

Bortezomib is packed in a glass vial with rubber stopper and a green flip-off cap.

Each pack contains 1 single-use vial.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Aspire Pharma Limited
Unit 4, Rotherbrook Court
Bedford Road
Petersfield
Hampshire
GU32 3QG
UK

Manufacturer

Synthon Hispania SL
C/Castelló no1
Pol. Las Salinas
Sant Boi de Llobregat
Barcelona
08830
Spain
or
Synthon s.r.o.
Brněnská 32/čp. 597
Blansko
678 01
Czech Republic

This leaflet was last revised 02/2021

1010391-P7.1G

 **ASPIRE**[®]
P H A R M A