

Package leaflet: Information for the user

Isotretinoin 10 mg soft capsules Isotretinoin 20 mg soft capsules

isotretinoin

WARNING

CAN SERIOUSLY HARM AN UNBORN BABY.

Women must use effective contraception.

Do not use if you are pregnant or you think you may be pregnant.

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Isotretinoin is and what it is used for
2. What you need to know before you take Isotretinoin
3. How to take Isotretinoin
4. Possible side effects
5. How to store Isotretinoin
6. Contents of the pack and other information

1. What Isotretinoin is and what it is used for

Isotretinoin contains *isotretinoin* – a substance related to vitamin A, and one of a group of medicines called *retinoids*.

Isotretinoin is used to treat severe types of acne (such as nodular or conglobate acne, or acne that is at risk of causing permanent scarring) in adults and adolescents from 12 years of age only after puberty. You will use Isotretinoin when your acne has not got better with anti-acne treatments, including antibiotics and skin treatments.

Isotretinoin treatment must be supervised by a dermatologist (a doctor specialised in the treatment of skin problems).

2. What you need to know before you take Isotretinoin

Do not take Isotretinoin:

- **if you are pregnant** or breast-feeding
- if there is any chance you could become pregnant, you must follow the precautions under "Pregnancy and prevention programme", see section on "Warnings and precautions".
- **if you are allergic to isotretinoin, peanut or soya or any of the other ingredients** of this medicine (listed in section 6).
- **if you have liver disease**
- **if you have very high levels of cholesterol or triglycerides in your blood**
- **if you have very high levels of vitamin A** in your body (*hypervitaminosis A*)
- **if you are receiving treatment with tetracyclines** (*a type of antibiotic*) at the same time (see "Other medicines and Isotretinoin")

Warnings and precautions

Talk to your doctor or pharmacist before taking Isotretinoin.

- If you have ever had any kind of mental health problems. This includes depression, aggressive tendencies or mood changes. It also includes thoughts about hurting yourself or ending your life. This is because your mood may be affected while taking Isotretinoin.

Mental health problems

You may not notice some changes in your mood and behaviour and so it is very important that you tell your friends and family that you are taking this medicine. They may notice these changes and help you quickly identify any problems that you need to talk to your doctor about.

Below, please find important advice first for women, then for men, and then for all patients.

Important advice for women

Pregnancy prevention programme

Women who are pregnant must not take Isotretinoin

This medicine can seriously harm an unborn baby (the medicine is said to be 'teratogenic') – it can cause serious abnormalities of the unborn baby's brain, face, ear, eye, heart and certain glands (thymus gland and parathyroid gland). It also makes a miscarriage more likely. This may happen even if Isotretinoin is taken only for a short time during pregnancy.

- You must not take Isotretinoin if you are pregnant or if you think you might be pregnant.
- You must not take Isotretinoin if you are breastfeeding. The medicine is likely to pass into your milk and may harm your baby.
- You must not take Isotretinoin if you could get pregnant during treatment.
- You must not get pregnant for one month after stopping this treatment because some medicine may still be left in your body.

Women who could get pregnant are only prescribed Isotretinoin under strict rules. This is because of the risk of serious harm to the unborn baby.

These are the rules:

- Your doctor must explain the risk of harm to the unborn baby - you must understand why you must not get pregnant and what you need to do to prevent getting pregnant.
- You must have talked about contraception (birth control) with your doctor. The doctor will give you information on how not to get pregnant. The doctor may send you to a specialist for contraception advice.
- Before you start treatment, your doctor will ask you to take a pregnancy test. The test must show that you are not pregnant when starting treatment with Isotretinoin.

Women must use effective contraception before, during and after taking Isotretinoin.

- You must agree to use at least one very reliable method of contraception (for example an intra uterine device or contraceptive implant) or, two effective methods that work in different ways (for example a hormonal contraceptive pill and a condom). Discuss with your doctor which methods would be suitable for you.
- You must use contraception for a month before taking Isotretinoin, during treatment and for a month afterwards
- You must use contraception even if you do not have periods or you are not sexually active (unless your doctor decides this is not necessary).

Women must agree to pregnancy testing before, during and after taking Isotretinoin

- You must agree to regular follow-up visits, ideally every month.
- You must agree to have regular pregnancy tests, ideally every month during treatment and, because some medicine may still be left in your body, 1 month after stopping Isotretinoin (unless your doctor decides this is not necessary in your case).
- You must agree to extra pregnancy tests if your doctor asks you.
- You must not get pregnant during treatment or for a month afterwards because some medicine may still be left in your body.
- Your doctor will discuss all these points with you, using a checklist and will ask you (or a parent/guardian) to sign it. This form confirms that you have been told about the risks and that you will follow the rules above.

If you get pregnant while taking Isotretinoin, **stop taking the medicine straight away**, and contact your doctor. Your doctor may send you to a specialist for advice.

Also, if you become pregnant within one month after you stop taking Isotretinoin, you should contact your doctor. Your doctor may send you to a specialist for advice.

Advice for men

The levels of oral retinoid in the semen of men taking Isotretinoin are too low to harm their partners' unborn baby. However, you must never share your medication with anyone.

Advice for all patients

- **Severe Skin reactions** (e.g. erythema multiforme (EM), Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN)) **have been reported with the use of Isotretinoin.** The rash may progress to widespread blistering or peeling of the skin. Also conjunctivitis (red and swollen eyes), ulcers in the mouth, throat, nose and genitals may occur.
- **Rarely, Isotretinoin may cause severe allergic reactions** some of which can affect skin in the form of eczema, hives and bruises or red patches on arms and legs. If you develop an allergic reaction, stop taking Isotretinoin, seek urgent advice from a doctor and tell him that you are taking this medicine.
- **Cut down on intensive exercise and physical activity.** Isotretinoin can cause muscle and joint pain particularly in children and teenagers undertaking vigorous physical activity.
- **Isotretinoin has been associated with inflammatory bowel disease.** Your doctor will take you off Isotretinoin if you have severe bloody diarrhoea without any history of gastrointestinal disorders.
- **Isotretinoin may cause dry eyes, problems using contact lenses and visual difficulties including decreased night vision.** Tell your doctor if you have any of these symptoms. Your doctor may ask you to use lubricating eye ointment or tear replacement therapy. If you use contact lenses and you have developed an intolerance to contact lenses, you may be advised to

- wear glasses during the treatment. Your doctor may refer you to a specialist for advice if you develop visual difficulties and you may be asked to stop taking Isotretinoin.
- **Benign intracranial hypertension has been reported with Isotretinoin use** and in some cases where Isotretinoin was used together with tetracyclines (a type of antibiotic). Stop taking Isotretinoin and seek urgent advice from your doctor if you develop symptoms like headache, nausea, vomiting and visual disturbances. Your doctor may refer you to a specialist to check for swelling of optic disk in the eye (papilloedema).
 - **Isotretinoin may increase liver enzyme levels.** Your doctor will do blood tests before, during and after Isotretinoin treatment to check these levels. If they stay high, your doctor may lower your dose or take you off Isotretinoin .
 - **Isotretinoin commonly increases blood fats**, such as cholesterol or triglycerides. Your doctor will test these levels before, during and after Isotretinoin treatment. It is best that you do not drink alcoholic drinks while on treatment. Tell your doctor if you know that you already have high blood fats, diabetes (high blood sugars), are overweight, or an alcoholic. You may need blood tests more often. If your blood fats stay high, your doctor may lower your dose, or take you off Isotretinoin.
 - **Tell your doctor if you have any kidney problems.** Your doctor may start you on a lower dose of Isotretinoin and then increase it to the maximum tolerated dose.
 - **Isotretinoin may increase blood sugar levels.** In rare cases, people become diabetic. Your doctor may monitor blood sugar levels during treatment, particularly if you already have diabetes, are overweight, or are an alcoholic.
 - **Your skin is likely to get dry.** Use a skin moisturising ointment or cream and a lip balm during treatment. To prevent skin irritation you should avoid using exfoliating or anti-acne products.
 - **Avoid too much sun and do not use a sun-lamp or sun-bed.** Your skin may become more sensitive to sunlight. Before you go out in the sun, use a sun-protection product with a high protection factor (SPF 15 or higher).
 - **Don't have any cosmetic skin treatments.** Isotretinoin may make your skin more fragile. Don't have any waxing (hair removal), dermabrasion or laser treatments (removing horny skin or scars) during treatment, or for at least 6 months after treatment. They could cause scarring, skin irritation, or rarely, changes in the colour of your skin.

Additional precautions

You should never give this medicinal product to another person. Please take any unused capsules to your pharmacist at the end of treatment.

You should not donate blood during treatment with this medicine and for 1 month after stopping Isotretinoin because an unborn baby could be harmed if a pregnant patient receives your blood.

Children and adolescents

The use of Isotretinoin in children under the age of 12 is not recommended. This is because it is not known if it is safe or effective in this age group.

Use in adolescents over 12 years of age only after puberty.

Other medicines and Isotretinoin

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines including herbal and non-prescription products.

- **Do not take vitamin A supplements or tetracyclines** (a type of antibiotic), or use any skin treatments for acne while you are on Isotretinoin. It is fine to use moisturisers and emollients (skin creams or preparations that prevent water loss and have a softening effect on the skin).
- **Avoid the use of topical keratolytic or exfoliative anti-acne agents** while you are on Isotretinoin.

Pregnancy, breast-feeding and fertility

Pregnancy

Isotretinoin must not be used during pregnancy. If you are able to get pregnant, you should use effective contraception during and up to one month after Isotretinoin treatment.

If you do get pregnant while taking Isotretinoin, or in the month after treatment has stopped, **stop taking the medicine straight away**, and contact your doctor. He or she may refer you to a specialist for advice.

If used during pregnancy, Isotretinoin is likely to damage an unborn baby. It also increases the risk of miscarriage.

Isotretinoin can cause serious abnormalities of the brain, face, ear, eye, heart and some glands (called the thymus gland and parathyroid gland) of the unborn baby.

Breast-feeding

You must not take Isotretinoin if you are breastfeeding. The medicine is likely to pass into your breast milk and may harm your baby.

For more information on pregnancy and contraception, see section 2 "Pregnancy and prevention programme".

Driving and using machines

You may not see as well at night during your treatment. This can happen suddenly. In rare cases this has continued after the treatment has stopped. Drowsiness and dizziness have been reported very rarely. If this happens to you, you should not drive or operate machinery.

Isotretinoin contains soya bean oil

- If you are allergic to peanut or soya, do not take this medicinal product.

3. How to take Isotretinoin

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The usual starting dose is 0.5 mg per kilogram body weight per day (0.5 mg/kg/day). So if you weigh 60 kg, your dose will usually start at 30 mg a day.

Take the capsules once or twice daily.

Take on a full stomach. Swallow them whole, with some water. Do not chew the capsule.

After a few weeks your doctor may adjust your dose. This depends on how you are getting on with your medicine. For most patients the dose will be between 0.5 and 1.0 mg/kg/day. If you think that Isotretinoin is too strong or too weak, talk to your doctor or pharmacist.

If you have severe kidney problems, you will usually start on a lower dose (such as 10 mg/day) which will be increased up to the highest dose your body can tolerate. If your body can't tolerate the recommended dose, you may be prescribed a lower dose: that can mean you are treated for longer and your acne might be more likely to come back.

A course of treatment usually lasts for 16 to 24 weeks. Most patients only need one course. Your acne may continue to improve for up to 8 weeks after the end of the treatment. You won't usually start another course until then.

Some people find their acne gets worse during the first weeks of treatment. It usually improves as treatment goes on.

If you take more Isotretinoin than you should

If you take too many capsules or someone else accidentally takes your medicine, contact your doctor, pharmacist or nearest hospital immediately.

If you forget to take Isotretinoin

If you miss a dose take it as soon as you can. However, if it is less than 4 hours until your next dose, skip the missed dose and carry on as before. Do not take a double dose (two doses close together). If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Below, you will first find a list of serious side effects that require immediate medical attention. After that you will find an overview of other side effects. Over time these side effects usually get better or disappear after the end of the treatment.

SIDE EFFECTS REQUIRING IMMEDIATE MEDICAL ATTENTION**Skin problems**

Frequency not known (*frequency cannot be estimated from available data*)

- Serious skin rashes (erythema multiforme, Stevens-Johnson syndrome, and toxic epidermal necrolysis), which are potentially life-threatening and require immediate medical attention. These appear initially as circular patches often with central blisters usually on arms and hands or legs and feet, more severe rashes may include blistering of the chest and back. Additional symptoms such as inflammation of the eye (conjunctivitis) or ulcers of the mouth, throat or nose may occur. Severe forms of rash may progress to widespread peeling of the skin which can be life threatening. These serious skin rashes are often preceded by headache, fever, body aches (flu-like symptoms).

If you develop a serious rash or these skin symptoms, stop taking Isotretinoin and contact your doctor immediately.

Mental problems

Rare effects (*may affect up to 1 in every 1000 people*)

- Depression or related disorders. Signs of this include sad or altered mood, anxiety, feelings of emotional discomfort.
- Existing depression getting worse.
- Becoming violent or aggressive.

Very rare effects (*may affect up to 1 in every 10,000 people*)

- Some people have had thoughts or feelings about hurting themselves or ending their own lives (suicidal thoughts), have tried to end their own lives (attempted suicide), or have ended their lives (suicide). These people may not appear to be depressed.
- Unusual behaviour.
- Signs of psychosis: a loss of contact with reality, such as hearing voices or seeing things that are not there.

Contact your doctor straight away if you get signs of any of these mental problems. Your doctor may tell you to stop taking Isotretinoin. That may not be enough to stop the effects: you may need more help, and your doctor can arrange this.

Allergic reactions

Rare effects (*may affect up to 1 in every 1000 people*)

- Serious (*anaphylactic*) reactions: difficulty breathing or swallowing caused by sudden swelling of the throat, face, lips and mouth. Also sudden swelling of the hands, feet and ankles.

Very rare effects (*may affect up to 1 in every 10,000 people*)

- Sudden tight chest, shortness of breath and wheezing, particularly if you have asthma.

If you have a serious reaction, get emergency medical help immediately.

If you have any allergic reaction, stop taking Isotretinoin and contact your doctor.

Bones and muscles

Frequency not known (*frequency cannot be estimated from available data*)

- Muscle weakness which can be potentially life-threatening, may be associated with trouble moving arms or legs, painful, swollen, bruised areas of the body, dark- coloured urine, reduced or no urine output, confusion or dehydration. These are sign of a breakdown of muscle tissue which can lead to kidney failure. This may occur if you are doing intensive physical activity while you're on Isotretinoin.

Liver and kidney problems

Very rare effects (*may affect up to 1 in every 10,000 people*)

- Yellow skin or eyes, and feeling tired. These can be signs of hepatitis. **Stop taking Isotretinoin straight away and contact your doctor.**
- Difficulty urinating (passing water), swollen and puffy eyelids, feeling excessively tired. These may be signs of kidney inflammation. **Stop taking Isotretinoin straight away and contact your doctor.**

Nervous system problems

Very rare effects (*may affect up to 1 in every 10,000 people*)

- Lasting headache, along with feeling sick (*nausea*), being sick (*vomiting*) and change in your eyesight including blurred vision. These may be signs of *benign intracranial hypertension*, especially if Isotretinoin is taken with antibiotics called tetracycline. **Stop taking Isotretinoin straight away and contact your doctor.**

Gut and stomach problems

Very rare effects (*may affect up to 1 in every 10,000 people*)

- Severe abdominal (tummy) pain, with or without severe bloody diarrhoea, feeling sick (*nausea*) and being sick (*vomiting*). These can be signs of serious gut conditions. **Stop taking Isotretinoin straight away and contact your doctor.**

Eye disorders

Very rare effects (*may affect up to 1 in every 10,000 people*)

- Blurred vision.

If you get blurred vision, stop taking Isotretinoin straight away and contact your doctor. If your sight is affected in any other way tell a doctor as soon as you can.

OTHER SIDE EFFECTS

Very common side effects with Isotretinoin: (*may affect more than 1 in 10 people*)

- Dryness of the skin, especially of the lips and face; inflamed skin, chapped and inflamed lips, rash, mild itching and slight peeling. Use a moisturising cream from the start of treatment.
- Skin becomes more fragile and redder than usual, especially the face.
- Back pain; muscle pain; joint pain particularly in children and teenagers. **To avoid making any bone or muscle problems worse**, cut down on intensive physical activity while you're on Isotretinoin.

- Inflammation of the eye (conjunctivitis) and eyelid area; eyes feel dry and irritated. Ask a pharmacist for suitable eye drops. If you get dry eyes and wear contact lenses, you may need to wear glasses instead.
- Raised liver enzymes seen in blood tests.
- Changed levels of fats in the blood (including HDL or triglycerides).
- Bruising, bleeding or clotting more easily - if clotting cells are affected.
- Anaemia – weakness, dizziness, pale skin – if red blood cells are affected.

Common side effects with Isotretinoin: *(may affect up to 1 in 10 people)*

- Headache.
- Higher levels of cholesterol in the blood.
- Protein or blood in the urine.
- More liable to get infections if the white blood cells are affected.
- Inside of the nose becomes dry and crusted, causing mild nosebleeds.
- Sore or inflamed throat and nose.
- Allergic reactions such as rash, itchiness. If you have any allergic reaction, stop taking Isotretinoin and contact your doctor.

Rare side effects with Isotretinoin: *(may affect up to 1 in 1,000 people)*

- Hair loss (*alopecia*). This is usually only temporary. Your hair should return to normal after the treatment ends.

Very rare side effects with Isotretinoin: *(may affect up to 1 in 10,000 people)*

- You may see less well at night; colour blindness and colour vision gets worse.
- Sensitivity to light may increase; you may find that you need to wear sunglasses to protect your eyes from too bright sunlight.
- Other sight problems including blurred vision, distorted vision, cloudy surface on the eye (corneal opacity, cataracts).
- Excessive thirst; frequent need to urinate; blood tests show an increase in your blood sugar. These can all be signs of diabetes.
- Acne can get worse in the first few weeks, but symptoms should improve with time.
- Skin inflamed, swollen, and darker than usual, especially on the face.
- Excess sweating or itching.
- Arthritis; bone disorders (delayed growth, extra growth and changes to bone density); growing bones may stop growing.
- Calcium deposits in soft tissue, sore tendons, high levels of muscle breakdown products in your blood if you exercise vigorously.
- Increased sensitivity to light.
- Inflammation at the base of the nail, changes to nails.
- Swellings, discharging, pus.
- Thickened scarring after surgery.
- Increased body hair.
- Convulsions, drowsiness, dizziness.
- Lymph glands may become swollen.
- Dry throat, hoarseness.
- Hearing difficulties.
- Generally feeling unwell.
- High levels of uric acid in the blood.
- Bacterial infections.
- Inflammation of the blood vessels (sometimes with bruising, red patches)
- Rhabdomyolysis.

Unknown frequency: *(frequency cannot be estimated from the available data)*

- Dark or cola-coloured urine
- Problems getting or maintaining an erection
- Lower libido
- Breast swelling with or without tenderness in males

- Vaginal dryness

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Isotretinoin

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.
Store in the original package. Keep the blister in the outer carton in order to protect from light.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Isotretinoin contains

- **The active substance** is isotretinoin.
- **The other ingredients** are soya-bean oil hydrogenated, hydrogenated vegetable oil, beeswax white, disodium edetate, butylhydroxyanisole (E320), soyabean oil refined, polysorbate 80, gelatin, glycerol (E422), iron oxide red (E172), titanium dioxide (E171), paraffin, light liquid printing ink containing shellac, iron oxide black (E172), propylene glycol (E1520).

What Isotretinoin looks like and contents of the pack

Isotretinoin comes in soft capsules containing either 10 mg or 20 mg isotretinoin.

The 10 mg capsules comprise of oval shaped, light pink colored, opaque soft gelatin capsules imprinted with `RR' in black edible ink, of approximately 9.30 ± 0.5 mm length and 6.60 ± 0.5 mm width, containing orange-yellow colored oily suspension.

The 20 mg capsules comprise of oval shaped, orange to reddish orange colored, opaque soft gelatin capsules imprinted with `RR' in black edible ink, of approximately 13.20 ± 0.5 mm length and 7.80 ± 0.5 mm width, containing orange-yellow colored oily suspension.

The capsules are packed in blister packs of 20, 30, 50, 60, 90 or 100 capsules.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Sun Pharmaceutical Industries Europe B.V.
Polarisavenue 87
2132 JH Hoofddorp
The Netherlands

Manufacturers

S.C. Terapia S.A.
124 Fabricii Street
400 632 Cluj-Napoca,
Romania

Alkaloida Chemical Company Zrt
Kabay János u.29,
Tiszavasvári, H-4440,
Hungary

Sun Pharmaceutical Industries Europe B.V.
Polarisavenue 87,
2132 JH Hoofddorp,
The Netherlands

This medicinal product is authorised in the Member States of the EEA under the following names:

Germany	Isotretinoin Basics 10 mg Weichkapseln Isotretinoin Basics 20 mg Weichkapseln
Spain:	Isotretinoína SUN 10 mg cápsulas blandas EFG Isotretinoína SUN 20 mg cápsulas blandas EFG
Hungary	Sotret Neo 10 mg lágy kapszula Sotret Neo 20 mg lágy kapszula
Italy	Isotretinoina SUN 10 mg capsule molli Isotretinoina SUN 20 mg capsule molli
The Netherlands	Isotretinoïne SUN 10 mg, zachte capsules Isotretinoïne SUN 20 mg, zachte capsules
Poland	Sotret 10 mg Sotret 20 mg
Romania	Sotret 10 mg capsule moi Sotret 20 mg capsule moi
United Kingdom	Isotretinoin 10 mg soft capsules Isotretinoin 20 mg soft capsules

This leaflet was last revised in February 2020.