

Package leaflet: Information for the patient

Otezla 10 mg film-coated tablets
Otezla 20 mg film-coated tablets
Otezla 30 mg film-coated tablets
apremilast

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Otezla is and what it is used for
2. What you need to know before you take Otezla
3. How to take Otezla
4. Possible side effects
5. How to store Otezla
6. Contents of the pack and other information

1. What Otezla is and what it is used for

What Otezla is

Otezla contains the active substance 'apremilast'. This belongs to a group of medicines called phosphodiesterase 4 inhibitors, which help to reduce inflammation.

What Otezla is used for

Otezla is used to treat adults with the following conditions:

- **Active psoriatic arthritis** - if you cannot use another type of medicine called 'Disease-Modifying Antirheumatic Drugs' (DMARDs) or when you have tried one of these medicines and it did not work.
- **Moderate to severe chronic plaque psoriasis** - if you cannot use one of the following treatments or when you have tried one of these treatments and it did not work:
 - phototherapy - a treatment where certain areas of skin are exposed to ultraviolet light
 - systemic therapy - a treatment that affects the entire body rather than just one local area, such as 'ciclosporin', 'methotrexate' or 'psoralen'.

What psoriatic arthritis is

Psoriatic arthritis is an inflammatory disease of the joints, usually accompanied by psoriasis, an inflammatory disease of the skin.

What plaque psoriasis is

Psoriasis is an inflammatory disease of the skin, which can cause red, scaly, thick, itchy, painful patches on your skin and can also affect your scalp and nails.

How Otezla works

Psoriatic arthritis and psoriasis are usually lifelong conditions and there is currently no cure. Otezla works by reducing the activity of an enzyme in the body called 'phosphodiesterase 4', which is involved in the process of inflammation. By reducing the activity of this enzyme, Otezla can help to control the inflammation

associated with psoriatic arthritis and psoriasis, and thereby reduce the signs and symptoms of these conditions.

In psoriatic arthritis, treatment with Otezla results in an improvement in swollen and painful joints, and can improve your general physical function.

In psoriasis, treatment with Otezla results in a reduction in psoriatic skin plaques and other signs and symptoms of the disease.

Otezla has also been shown to improve the quality of life in patients with psoriasis or psoriatic arthritis. This means that the impact of your condition on daily activities, relationships and other factors should be less than it was before.

2. What you need to know before you take Otezla

Do not take Otezla:

- if you are allergic to apremilast or any of the other ingredients of this medicine (listed in section 6).
- if you are pregnant or think you may be pregnant.

Warnings and precautions

Talk to your doctor or pharmacist before taking Otezla.

Symptoms you should be aware of

If your doctor considers you to be underweight, and you observe an unintentional loss of body weight while being treated with Otezla, you should talk to your doctor.

If you have severe kidney problems then the recommended dose of Otezla is 30 mg **once a day (morning dose)**. Your doctor will talk to you about how to increase your dose when you first start taking Otezla.

Before starting treatment with Otezla, inform your doctor if you are suffering from symptoms of worsening depression with suicidal thinking or behaviour, especially if you take any additional medicines since some of those could increase the probability of these side effects. You or your caregiver should also immediately inform your doctor of any changes in behaviour or mood and of any suicidal thoughts you may have. You may also experience sleeplessness, or depressive mood.

If you experience severe diarrhoea, nausea, or vomiting, you should talk to your doctor.

Children and adolescents

Otezla has not been studied in children and adolescents, therefore it is not recommended for use in children and adolescents aged 17 years and under.

Other medicines and Otezla

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes medicines obtained without a prescription and herbal medicines. This is because Otezla can affect the way some other medicines work. Also some other medicines can affect the way Otezla works.

In particular, tell your doctor or pharmacist before taking Otezla if you are taking any of the following medicines:

- rifampicin – an antibiotic used for tuberculosis
- phenytoin, phenobarbital and carbamazepine - medicines used in the treatment of seizures or epilepsy
- St John's Wort – a herbal medicine for mild anxiety and depression.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

There is little information about the effects of Otezla in pregnancy. You should not become pregnant while taking this medicine and should use an effective method of contraception during treatment with Otezla. It is not known if this medicine passes into human milk. You should not use Otezla while breast-feeding.

Driving and using machines

Otezla has no effect on the ability to drive and use machines.

Otezla contains lactose

Otezla contains lactose (a type of sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Otezla

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

How much to take

- When you first start taking Otezla, you will receive a ‘treatment initiation pack’ which contains all the doses as listed in the table below.
- The ‘treatment initiation pack’ is clearly labelled to make sure you take the correct tablet at the correct time.
- Your treatment will start at a lower dose and will gradually be increased over the first 6 days of treatment.
- The ‘treatment initiation pack’ will also contain enough tablets for another 8 days at the recommended dose (days 7 to 14).
- The recommended dose of Otezla is 30 mg twice a day after the titration phase is complete - one 30 mg dose in the morning and one 30 mg dose in the evening, approximately 12 hours apart, with or without food.
- This is a total daily dose of 60 mg. By the end of day 6 you will have reached this recommended dose.
- Once the recommended dose has been reached, you will only get the 30 mg tablet strength in your prescribed packs. You will only ever need to go through this stage of gradually increasing your dose once even if you re-start treatment.

Day	Morning Dose	Evening Dose	Total Daily Dose
Day 1	10 mg (pink)	Do not take a dose	10 mg
Day 2	10 mg (pink)	10 mg (pink)	20 mg
Day 3	10 mg (pink)	20 mg (brown)	30 mg
Day 4	20 mg (brown)	20 mg (brown)	40 mg
Day 5	20 mg (brown)	30 mg (beige)	50 mg
Day 6 onwards	30 mg (beige)	30 mg (beige)	60 mg

People with severe kidney problems

If you have severe kidney problems then the recommended dose of Otezla is 30 mg **once a day (morning dose)**. Your doctor will talk to you about how to increase your dose when you first start taking Otezla.

How and when to take Otezla

- Otezla is for oral use.
- Swallow the tablets whole, preferably with water.
- You can take the tablets either with or without food.
- Take Otezla at about the same time each day, one tablet in the morning and one tablet in the evening.

If your condition has not improved after six months of treatment, you should talk to your doctor.

If you take more Otezla than you should

If you take more Otezla than you should, talk to a doctor or go to a hospital straight away. Take the medicine pack and this leaflet with you.

If you forget to take Otezla

- If you miss a dose of Otezla, take it as soon as you remember. If it is close to the time for your next dose, just skip the missed dose. Take the next dose at your regular time.
- Do not take a double dose to make up for a forgotten dose.

If you stop taking Otezla

- You should continue taking Otezla until your doctor tells you to stop.
- Do not stop taking Otezla without talking to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Some side effects could be serious. Uncommon instances of suicidal thinking and behaviour (including suicide) were reported. Please notify your doctor immediately of any feelings of depression, suicidal thoughts or suicidal behaviour you may have. You may also experience sleeplessness (common), or depressive mood (common).

Very common side effects (may affect more than 1 in 10 people)

- diarrhoea
- nausea

Common side effects (may affect up to 1 in 10 people)

- cough
- back pain
- vomiting
- feeling tired
- stomach pain
- loss of appetite
- frequent bowel movements
- difficulty sleeping (insomnia)
- indigestion or heartburn
- headaches, migraines or tension headaches
- upper respiratory tract infections such as cold, runny nose, sinus infection
- inflammation and swelling of the tubes in your lungs (bronchitis)
- common cold (nasopharyngitis)
- depression

Uncommon side effects (may affect up to 1 in 100 people)

- rash
- hives (urticaria)
- weight loss
- allergic reaction
- bleeding in the bowel or in the stomach
- suicidal ideation or behaviour

Not known side effects (frequency cannot be estimated from the available data):

- severe allergic reaction (may include swelling of the face, lips, mouth, tongue, or throat that may lead to difficulty breathing or swallowing)

If you are 65 years of age or older, you might have a higher risk of complications of severe diarrhoea, nausea and vomiting.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme

Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Otezla

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the blister or on the wallet or on the carton after EXP. The expiry date refers to the last day of that month.
- Do not store above 30°C.
- Do not use this medicine if you notice any damage or signs of tampering to the medicine packaging.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Otezla contains

- The active substance is apremilast.
- Otezla 10 mg film-coated tablets: each film-coated tablet contains 10 mg of apremilast.
- Otezla 20 mg film-coated tablets: each film-coated tablet contains 20 mg of apremilast.
- Otezla 30 mg film-coated tablets: each film-coated tablet contains 30 mg of apremilast.
- The other ingredients in the tablet core are cellulose microcrystalline, lactose monohydrate, croscarmellose sodium and magnesium stearate.
- The film-coating contains poly(vinyl alcohol), titanium dioxide (E171), macrogol (3350), talc, iron oxide red (E172).
- The 20 mg film-coated tablet also contains iron oxide yellow (E172).
- The 30 mg film-coated tablet also contains iron oxide yellow (E172) and iron oxide black (E172).

What Otezla looks like and contents of the pack

The Otezla 10 mg film-coated tablet is a pink, diamond shaped film-coated tablet with “APR” engraved on one side and “10” on the opposite side.

The Otezla 20 mg film-coated tablet is a brown, diamond shaped film-coated tablet with “APR” engraved on one side and “20” on the opposite side.

The Otezla 30 mg film-coated tablet is a beige, diamond shaped film-coated tablet with “APR” engraved on one side and “30” on the opposite side.

Pack sizes

- The treatment initiation pack is a folding wallet containing 27 film-coated tablets: 4 x 10 mg tablets, 4 x 20 mg tablets and 19 x 30 mg tablets.
- The one-month standard pack contains 56 x 30 mg film-coated tablets.
- The three-month standard pack contains 168 x 30 mg film-coated tablets.

Marketing Authorisation Holder

Celgene Europe B.V.
Winthontlaan 6 N
3526 KV Utrecht
Netherlands

Manufacturer

Celgene Distribution B.V.
Winthontlaan 6 N
3526 KV Utrecht
Netherlands

This leaflet was last revised in

August 2019

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site:
<http://www.ema.europa.eu>.