

# MEGACE 160 mg Tablets

## Megestrol acetate

### Package leaflet: Information for the user

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What Megace tablets are and what they are used for
2. What you need to know before you take Megace tablets
3. How to take Megace tablets
4. Possible side effects
5. How to store Megace tablets
6. Contents of the pack and other information

#### 1. What Megace tablets are and what they are used for

The name of your medicine is Megace tablets. Megace tablets contain the active ingredient megestrol acetate. Megestrol acetate belongs to a group of medicines called progestogens and is used to treat certain types of cancer known as hormone-dependent cancer, which include cancer of the breast.

#### 2. What you need to know before you take Megace tablets

##### Do not take Megace tablets if you:

- are allergic to megestrol acetate or any of the other ingredients of this medicine (listed in section 6).

##### Warnings and precautions

Tell your doctor if you:

- have severe liver problems
- currently suffer from, or have a history of suffering from blood clots (thrombophlebitis)
- have diabetes, as Megace tablets can increase your blood sugar levels

If you are over 65 years of age your doctor may want to monitor your kidney function from time to time, as this can sometimes decrease with age.

##### Other medicines and Megace tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

##### Pregnancy and breast-feeding

Do not take Megace tablets if you are pregnant, think you may be pregnant, or if you are breast-feeding.

##### Driving and using machines

Megace tablets do not usually affect your ability to drive.

##### Megace tablets contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

#### 3. How to take Megace tablets

Always take Megace tablets exactly as your doctor has told you. You should check with your doctor if you are not sure. The usual dosage is 160 mg megestrol acetate per day.

Tablets should be swallowed with a glass of water.

This medicine is not recommended for children.

Megace tablets should be taken for at least two months to see if they are effective. Some patients take them for many years.

**Do not stop** taking them, unless advised to do so by your doctor.

#### If you take more Megace tablets than you should

If you take more Megace than you should, go to your nearest hospital emergency department or contact your doctor immediately.

Take the empty container and any remaining medicine with you.

#### If you forget to take Megace tablets

If you miss a dose, do not worry. If you remember within a few hours of the missed dose take it as soon as possible, but if it is almost time for your next dose then skip the forgotten one and continue as before. Do not take a double dose to make up for the one you missed.

#### 4. Possible side effects

As with all medicines, Megace can cause side effects although not everybody gets them.

If you notice any of the following, stop taking Megace and contact your doctor **immediately**:

- swelling of the face, lips, or tongue
- skin rash
- difficulty breathing

As these may be signs of an allergic reaction.

P1UK01

Other side effects which may occur with Megace tablets are:

Very Common (affects more than 1 in 10 people)

- increase in appetite and weight gain (not associated with water retention)
- shortness of breath
- hot flushes, reddening of the face and neck
- high blood pressure
- increased blood sugar levels
- rounded face, sometimes called 'moon' face
- constipation
- swelling of a vein due to a blood clot
- blood clot in the lungs

Signs of a blood clot can include loss of co-ordination, slurred speech, pains in your groin, pains or numbness in your arm or leg (especially in calf of leg), shortness of breath, weakness, headache, and fainting. If you experience any of these signs tell your doctor **immediately**.

Common (affects between 1 in 10 to 1 in 100 people)

- temporary increase in the signs and symptoms of your disease.
- changes in mood
- numbness, pain or tingling in fingers or thumb
- heart attack
- hair loss
- diarrhoea
- tiredness
- feeling weak
- impotence
- passing wind
- needing to urinate more often
- feeling sick and being sick
- changes in vaginal bleeding (breakthrough bleeding, spotting)
- build up of fluid which causes swelling (mainly in the ankles)
- pain
- rash, including hives or nettle rash (urticaria)

#### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see contact details below). By reporting side effects you can help provide more information on the safety of this medicine.

##### Ireland:

HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517.

Website: [www.hpra.ie](http://www.hpra.ie); E-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie)

##### United Kingdom:

Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

#### 5. How to store Megace tablets

Do not store above 25°C. Store in the original package in order to protect from moisture. They should not get too hot or damp; so do not leave your tablets near a radiator, on a window sill or in the bathroom.

Please check the expiry date of the product, shown on the carton and blister of Megace tablets. Do not use after the month stated.

Keep this medicine out of the sight and reach of children.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

#### 6. Contents of the pack and other information

##### What Megace tablets contain

Each Megace tablet contains 160 mg of the active ingredient megestrol acetate. The other ingredients are: microcrystalline cellulose, lactose monohydrate 224.5 mg, magnesium stearate, povidone, colloidal anhydrous silica, and sodium starch glycolate.

##### What Megace tablets look like and contents of the pack

Megestrol acetate tablets (160 mg) are oval shaped biconvex tablets scored on one side and engraved with "160" on the other side and are available in blister packs of 30 tablets.

##### Marketing Authorisation Holder and Manufacturer

###### Marketing Authorisation Holder

IE:  
PharmaSwiss Česká republika s.r.o.,  
Jankovcova 1569/2c  
170 00, Prague 7  
Czech Republic

UK:  
Bausch and Lomb UK Ltd.  
Bausch & Lomb House  
106 London Road  
Kingston-Upon-Thames, Surrey, KT2 6TN, UK


###### Manufacturer

Haupt Pharma Amareg GmbH  
Donaustauffer Strasse 378  
93055 Regensburg, Germany

This leaflet was last revised in September 2020

P1UK01

Artwork prepared by  
 PharmaSwiss

Product Title	PIL - Megace 160 mg, tablets			Draft date	Reason
Market	UK/IE	Printing paper	- gr/m <sup>2</sup>	21.03.2019.	Version 02 - text correction
Item Code	P1UK01	Dimension (width)	148 mm	04.09.2020.	Version 02 - new item code, MAH change
EAN Code	-	Dimension (height)	297 mm		
Pharma Code	-	Final dimension	-		
Material Code	-	Scale	1:1		
Printing colors:					
					
Typefaces: Helvetica Nue LT Std Condensed (8pt)					

