

## PACKAGE LEAFLET: INFORMATION FOR THE USER

Dutasteride/Tamsulosin hydrochloride 0.5 mg/0.4 mg Capsules, Hard  
Dutasteride/tamsulosin hydrochloride

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

## What is in this leaflet:

1. What Dutasteride/Tamsulosin is and what it is used for
2. What you need to know before you take Dutasteride/Tamsulosin
3. How to take Dutasteride/Tamsulosin
4. Possible side effects
5. How to store Dutasteride/Tamsulosin
6. Contents of the pack and other information

## 1. What Dutasteride/Tamsulosin is and what it is used for

Dutasteride/Tamsulosin is used to treat men with an enlarged prostate (*benign prostatic hyperplasia*) - a non-cancerous growth of the prostate gland, caused by producing too much of a hormone called dihydrotestosterone.

Dutasteride/Tamsulosin is a combination of two different medicines called dutasteride and tamsulosin. Dutasteride belongs to a group of medicines called *5-alpha reductase inhibitors* and tamsulosin belongs to a group of medicines called *alpha-blockers*.

As the prostate grows, it can lead to urinary problems, such as difficulty in passing urine and a need to go to the toilet frequently. It can also cause the flow of the urine to be slower and less forceful. If left untreated, there is a risk that your urine flow will be completely blocked (*acute urinary retention*). This requires immediate medical treatment. Sometimes surgery is necessary to remove or reduce the size of the prostate gland.

Dutasteride lowers the production of a hormone called dihydrotestosterone, which helps to shrink the prostate and relieve the symptoms. This will reduce the risk of acute urinary retention and the need for surgery. Tamsulosin acts by relaxing the muscles in your prostate gland, making it easier to pass urine and rapidly improving your symptoms.

## 2. What you need to know before you take Dutasteride/Tamsulosin

## Do not take Dutasteride/Tamsulosin

- if you're a woman (because this medicine is for men only).
  - if you're a child or adolescent less than 18 years old.
  - if you're allergic to dutasteride, other 5-alpha reductase inhibitors, tamsulosin, soya, peanut or to any of the other ingredients of this medicine (listed in section 6).
  - if you have low blood pressure which makes you feel dizzy, lightheaded or faint (*orthostatic hypotension*).
  - if you have a severe liver disease.
- If you think any of these apply to you, don't take this medicine until you have checked with your doctor.

## Warnings and precautions

Talk to your doctor before taking Dutasteride/Tamsulosin

In some clinical studies, more patients taking dutasteride and another medicine called an alpha-blocker, like tamsulosin, experienced heart failure than patients taking only dutasteride or only an alpha blocker. Heart failure means your heart does not pump blood as well as it should.

**Make sure your doctor knows about liver problems.** If you have had any illness affecting your liver, you may need some additional check-ups while you are taking Dutasteride/Tamsulosin.

**Make sure your doctor knows if you have severe problems with your kidney.**

**Cataract (cloudy lens) surgery.** If you are going to have surgery to remove a cataract, your doctor may ask you to stop taking Dutasteride/Tamsulosin for a while before your operation. Tell your eye specialist before your operation that you are taking Dutasteride/Tamsulosin or tamsulosin (or have previously taken it). Your specialist will need to take appropriate precautions to help prevent complications during your operation.

**Women, children and adolescents** must not handle leaking Dutasteride/Tamsulosin capsules, because the active ingredient can be absorbed through the skin.

**Wash the affected area immediately** with soap and water if there is any contact with the skin.

**Use a condom during sexual intercourse.** Dutasteride has been found in the semen of men taking Dutasteride/Tamsulosin. If your partner is or may be pregnant, you must avoid exposing her to your semen as dutasteride may affect the normal development of a male baby. Dutasteride has been shown to decrease sperm count, semen volume and sperm motility. This could reduce your fertility.

Dutasteride/Tamsulosin affects a blood test for PSA (*prostate-specific antigen*), which is sometimes used to detect prostate cancer. Your doctor should be aware of this effect and can still use the test to detect prostate cancer. If you are having a blood test for PSA, tell your doctor that you are taking Dutasteride/Tamsulosin. **Men taking Dutasteride/Tamsulosin should have their PSA tested regularly.**

In a clinical study of men at increased risk of prostate cancer, men taking dutasteride had a **serious form of prostate cancer more often** than men who did not take dutasteride. The effect of dutasteride on this serious form of prostate cancer is not clear.

Dutasteride/Tamsulosin may cause **breast enlargement and tenderness**. If this becomes troublesome, or if you notice **breast lumps or nipple discharge** you should talk to your doctor about these changes as these may be signs of a serious condition, such as breast cancer.

→ **Contact your doctor or pharmacist** if you have any questions about taking Dutasteride/Tamsulosin.

## Other medicines and Dutasteride/Tamsulosin

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Don't take Dutasteride/Tamsulosin with these medicines:

- other alpha blockers (for enlarged prostate or high blood pressure)

Dutasteride/Tamsulosin is not recommended with these medicines:

- ketoconazole (used to treat fungal infections)

Some medicines can react with Dutasteride/Tamsulosin and may make it more likely that you'll have side-effects.

These medicines include:

- PDE5 inhibitors (used to help achieve or maintain an erection) such as vardenafil, sildenafil citrate and tadalafil
- verapamil or diltiazem (for high blood pressure)
- ritonavir or indinavir (for HIV)
- itraconazole or ketoconazole (for fungal infections)
- nefazodone (an antidepressant)
- cimetidine (for stomach ulcers)
- warfarin (for blood clotting)
- erythromycin (an antibiotic used to treat infections)
- paroxetine (an antidepressant)
- terbinafine (used to treat fungal infections)
- diclofenac (used to treat pain and inflammation)

→ **Tell your doctor** if you are taking any of these medicines.

**Dutasteride/Tamsulosin with food** Dutasteride/Tamsulosin should be taken 30 minutes after the same meal each day.

**Pregnancy, breastfeeding and fertility** Dutasteride/Tamsulosin **must not be taken by women.**

**Women who are pregnant (or may be) must not handle leaking capsules.** Dutasteride is absorbed through the skin and can affect the normal development of a male baby. This is a particular risk in the first 16 weeks of pregnancy.

**Use a condom during sexual intercourse.** Dutasteride has been found in the semen of men taking Dutasteride/Tamsulosin. If your partner is or may be pregnant, you must avoid exposing her to your semen.

Dutasteride/Tamsulosin has been shown to reduce sperm count, semen volume and sperm movement. Therefore male fertility may be reduced.

→ **Contact your doctor** for advice if a pregnant woman has come into contact with Dutasteride/Tamsulosin.

## Driving and using machines

Dutasteride/Tamsulosin makes some people feel dizzy, so it may affect your ability to drive or operate machinery safely.

→ **Don't drive or operate machinery** if you are affected in this way.

**Dutasteride/Tamsulosin Capsule contains lecithin from soya and propylene glycol.**

This medicine contains lecithin from soya, which may contain soya oil. If you are allergic to peanut or soya, do not use this medicinal product.

This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

This medicine contains 299 mg propylene glycol in each capsule.

## 3. How to take Dutasteride/Tamsulosin

**Always take Dutasteride/Tamsulosin exactly as your doctor or pharmacist has told you to.** If you do not take it regularly the monitoring of your PSA levels may be affected. Check with your doctor or pharmacist if you are not sure.

## How much to take

**The recommended dose is one capsule taken once a day, 30 minutes after the same meal each day.**

## How to take

**Swallow the capsules whole with water.** Do not chew or break open the capsule. Contact with the contents of the capsules may make your mouth or throat sore.

**If you take more Dutasteride/Tamsulosin than you should**

Contact your doctor or pharmacist for advice if you take too many Dutasteride/Tamsulosin capsules.

## If you forget to take

**Dutasteride/Tamsulosin** Do not take a double dose to make up for a forgotten dose. Just take your next dose at the usual time.

**Don't stop Dutasteride/Tamsulosin without advice**

Don't stop taking Dutasteride/Tamsulosin without talking to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## 4. Possible side effects

**Like all medicines, this medicine can cause side effects, although not everybody gets them.**

## Allergic reaction

The signs of allergic reactions can include:

- skin rash (which can be itchy)
- hives (like a nettle rash)
- swelling of the eyelids, face, lips, arms or legs

→ **Contact your doctor immediately** if you get any of these symptoms, and **stop using Dutasteride/Tamsulosin.**

## Dizziness, light-headedness and fainting

Dutasteride/Tamsulosin can cause dizziness, lightheadedness and on rare occasions fainting. Take care when moving from a lying down or sitting position to sitting or standing, particularly if you wake up in the night, until you know how this medicine affects you. If you feel dizzy or lightheaded at any time during treatment, **sit or lie down until the symptoms pass.**

## Serious skin reactions

The signs of serious skin reactions can include:

- a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (Stevens-Johnson syndrome).

→ **Contact a doctor immediately** if you get these symptoms and **stop using Dutasteride/Tamsulosin.**

## Common side effects

These may affect up to 1 in 10 men taking Dutasteride/Tamsulosin:

- impotence (not able to achieve or maintain an erection)\*
- decreased sex drive (libido)\*
- difficulty with ejaculation, such as a decrease in the amount of semen released during sex\*
- breast enlargement or tenderness (gynecomastia)
- dizziness
- \*In a small number of people some of these events may continue after you stop taking Dutasteride/Tamsulosin.

## Uncommon side effects

These may affect up to 1 in 100 men

- heart failure (heart becomes less efficient at pumping blood around the body. You may have symptoms such as shortness of breath, extreme tiredness and swelling in your ankles and legs)
- low blood pressure on standing
- fast heart beat (palpitations)
- constipation, diarrhoea, vomiting, feeling sick (nausea)
- weakness or loss of strength
- headache
- itchy, blocked or runny nose (rhinitis)
- skin rash, hives, itching
- hair loss (usually from the body) or hair growth

## Rare side effects

These may affect up to 1 in 1,000 men

- swelling of the eyelids, face, lips, arms or legs (*angioedema*)
- fainting

## Very rare side effects

These may affect up to 1 in 10,000 men

- persistent painful erection of the penis (*priapism*)
- serious skin reactions (*Stevens-Johnson syndrome*)

## Other side effects

Other side effects have occurred in a small number of men, but their exact frequency is not known (the frequency cannot be estimated from the available data):

- abnormal or fast heartbeat (*arrhythmia* or *tachycardia* or *atrial fibrillation*)
- shortness of breath (*dyspnoea*)
- depression
- pain and swelling in your testicles
- nose bleeds
- severe skin rash
- changes in vision (*blurred vision* or *visual impairment*)
- dry mouth

## Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme website [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Dutasteride/Tamsulosin

Keep this medicine out of the sight and reach of children.

Do not use Dutasteride/Tamsulosin after the expiry date which is stated on the carton and the bottle after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage condition.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

## What Dutasteride/Tamsulosin Capsule contains

The active substances are dutasteride and tamsulosin hydrochloride. Each capsule contains 0.5 mg dutasteride and 0.4 mg tamsulosin hydrochloride. (equivalent to 0.367 mg of tamsulosin).

The other ingredients are:

## Hard capsule shell:

Black iron oxide (E172)  
Red iron oxide (E172)  
Titanium dioxide (E171)  
Yellow iron oxide (E172)  
Gelatin

## Contents in dutasteride soft capsule:

Propylene glycol monooctylate, type II  
Butylhydroxytoluene (E321)

## Soft Capsule Shell:

Gelatin  
Glycerol  
Titanium dioxide (E171)  
Triglycerides (medium chain)  
Lecithin (may contain soya oil) (see section 2)

## Tamsulosin pellets:

Methacrylic acid - ethyl acrylate copolymer 1:1 dispersion 30 per cent (contains sodium laurylsulfate, polysorbate 80)  
Microcrystalline cellulose  
Dibutyl sebacate  
Polysorbate 80  
Colloidal hydrated silica  
Calcium stearate

## Black ink:

Shellac (E904)  
Black iron oxide (E172)  
Propylene glycol (E1520)  
Concentrated ammonia solution (E527)  
Potassium hydroxide (E525)

See section 2 "Dutasteride/Tamsulosin Capsule contains lecithin from soya and propylene glycol".

## What Dutasteride/Tamsulosin Capsule looks like and contents of the pack

This medicine is an oblong hard gelatin capsule, of 24.2 mm x 7.7 mm approx., with brown body and beige cap printed with C001 in black ink.

They are available in packs of 7, 30 and 90 capsules.

Not all pack sizes may be marketed.

## Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:  
Dr. Reddy's Laboratories (UK) Ltd.,  
6 Riverview Road, Beverley, East Yorkshire,  
HU17 0LD, United Kingdom

Manufacturer:  
Laboratorios LEÓN FARMA, SA  
C/La Vallina, s/n, Polígono Industrial  
Navatejera  
Villaquilambre -24008 (León)  
SPAIN

This leaflet was last revised in 01/2020

## Artwork

Brand:	Dr Reddy's
Product Name:	Dutastride Tamsulosine
Strength:	0,5mg/0,4mg
Form:	Capsules, Hard
Component:	Leaflet
Pack Size:	30's
Country:	UK
Date Created:	26 JUL 2019
Date Modified:	22 JAN 2020
Version:	1.3

## Technical Information

Die Cut	Guides
---------	--------

Min Font Size:	9pt
Cutter Ref	DUTA TAMSU CHEMO 540x140_PIL
Printer:	CHEMO
Dimensions:	540 x 140mm
Project:	NPL - Change Control EXC 016/2020

## Colours

Process Black
---------------

**Dr.Reddy's**  
Good Health Can't Wait

Dr. Reddy's Laboratories (UK) Ltd  
6 Riverview Road, Beverley, HU17 0LD, UK