

experience some abdominal discomfort. It is unlikely to

cause you any harm, but please consult your doctor or

If symptoms persist after 7 days consult your doctor. 4. <u>Possible side effects</u> Very rarely (less than 1 in 10,000 patients) there is a chance of an allergic reaction to the ingredients.

Symptoms of this may include skin rash, itching, difficulty breathing, dizziness or swelling of the face, lips, tongue or throat. If you experience these or any other side effects stop taking the product and consult your doctor immediately. If you get any side effects, talk to your

doctor, pharmacist or nurse. This includes any possible

side effects not listed in this leaflet. You can also report

side effects directly via the Yellow Card Scheme at: "http://www.mhra.gov.uk/yellowcard" or search for MHRA Yellow Card in the Google Play or Apple App

Store. By reporting side effects, you can help provide

(EXP: month/year) shown. Keep out of the sight and reach

more information on the safety of this medicine.

5. How to store this medicine Do not use this product after the expiry date

pharmacist if this doesn't go away.

of children.

Do not refrigerate

help to protect the environment.

180ml, 200ml, 250ml and 300ml.

Hull, HU87DS.

Not all pack sizes may be marketed.

Date of preparation: November 2020

Manufacturer and Product Licence Holder: Reckitt Benckiser Healthcare (UK) Limited,

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will

6. Contents of the pack and other information Each 10 ml of oral suspension contains 1000 mg sodium

alginate and 200 mg potassium hydrogen carbonate as the active ingredients. The other ingredients are calcium carbonate, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, sodium

hydroxide, peppermint flavour and purified water. This product does not contain sugar or gluten. this medicine is available in bottles of 80ml, 100ml, 125ml, 140ml, 150ml,

Gaviscon is a trademark of the RB group of companies

1.3.2 Mock-up Label-Leaflet - Clean

(a medication to treat prostate cancer). **Pregnancy and breast-feeding:**You can take this product if you are pregnant or breast-

feeding. As with all medicines, the treatment duration should be limited as much as possible.

Important information about some of the ingredients of Gaviscon Advance Peppermint flavour Oral Suspension This product contains methyl (E218) and propyl (E216)

This medicine has no influence on the ability to drive and use machines.

Check that the cap seal is unbroken before first using this

product. Shake well before use. Adults including the

elderly and children 12 years and over: 5-10 ml (one to two 5ml spoonfuls) after meals and at bedtime. Children

under 12 years: Should only be taken on medical advice.

If you forget a dose it is not necessary to double the dose

much of this product you may feel bloated and

para-hydroxybenzoates which may cause allergic

reactions (possibly delayed). **Driving and using machines:**

3. How to take this medicine