

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed on this carton. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

92776792401 GB

SUPPOSITORIES

12

> Internal Relief
> Contains local anaesthetic

SOOTHES PAIN, RELIEVES ITCHING
& PROMOTES HEALING OF PILES

Haemorrhoid Relief
Suppositories
Allantoin, Lidocaine Hydrochloride Monohydrate

Boots

26-03-063



If you need more advice
ask your pharmacist.

Boots

Haemorrhoid Relief
Suppositories
Allantoin, Lidocaine
Hydrochloride Monohydrate



How to store this medicine
Store at or below 25°C.
Keep all medicines out of the sight and reach of children.
Use by the date on the side panel of the carton.
Active ingredients: Each suppository contains Allantoin 10.25 mg, Lidocaine Hydrochloride Monohydrate 10.25 mg.

Also contains: colloidal silicon dioxide, witepsol w35, purified water.
PL 00240/0071 UK Only
Manufactured for The Boots Company PLC Nottingham NG2 3AA by the Marketing Authorisation holder 8 Ross Ltd Linthwaite Huddersfield West Yorkshire HD7 5QH
Text prepared 03/2024

Read all of this carton for full instructions.

Uses: This medicine contains a local anaesthetic, Lidocaine, to relieve the pain and discomfort of haemorrhoids (also known as piles). It also contains Allantoin which helps healing.

Before you use this medicine

✗ Do not use:

- If you are allergic to any of the ingredients
- **Talk to your pharmacist or doctor:**
 - If you have continuous pain or bleeding from the anus, a change in bowel habit, a swollen stomach or have lost weight without trying
 - If you are pregnant or breastfeeding

✓ How to use this medicine

Check the blister is not broken before use. If it is, do not use that suppository.

Wash hands thoroughly before and after inserting the suppository. Remove the protective film.

Adults and the elderly: If possible empty bowels before use. Gently insert one suppository deep into the rectum morning and night and after each bowel movement.

Use only in the rectum. Do not swallow the suppository. Do not use this medicine for more than 7 days, unless your doctor tells you to.

If symptoms do not go away within 7 days talk to your doctor.

- ! **If anyone accidentally swallows a suppository:**
Talk to a doctor straight away.

Possible side effects

Most people will not have problems, but some may get some.

- ! **If you get any of these serious side effects, stop using the suppositories. See a doctor:**
 - Allergic reactions (e.g. skin rash, redness or itchy skin)
- If you get any of the following side effects see your pharmacist or doctor:**
 - Diarrhoea

UK ONLY



BOX Recycle



BLISTER Do Not Recycle