

**Package leaflet: Information for the user**  
**REKAMBYS 900 mg prolonged-release suspension for injection**  
rilpivirine

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What is REKAMBYS and what it is used for
2. What you need to know before you use REKAMBYS
3. How REKAMBYS is given
4. Possible side effects
5. How to store REKAMBYS
6. Contents of the pack and other information

**1. What REKAMBYS is and what it is used for**

REKAMBYS contains the active ingredient rilpivirine. It is one of a group of medicines called non-nucleoside reverse transcriptase inhibitors (NNRTIs) that are used for the treatment of human immunodeficiency virus type 1 (HIV-1) infection.

REKAMBYS works together with other HIV medicines to block the ability of the virus to make more copies of itself. REKAMBYS injections do not cure HIV infection but help reduce the amount of HIV in your body and keeps it at a low level. This holds off damage to the immune system and the development of infections and diseases associated with AIDS.

REKAMBYS is always given with another HIV medicine called cabotegravir injection. They are used together in adults and adolescents (at least 12 years of age and weighing at least 35 kg) whose HIV-1 infection is already under control.

**2. What you need to know before you use REKAMBYS**

**Do not use REKAMBYS** if you are allergic to rilpivirine or any of the other ingredients of this medicine (listed in section 6).

**Do not use REKAMBYS if you are taking any of the following medicines** as they may affect the way REKAMBYS or the other medicine works:

- carbamazepine, oxcarbazepine, phenobarbital, phenytoin (medicines to treat epilepsy and prevent seizures)
- rifabutin, rifampicin, rifapentine (medicines to treat some bacterial infections such as tuberculosis)
- dexamethasone (a corticosteroid used in a variety of conditions such as inflammation and allergic reactions) as a course of treatment by mouth or injection

- products that contain St John's wort (*Hypericum perforatum*, a herbal remedy used for depression).

If you are taking any of the above, ask your doctor about alternatives.

### **Warnings and precautions**

Talk to your doctor or pharmacist before using REKAMBYS.

REKAMBYS is not a cure for HIV infection. It is part of a treatment to reduce the amount of virus in the blood.

### **Tell your doctor about your situation**

Check the following points and tell your doctor if any of them apply to you.

- You must attend all the planned visits for injections, do not miss any visits, it is very important for the success of your treatment. If you cannot attend a planned visit, inform your doctor as soon as possible.
- Tell your doctor if you have ever had **problems with your liver**, including hepatitis B or hepatitis C, or **problems with your kidneys**. Your doctor may check how well your liver or kidneys work to decide if you can use REKAMBYS. See 'Uncommon side effects' in section 4 of this leaflet for signs of liver damage.
- Tell your doctor immediately if you notice any **symptoms of infections** (for example, fever, chills, sweats). In some patients with HIV, inflammation from previous infections may occur soon after starting HIV treatment. It is believed that these symptoms are due to an improvement in the body's immune response, enabling the body to fight infections that were present previously but caused no obvious symptoms.
- Also tell your doctor straight away if you notice any symptoms such as muscle weakness, weakness beginning in the hands and feet and moving up towards the trunk of the body, palpitations, tremor or hyperactivity. This is because autoimmune disorders (conditions in which the immune system mistakenly attacks healthy body tissue) may also occur after you start taking medicines for the treatment of your HIV infection. Autoimmune disorders may occur many months after the start of treatment.
- Tell your doctor if you are taking any medicines that you have been told may cause a life-threatening irregular heartbeat (torsade de pointes).

### **Reactions to Injections**

Post-injection reaction symptoms have happened within minutes in some people after receiving their rilpivirine injection. Most symptoms resolved within a few minutes after the injection. Symptoms of post-injection reactions may include: difficulty breathing, stomach cramps, rash, sweating, numbness of your mouth, feeling anxious, feeling warm, feeling lightheaded or feeling like you are going to pass out (faint), blood pressure changes, and pain (e.g., back and chest). Tell your healthcare professional if you experience these symptoms after you receive your injections.

### **Regular appointments are important**

It is important that you **attend your planned appointments** to receive REKAMBYS, to control your HIV infection and to stop your illness from getting worse. Do not miss any visits, it is very important for the success of your treatment. If you cannot attend a planned visit, inform your doctor as soon as possible. Talk to your doctor if you are thinking about stopping treatment. If you are late receiving your REKAMBYS injection, or if you stop receiving REKAMBYS, you will need to take other medicines to treat HIV infection and to reduce the risk of the virus becoming resistant as the drug levels in your body will be too low to treat the HIV infection.

### **Children**

REKAMBYS is not for use in children less than 12 years of age or adolescents weighing less than 35 kg, because it has not been studied in these patients.

### Other medicines and REKAMBYS

Tell your healthcare provider if you are taking, have recently taken or might take any other medicines. Some medicines may affect the levels of REKAMBYS in the blood if you are taking them while being treated with REKAMBYS, or REKAMBYS may affect how well the other medicine works.

**REKAMBYS must not be given with some other medicines (see ‘Do not use REKAMBYS’ in section 2).**

**The effects of REKAMBYS or other medicines might change** if you use REKAMBYS together with any of the following medicines:

- clarithromycin, erythromycin (antibiotics)
- methadone (used to treat narcotic withdrawal and dependence)

If you are taking any of the above, ask your doctor about alternatives.

### Pregnancy and breast-feeding

Tell your doctor immediately if you are pregnant or if you plan to become pregnant. Your doctor will consider the benefit and the risk to you and your baby of using REKAMBYS while you are pregnant. If you are planning to have a baby, talk to your doctor in advance, as rilpivirine can remain in your body for up to 4 years after the last injection of REKAMBYS.

Breast-feeding is not recommended in women living with HIV because HIV infection can be passed on to the baby in breast milk.

If you are breast-feeding, or thinking about breast-feeding, you should discuss it with your doctor as soon as possible.

### Driving and using machines

Some patients may feel tired, dizzy or drowsy during treatment with REKAMBYS. Do not drive or operate machinery if you have any of these side effects.

### REKAMBYS contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per 3 mL injection, that is to say essentially ‘sodium-free’.

## 3. How REKAMBYS is given

A nurse or doctor will give you REKAMBYS as an injection in the muscle of your buttock (*intramuscular, or IM injection*).

You will be given your injection **either once every month or once every 2 months**, together with another injectable medicine called cabotegravir. Your doctor will explain how often the medicine will be given.

**When you start treatment with REKAMBYS**, you and your doctor may decide to start with daily treatment of one 25 mg rilpivirine tablet with a meal and one 30 mg cabotegravir tablet for one month before your first REKAMBYS injection. This is called the *lead-in period* - taking the tablets before you receive REKAMBYS and cabotegravir injections will allow your doctor to test how well these medicines suit you.

The other option is that you and your doctor may decide to start directly with REKAMBYS injections.

If you are going to be given REKAMBYS every month, your treatment will be as follows:

	When	
Medicine	First injection	Second injection onwards, every month
Rilpivirine	single injection of 900 mg	600 mg by injection every month

Cabotegravir	single injection of 600 mg	400 mg by injection every month
--------------	----------------------------	---------------------------------

If you are going to be given REKAMBYS every 2 months, your treatment will be as follows:

Medicine	When	
	First and second injections, one month apart	Third injection onwards, every two months
Rilpivirine	single injection of 900 mg	900 mg by injection, every 2 months
Cabotegravir	single injection of 600 mg	600 mg by injection, every 2 months

### **If you miss a REKAMBYS injection**

It is important that you keep your regular planned appointments to receive your injection. If you miss an appointment, contact your doctor immediately to make a new appointment.

**Talk to your doctor** if you think you will not be able to receive your REKAMBYS injection at the usual time. Your doctor may recommend you take tablets instead, until you are able to have a REKAMBYS injection again.

### **If you are given too much REKAMBYS**

A doctor or nurse will give this medicine to you, so it is unlikely that you will be given too much. If you are worried, tell the doctor or nurse.

### **Don't stop using REKAMBYS without advice from your doctor.**

Use REKAMBYS for as long as your doctor recommends. Don't stop unless your doctor advises you to.

Low levels of rilpivirine (the active ingredient of REKAMBYS) can remain in your body for up to 4 years after stopping treatment. However, once you received your last REKAMBYS injection, the low levels of rilpivirine that remain will not work well enough against the virus which then can become resistant. To keep your HIV-1 infection under control and to stop the virus becoming resistant, you must start a different HIV treatment by the time your next REKAMBYS injection was planned.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The following is a list of side effects that have been reported when REKAMBYS is used with cabotegravir injection.

Very common side effects (affects at least 1 in 10 people)

- headache
- injection site reactions - these are generally mild to moderate and became less frequent over time. Symptoms may include:
  - very common: pain and discomfort, a hardened mass or lump
  - common: redness, itching, swelling, warmth or bruising (which may include discolouration or a collection of blood under the skin).
  - uncommon: numbness, minor bleeding, an abscess (collection of pus) or cellulitis (heat, swelling or redness).
- feeling hot/feverish (*pyrexia*), which may occur within one week after injections.

Common side effects (affects less than 1 in 10 people)

- depression
- anxiety
- abnormal dreams
- sleeping difficulty (*insomnia*)

- dizziness
- feeling sick (*nausea*)
- vomiting
- belly pain (*abdominal pain*)
- wind (*flatulence*)
- diarrhoea
- rash
- muscle pain (*myalgia*)
- tiredness (*fatigue*)
- feeling weak (*asthenia*)
- generally feeling unwell (*malaise*)
- weight gain

Uncommon side effects (affects less than 1 in 100 people)

- feeling drowsy (*somnolence*)
- feeling lightheaded, during or after an injection. This may lead to fainting.
- liver damage (signs may include yellowing of the skin and the whites of the eyes loss of appetite, itching, tenderness in the belly, light-coloured stools or unusually dark urine).
- changes in liver blood tests (increase in *transaminases*)
- an increase in *bilirubin* (a substance produced by the liver) in the blood.

Other side effects

- Severe abdominal pain caused by inflammation of the pancreas (*pancreatitis*).

The following side effects that can occur with rilpivirine tablets may also occur with REKAMBYS injection:

Very Common side effects (affects at least 1 in 10 people)

- increase in cholesterol and/or pancreatic amylase in your blood

Common side effects (affects less than 1 in 10 people)

- decreased appetite
- sleep disorders
- depressed mood
- stomach discomfort
- dry mouth
- low white blood cell and/or platelet count, decrease in haemoglobin in your blood, increase in triglycerides and/or lipase in your blood

Uncommon side effects (affects less than 1 in 100 people)

- signs or symptoms of inflammation or infection, for example fever, chills, sweats (*immune reactivation syndrome, see section 2 for more details*)

### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store REKAMBYS

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

Store in a refrigerator (2°C - 8°C). Do not freeze.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What REKAMBYS contains**

- The active substance is rilpivirine. Each 3 mL vial contains 900 mg rilpivirine.
- The excipients are poloxamer 338, citric acid monohydrate (E330), glucose monohydrate, sodium dihydrogen phosphate monohydrate, sodium hydroxide (E524) to adjust pH and ensure isotonicity, and water for injections.

### **What REKAMBYS looks like and contents of the pack**

Prolonged-release suspension for injection. REKAMBYS is presented in a glass vial. The pack also contains 1 syringe, 1 vial adaptor, and 1 injection needle.

### **Marketing Authorisation Holder**

Janssen-Cilag Ltd  
50-100 Holmers Farm Way  
High Wycombe  
Buckinghamshire  
HP12 4EG  
UK

### **Manufacturer**

Janssen Pharmaceutica NV  
Turnhoutseweg 30  
B-2340 Beerse  
Belgium

**For information in large print, tape, CD or Braille, telephone 0800 7318450.**

**This leaflet was last revised in July 2024**

**The following information is intended for healthcare professionals only:**

**REKAMBYS 3 mL injection Instructions for Use:**

**Overview**

A complete dose requires two injections:

3 mL of cabotegravir and 3 mL of rilpivirine.

Cabotegravir and rilpivirine are suspensions that do not need further dilution or reconstitution. The preparation steps for both medicines are the same. Carefully follow these instructions when preparing the suspension for injection to avoid leakage.

Cabotegravir and rilpivirine are for intramuscular use only. Both injections must be administered to separate gluteal injection sites.

**Note:** The ventrogluteal site is recommended. **The administration order is not important.**



**Storage information**

- Store in refrigerator at 2°C to 8°C.


**Do not freeze.**

**Your pack contains**

- 1 vial of rilpivirine
- 1 vial adaptor
- 1 syringe
- 1 injection needle (23 gauge, 1½ inch)

Consider the patient's build and use medical judgment to select an appropriate injection needle length.

**Rilpivirine vial**




**Vial cap**  
(Rubber stopper under cap)

**Vial adaptor**




**Syringe**



**Plunger**

**Injection needle**

**Needle guard**



**Needle cap**

## You will also need

- Non-sterile gloves
- 2 alcohol swabs
- 2 gauze pads
- A suitable sharps container

**+** 1 Cabotegravir 3 mL pack




Make sure to have the cabotegravir pack close by before starting.

## Preparation

### 1. Inspect vial

**Check expiry date  
and medicine**




- Check that the expiry date has not passed.
- Inspect the vials immediately. If you can see foreign matter, do not use the product.

**Do not** use if the expiry date has passed.

### 2. Wait 15 minutes



**Wait 15 minutes**



- Wait at least 15 minutes before you are ready to give the injection to allow the medicine to come to room temperature.




### 3. Shake vigorously



- Hold the vial firmly and vigorously shake for a full 10 seconds as shown.


### 4. Inspect suspension



- Invert the vial and check the resuspension. It should look uniform. If the suspension is not uniform, shake the vial again.
- It is also normal to see small air bubbles.

**Note:** Vial preparation order is not important.


### 5. Remove vial cap



- Remove the cap from the vial.
- Wipe the rubber stopper with an alcohol swab.

**Do not** allow anything to touch the rubber stopper after wiping it.


### 6. Peel open vial adaptor



- Peel off the paper backing from the vial adaptor packaging.

**Note: Do not** remove the adaptor from in its packaging for the next step. The adaptor **will not** fall out when the packaging is turned upside down.

## 7. Attach vial adaptor




- Place the vial on a flat surface.
- Press the vial adaptor straight down onto the vial, as shown.
- The vial adaptor should click securely into place.

## 8. Lift off the packaging




- Lift off the vial adaptor packaging, as shown.

## 9. Prepare syringe








- Remove the syringe from its packaging.
- Draw 1 mL of air into the syringe. This will make it easier to draw up the liquid later.

## 10. Attach syringe




- Hold the vial adaptor and vial firmly, as shown.
- Screw the syringe firmly onto the vial adaptor.

## 11. Press the plunger

	<ul style="list-style-type: none"> <li>• Press the plunger all the way down to push the air into the vial.</li> </ul>
<b>12. Slowly draw up dose</b>	
	<ul style="list-style-type: none"> <li>• Invert the syringe and vial, and slowly withdraw as much of the liquid as possible into the syringe. There might be more liquid than dose amount.</li> </ul> <p>  <b>Note:</b> Keep the syringe upright to avoid leakage.         </p>
<b>13. Unscrew syringe</b>	
	<ul style="list-style-type: none"> <li>• Hold the syringe plunger firmly in place as shown to prevent leakage. It is normal to feel some back pressure.</li> <li>• Screw the syringe off the vial adaptor, holding the vial adaptor as shown.</li> </ul> <p>  <b>Note:</b> Check that the suspension looks uniform and milky white.         </p>


## 14. Attach needle



- Peel open the needle packaging part way to expose the needle base.
- Keeping the syringe upright, firmly twist the syringe onto the needle.
- Remove the needle packaging from the needle.

## Injection

## 15. Prepare injection site



Ventrogluteal


Dorsogluteal

Injections must be administered to the gluteal sites. Select from the following areas for the injection:

- Ventrogluteal (recommended)
- Dorsogluteal (upper outer quadrant)


**Note:** For gluteal intramuscular use only.  
**Do not** inject intravenously.

## 16. Remove cap



- Fold the needle guard away from the needle.
- Pull off the injection needle cap.


## 17. Remove extra liquid



- Hold the syringe with the needle pointing up. Press the plunger to the 3 mL dose to remove extra liquid and any air bubbles.

**Note:** Clean the injection site with an alcohol swab. Allow the skin to air dry before continuing.


## 18. Stretch skin



Use the z-track injection technique to minimise medicine leakage from the injection site.


- Firmly drag the skin covering the injection site, displacing it by about 2.5 cm (1 inch).
- Keep it held in this position for the injection.

## 19. Insert needle



- Insert the needle to its full depth, or deep enough to reach the muscle.

## 20. Inject dose



- Still holding the skin stretched – slowly press the plunger all the way down.
- Ensure the syringe is empty.
- Withdraw the needle and release the stretched skin immediately.


## 21. Assess the injection site



- Apply pressure to the injection site using a gauze.
- A small bandage may be used if a bleed occurs.

**Do not** massage the area.


## 22. Make needle safe



- Fold the needle guard over the needle.
- Gently apply pressure using a hard surface to lock the needle guard in place.
- The needle guard will make a click when it locks.

## After injection

## 23. Dispose safely



- Dispose of used needles, syringes, vials and vial adaptors according to local health and safety laws.

## Repeat for 2nd medicine



**Repeat all steps  
for 2nd medicine**

If you have not yet injected cabotegravir follow its own specific instructions for use for preparation and injection of the medicine.

## Questions and Answers

### 1. How long can the medicine be left out of the refrigerator?

It is best to inject the medicine as soon as it reaches room temperature. However, the vial may sit in the carton at room temperature (maximum temperature of 25°C) for up to 6 hours; do not put back into the refrigerator. If not used within 6 hours, the vial must be discarded.

### 2. How long can the medicine be left in the syringe?

It is best to inject the (room temperature) medicine as soon as possible after drawing it up. However, the medicine can remain in the syringe for up to 2 hours before injecting.

If 2 hours are exceeded, the medicine, syringe and needle must be discarded.

### 3. Why do I need to inject air into the vial?

Injecting 1 mL of air into the vial makes it easier to draw up the dose into the syringe. Without the air, some liquid may flow back into the vial unintentionally, leaving less than intended in the syringe.

### 4. Does the order in which I give the medicines matter?

No, the order is unimportant.

### 5. Is it safe to warm the vial up to room temperature more quickly?

It is best to let the vial come to room temperature naturally. However, you can use the warmth of your hands to speed up the warm up time, but make sure the vial does not get above 25°C.

Do not use any other heating methods.

### 6. Why is the ventrogluteal administration approach recommended?

The ventrogluteal approach, into the gluteus medius muscle, is recommended because it is located away from major nerves and blood vessels. A dorsogluteal approach, into the gluteus maximus muscle, is acceptable, if preferred by the healthcare professional. The injection should not be administered in any other site.