



HEARTBURN, INDIGESTION & WIND RELIEF
680 mg/80 mg/25 mg Chewable Tablets



Calcium Carbonate, Magnesium Carbonate, Simeticone

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must contact a doctor if you do not feel better or if you feel worse after 14 days.

WHAT IS IN THIS LEAFLET

1. What Rennie Heartburn, Indigestion & Wind Relief is and what it is used for
2. What you need to know before you take Rennie Heartburn, Indigestion & Wind Relief
3. How to take Rennie Heartburn, Indigestion & Wind Relief
4. Possible side effects
5. How to store Rennie Heartburn, Indigestion & Wind Relief
6. Contents of the pack and other information

1. WHAT RENNIE® HEARTBURN, INDIGESTION & WIND RELIEF IS AND WHAT IT IS USED FOR

Rennie Heartburn, Indigestion & Wind Relief works quickly and effectively in two ways:

1. Antacids (calcium carbonate and magnesium carbonate) get to work within minutes, powerfully neutralising excess acid in the stomach.
2. Simeticone, a highly effective anti-foaming agent, causes the trapped air bubbles to combine and disperse, helping to eliminate the symptoms discreetly and without embarrassment.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE RENNIE® HEARTBURN, INDIGESTION & WIND RELIEF

Do not take if you:

- Have severe kidney disease.
- Have high calcium in the urine or kidney stones.
- Have high calcium or low phosphate levels in the blood.
- Are allergic (hypersensitive) to any of the ingredients (listed in Section 6).

Warnings and precautions:

Please consult your doctor if you suffer frequently or need to use this medicine for more than 14 days, in order to rule out more serious conditions.

Please consult your doctor or pharmacist if you have mild or moderate kidney problems as regular blood tests should be carried out.

Other medicines and Rennie Heartburn, Indigestion & Wind Relief:

- Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.
- If you are taking any of the following medicines speak to your doctor or pharmacist before taking Rennie Heartburn, Indigestion & Wind Relief because it can affect how these medicines work.
 - Tetracyclines and quinolones (antibiotics)
 - Bisphosphonates (used for the treatment of osteoporosis)
 - Dolutegravir (an anti-retroviral medicine used for the treatment of HIV infection)
 - Levothyroxine (a thyroid hormone)
 - Eltrombopag (used to increase the number of platelets)
 - Digoxin (used for some heart conditions)
 - Thiazide diuretics (water tablets to treat fluid retention and high blood pressure)
 - Fluorides (for preventing dental caries)
 - Phosphates (for clearing the bowel before a surgical procedure)
 - Iron supplements
- To maximise the benefit of all medicines being used, take Rennie Heartburn, Indigestion & Wind Relief at least 4 hours before or after taking eltrombopag and 1 to 2 hours after taking any other medicines.

Rennie Heartburn, Indigestion & Wind Relief with food and drink

Avoid taking this medicine with large amounts of milk or dairy products.

Pregnancy and breastfeeding:

Rennie Heartburn, Indigestion & Wind Relief can be used during pregnancy and breastfeeding if taken as instructed.

7001455 / 1122GB

As this product provides a substantial amount of calcium (272 mg per tablet) in addition to dietary calcium intake, pregnant women should strictly limit their use of this product to the maximum recommended daily dose and avoid concomitant, excessive intake of milk (1 litre contains up to 1.2 g elemental calcium) and dairy products.

Rennie Heartburn, Indigestion & Wind Relief contains sorbitol

This product contains 430 mg sorbitol in each tablet.

Sorbitol is a source of fructose. If your doctor has told you that you (or your child) have an intolerance to some sugars or if you have been diagnosed with hereditary fructose intolerance (HFI), a rare genetic disorder in which a person cannot break down fructose, talk to your doctor before you (or your child) take or receive this medicine.

3. HOW TO TAKE RENNIE® HEARTBURN, INDIGESTION & WIND RELIEF

Adults and children over 12 years only: Suck or chew 1-2 tablets, up to a maximum of 11 tablets a day.

Do not take more than the label tells you to.

Children under 12 years: Not recommended.

If symptoms persist after 14 days consult your pharmacist or doctor. Prolonged use should be avoided.

If you take more Rennie Heartburn, Indigestion & Wind Relief than you should:

Drink plenty of water and consult your doctor or pharmacist. Symptoms of an overdose include nausea and vomiting, tiredness, increased urine production, increased thirst, dehydration and abnormal muscular weakness.

If you have any questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them. Side effects are unlikely at the recommended dose.

If you experience the following side effects stop treatment immediately and seek medical advice.

Rarely allergic reactions to ingredients have been reported, e.g. rashes, itching, difficulty in breathing and swelling of the face, mouth or throat and anaphylactic shock (anaphylactic shock is a severe sudden allergic reaction, symptoms of which are low blood pressure, shock, palpitations, difficulty in breathing, bronchospasm, skin reactions, abdominal pain or cramps, vomiting, and diarrhoea).

Long term use of high doses can cause high blood levels of calcium and magnesium, especially in people with kidney conditions. Symptoms of this may include nausea, vomiting, upset stomach, constipation, diarrhoea, tiredness, muscular weakness, headache, kidney problems and impaired sense of taste. In exceptional cases, long term use of high doses can lead to milk alkali syndrome, which can cause high blood levels of calcium.

Reporting of side effects:

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at:

www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE RENNIE® HEARTBURN, INDIGESTION & WIND RELIEF

Keep this medicine out of the sight and reach of children. Do not store above 25°C and keep in the original pack.

Do not use this medicine after the expiry date which is stated on the carton and blister. The expiry date refers to the last day of that month.

Do not throw away any medicines via waste water or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Rennie Heartburn, Indigestion & Wind Relief contains:

The active substances are: simeticone 25mg, calcium carbonate 680mg and heavy magnesium carbonate 80mg. The other ingredients are sorbitol (E420), magnesium stearate, potato starch, pregelatinised maize starch, talc, mint flavour, lemon flavour.

What Rennie Heartburn, Indigestion & Wind Relief looks like and contents of the pack:

Rennie Heartburn, Indigestion & Wind Relief are cream white flat round tablets, scored on one side.

They are available in packs of 2, 4, 6, 12, 16, 18, 20, 24, 36, 48, 60, 96 tablets. Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer:

Marketing Authorisation Holder: Bayer plc, 400 South Oak Way, Reading, RG2 6AD, UK. consumerhealthuk@bayer.com

Manufacturer: Delpharm Gaillard, 74240 Gaillard, France.

This leaflet was last revised in January 2022. Bayer