


85 mm

Pharmacode Reading Direction



360 mm

PACKAGE LEAFLET:
INFORMATION FOR THE USER

MORNINGSIDE
HEALTHCARE

Prucalopride 1 mg Film-coated Tablets
Prucalopride 2 mg Film-coated Tablets
Prucalopride succinate

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The names of your medicines are Prucalopride 1 mg Film-coated Tablets and Prucalopride 2 mg Film-coated Tablets (called Prucalopride Tablets in the rest of this leaflet).

What is in this leaflet:

1. What Prucalopride Tablets are and what they are used for
2. What you need to know before you take Prucalopride Tablets
3. How to take Prucalopride Tablets
4. Possible side effects
5. How to store Prucalopride Tablets
6. Contents of the pack and other information.

Prucalopride Tablets with food and drink
Prucalopride Tablets can be taken with or without food and drinks, at any time of the day.

Pregnancy and breast-feeding
Prucalopride Tablets are not recommended for use during pregnancy.

- Tell your doctor if you are pregnant or planning to become pregnant.
- Use a reliable method of contraception while you're taking Prucalopride Tablets, to prevent pregnancy.
- If you do become pregnant during treatment with Prucalopride Tablets, tell your doctor.

When breast-feeding, prucalopride can pass into breast milk. Breast-feeding is not recommended during treatment with Prucalopride Tablets. Talk to your doctor about this.

Ask your doctor for advice before taking any medicine.

Driving and using machines
Prucalopride Tablets are unlikely to affect your ability to drive or use machines. However, sometimes Prucalopride may cause dizziness and tiredness, especially on the first day of treatment, and this may have an effect on driving and use of machines.

Prucalopride Tablets contain lactose
This medicine contains lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Prucalopride Tablets

Always take this medicine exactly as described in this leaflet or as your doctor has told you. Check with your doctor or pharmacist if you are not sure. Take Prucalopride Tablets every day for as long as your doctor prescribes it.

The doctor may want to reassess your condition and the benefit of continued treatment after the first 4 weeks and thereafter at regular intervals.

The recommended dose of Prucalopride Tablets for most patients is one 2 mg tablet once a day.

If you are older than 65 years or have severe liver disease, the starting dose is one 1 mg tablet once a day, which your doctor may increase to 2 mg once a day if needed.

Your doctor may also recommend a lower dose of one 1 mg tablet daily if you have severe kidney disease.

Taking a higher dose than recommended will not make the product work better.

Prucalopride Tablets are only for adults and should not be taken by children and adolescents up to 18 years.

If you take more Prucalopride Tablets than you should
It is important to keep to the dose as prescribed by your doctor. If you have taken more Prucalopride Tablets than you should, it is possible that you will get diarrhoea, headache and/or nausea. In case of diarrhoea, make sure that you drink enough water.

If you forget to take Prucalopride Tablets
Do not take a double dose to make up for a forgotten tablet. Just take your next dose at the usual time.

If you stop taking Prucalopride Tablets
If you stop taking Prucalopride Tablets, your constipation symptoms may come back again.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. The side effects mostly occur at the start of treatment and usually disappear within a few days with continued treatment.

52 mm

30 mm

1. What Prucalopride Tablets are and what they are used for

Prucalopride Tablets contain the active substance prucalopride.

Prucalopride Tablets belong to a group of gut motility enhancing medicines (gastrointestinal prokinetics). It acts on the muscle wall of the gut, helping to restore the normal functioning of the bowel. Prucalopride Tablets are used for the treatment of chronic constipation in adults in whom laxatives do not work well enough.

Not for use in children and adolescents younger than 18 years.

2. What you need to know before you take Prucalopride Tablets

Do not take Prucalopride Tablets if you:

- are allergic to prucalopride or any of the other ingredients of this medicine (listed in section 6).
- are on renal dialysis.
- suffer from perforation or obstruction of the gut wall, severe inflammation of the intestinal tract, such as Crohn's disease, ulcerative colitis or toxic megacolon/megarectum.

Warnings and precautions
Talk to your doctor or pharmacist before taking Prucalopride Tablets.

Take special care with Prucalopride Tablets and tell your doctor if you:

- suffer from severe kidney disease,
- suffer from severe liver disease,
- are currently under supervision by a doctor for a serious medical problem such as lung or heart disease, nervous system or mental health problems, cancer, AIDS or a hormonal disorder.

If you have very bad diarrhoea, the contraceptive pill may not work properly and the use of an extra method of contraception is recommended. See the instructions in the patient leaflet of the contraceptive pill you are taking.

Other medicines and Prucalopride Tablets
Tell your doctor if you are taking, or have recently taken, or might take any other medicines.

170 mm

5 mm

85 mm

30 mm
52 mm

The following side effects have been reported very commonly (may affect more than 1 in 10 people): headache, feeling sick, diarrhoea and abdominal pain.

The following side effects have been reported commonly (may affect up to 1 in 10 people): decreased appetite, dizziness, vomiting, disturbed digestion (dyspepsia), windiness, abnormal bowel sounds, tiredness.

The following uncommon side effects have also been seen (may affect up to 1 in 100 people): tremors, pounding heart, rectal bleeding, increase in frequency of passing urine (pollakiuria), fever and feeling unwell. If pounding heart occurs, please tell your doctor.

Reporting of side effects
If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via Yellow Card Scheme on the MHRA website www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Prucalopride Tablets

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and carton. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Prucalopride Tablets contain
The active substance is prucalopride.
Each film-coated tablet contains either 1 mg or 2 mg prucalopride (as succinate).

The other ingredients are lactose monohydrate (see section 2), cellulose microcrystalline, silica colloidal anhydrous and magnesium stearate, hypromellose, titanium dioxide, macrogol and triacetin. The 2 mg tablet also contains iron oxide red, iron oxide yellow, indigo carmine aluminium lake.

What Prucalopride Tablets look like and contents of the pack
Prucalopride 1 mg Film-coated Tablets are white coloured, round shaped, biconvex film-coated tablets, debossed with "31" on one side and plain on the other side.

Prucalopride 2 mg Film-coated Tablets are pink coloured, round shaped, biconvex film-coated tablets, debossed with "32" on one side and plain on the other side.

Prucalopride Tablets are provided in OPA/Alu/PVC-Alu blister in packs of 7, 14, 28 or 84 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder
Morningside Healthcare Ltd.
Unit C, Harcourt Way
Leicester, LE19 1WP
UK

Manufacturer
Morningside Pharmaceuticals Ltd.
Unit K, L, S & T2, Castle Business Park
Loughborough, LE11 5GW
UK

For information in large print, tape, CD or Braille, please contact medical enquiries on 0116 478 0322.

This leaflet was last revised in March 2022.

PP000584
M0402LAMUKNA-P1-002

5 mm

170 mm

5 mm
Pharmacode Reading Direction ←

360 mm

