

PACKAGE LEAFLET

Package leaflet: Information for the patient
Isoprenaline hydrochloride Tillomed 0.2 mg/ml solution for infusion
Isoprenaline hydrochloride Tillomed 1 mg/5ml solution for infusion

Isoprenaline hydrochloride

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Isoprenaline hydrochloride Tillomed is and what it is used for
2. What you need to know before you use Isoprenaline hydrochloride Tillomed
3. How to use Isoprenaline hydrochloride Tillomed
4. Possible side effects
5. How to store Isoprenaline hydrochloride Tillomed
6. Contents of the pack and other information

1. What Isoprenaline hydrochloride Tillomed is and what it is used for

This medicine is used in certain heart rhythm disorders and some cardiac emergencies.

2. What you need to know before you take Isoprenaline hydrochloride Tillomed

Do not take Isoprenaline hydrochloride Tillomed

- If you are allergic to the active substance or any of the other ingredients of this medicine, mentioned in section 6.
- Certain heart rhythm disorders;
- Certain forms of angina pectoris;
- In case of digitalis intoxication;

This medicine is usually not recommended in case of association with volatile halogenated anaesthetics.

Warnings and precautions

Please talk to your doctor, pharmacist or nurse before taking Isoprenaline hydrochloride Tillomed.

In case of hyperthyroidism, caution is recommended.

- In case of angina pectoris, in diabetic or digitalized patient, in case of hyperthyroidism.

Follow the advices of your doctor.

Children and adolescents

No data available

Other medicines and Isoprenaline hydrochloride Tillomed

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

Isoprenaline hydrochloride Tillomed with food and drink

No data available

Pregnancy and breast-feeding

It is preferable not to use this medicine during pregnancy.

If you discover that you are pregnant during treatment, consult your doctor as he alone can judge the need to continue it.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and ability to use machines

As this medicine will be administered to you by a healthcare professional as an emergency medicine, this does not apply to you.

Isoprenaline hydrochloride Tillomed contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per vial that is essentially “sodium-free”.

3. How to use Isoprenaline hydrochloride Tillomed

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Dosage

Dosage is adapted according to the patient’s state.

Mode of administration

Continuous intravenous route.

If you have used more Isoprenaline hydrochloride Tillomed than you should have:

As this medicine will be administered to you by a healthcare professional, it is extremely unlikely that you will receive too high or too low a dose. However, consult your doctor if you have any queries.

In the event of an overdose, you might experience an increase in the number and/or intensity of side effects. Your doctor will treat these side effects accordingly.

If you forget to use Isoprenaline hydrochloride Tillomed:

As this medicine will be administered to you by a healthcare professional, this does not apply to you.

If you stop using Isoprenaline hydrochloride Tillomed:

As this medicine will be administered to you by a healthcare professional, this does not apply to you.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Hot flashes, increased heart rate, hypotension, ventricular rhythm disorders, anginal pain, tremor, headache, nausea.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine

5. How to store Isoprenaline hydrochloride Tillomed

Keep this medicine out of the sight and reach of children.

Do not take this medicine after the expiry date which is stated on the carton box or vial after EXP. The expiry date refers to the last day of that month.

Before opening: Do not refrigerate. Keep vials in the outer carton in order to protect from light.

After opening: The product must be used immediately.

After dilution:

Chemical and physical in-use stability of the solution diluted in 5% glucose or 0.9% sodium chloride has been demonstrated for 24 hours at 25°C.

From a microbiological point of view, unless the method of opening precludes the risk of microbial contamination, the product should be used immediately. If not used immediately, in-use storage times and conditions are the responsibility of the user.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist to dispose of the medicines that you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Isoprenaline hydrochloride Tillomed contains

The active substance is Isoprenaline hydrochloride, 0.2 mg/ml and 1mg/5ml.

The other ingredients are:

Disodium Edetate, Sodium Citrate dihydrate, Citric acid anhydrous, sodium chloride, Hydrochloric acid, Sodium hydroxide, Water for injections.

What Isoprenaline hydrochloride Tillomed looks like and content of the pack

Isoprenaline hydrochloride Tillomed 0.2 mg/ml solution for infusion is a clear, colourless to slightly yellow coloured solution, free from visible particles. It is available as 1 ml fill in 2 ml clear type-I glass vial along with 13 mm rubber stopper and 13 mm flip off seal and, is available in pack sizes of 1 vial or 5 vials.

Isoprenaline hydrochloride Tillomed 1mg/5ml solution for infusion is a clear, colourless to slightly yellow coloured solution, free from visible particles. It is available as 5 ml fill in 5 ml clear type-I glass vial along with 13 mm rubber stopper and 13 mm flip off seal and, is available in pack sizes of 1 vial or 5 vials.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Tillomed Laboratories Limited
220 Butterfield, Great Marlings
Luton, LU2 8DL
United Kingdom

Manufacturer

MIAS Pharma Limited
Suite 2, Stafford House, Strand Road
Portmarnock, Co. Dublin
Ireland

This leaflet was last revised in 08/2021