


FRONT



Package leaflet: Information for the user Atenolol 25 mg film-coated tablets Atenolol 50 mg film-coated tablets Atenolol 100 mg film-coated tablets

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Atenolol is and what it is used for
2. What you need to know before you take Atenolol
3. How to take Atenolol
4. Possible side effects
5. How to store Atenolol
6. Contents of the pack and other information

1. What Atenolol is and what it is used for

Atenolol tablets contains a medicine called atenolol. This belongs to a group of medicines called beta-blockers.

Atenolol tablets are used to treat:

- Treat high blood pressure (hypertension)
- Treat uneven heart beats (arrhythmias).
- Help prevent chest pain (angina).
- Protect the heart in the early treatment after a heart attack (myocardial infarction).

It works by making your heart beat more slowly and with less force.

2. What you need to know before you take Atenolol

Do not take Atenolol

- If you are allergic to atenolol or any of the other ingredients of this medicine (listed in Section 6).
- If you have ever had any of the following heart problems:
 - heart failure which is not under control (this usually makes you breathless and causes your ankles to swell)
 - second- or third-degree heart block (a condition which may be treated by a pacemaker)
 - very slow or very uneven heart beats, very low blood pressure or very poor circulation.
- If you have a tumour called phaeochromocytoma that is not being treated. This is usually near your kidney and can cause high blood pressure. If you are being treated for phaeochromocytoma, your doctor will give you another medicine, called an alpha-blocker, to take as well as Atenolol.
- If you have been told that you have higher than normal levels of acid in your blood (metabolic acidosis).

Do not take Atenolol if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Atenolol.

Warnings and precautions

Talk to your doctor or pharmacist before taking Atenolol if:

- **You have asthma, wheezing or any other similar breathing problems, or you get allergic reactions, for example to insect stings. If you have ever had asthma or wheezing, do not take this medicine without first checking with your doctor.**
- You have a type of chest pain (angina) called Prinzmetal's angina.
- You have poor blood circulation or controlled heart failure.
- You have first-degree heart block.
- You have diabetes. Your medicine may change how you respond to having low blood sugar. You may feel your heart beating faster.
- You have thyrotoxicosis (a condition caused by an overactive thyroid gland). Your medicine may hide the symptoms of thyrotoxicosis.
- You have problems with your kidneys. You may need to have some check-ups during your treatment.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Atenolol.

Other medicines and Atenolol

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines. This includes medicines that you buy without a prescription and herbal medicines. This is because atenolol tablets can affect the way some other medicines can have an effect on atenolol.

In particular, tell your doctor if you are taking any of the following medicines:

- Verapamil, diltiazem and nifedipine (for high blood pressure or chest pain).
- Clonidine (for high blood pressure or migraine). If you are taking clonidine and Tenormin together, do not stop taking clonidine unless your doctor tells you to do so. If you have to stop taking clonidine, your doctor will give you careful instructions about how to do it.
- Digoxin (for heart problems).
- Disopyramide, quinidine or amiodarone (for an uneven heart beat).
- Adrenaline, also known as epinephrine (a medicine that stimulate the heart)
- Ibuprofen or indometacin (for pain and inflammation)
- Insulin or medicines that you take by mouth for diabetes
- Medicines to treat nose or sinus congestion or other cold remedies (including those you can buy in pharmacy).

Operations

- if you are going to a hospital to have an operation, tell the anaesthetist or medical staff that you are taking Atenolol. This is because you may get low blood pressure (hypotension) if you are given certain anaesthetics while you are taking this medicine

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

- Your medicine is not likely to affect you being able to drive or use any tools or machines. However, it is best to wait to see how your medicine affects you before trying these activities.
- If you feel dizzy or tired when taking this medicine, do not drive or use any tools or machines.

Information on sodium content

This medicine contains less than 1 mmol (23 mg) of sodium per tablet, that is to say it is essentially 'sodium-free.'

3. How to take Atenolol

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

- Your doctor will tell you how many tablets to take each day and when to take them. Read the label on the carton to remind you what the doctor said.
- Try to take your tablet at the same time each day.
- Swallow the Atenolol tablet whole with a drink of water.

Atenolol 25 mg film-coated tablets: The score line is only there to help you break the tablet if you have difficulty swallowing it whole.

Atenolol 50mg and 100mg film-coated tablets: The tablet can be divided into equal doses.

Adults

- **High blood pressure (hypertension):** the recommended dose is 50 mg to 100 mg a day.
- **Chest pain (angina):** the recommended dose is 100 mg a day or 50 mg twice a day.
- **Uneven heart beats (arrhythmias):** the recommended dose is 50 mg to 100 mg a day.
- **The early treatment of a heart attack (myocardial infarction):** the recommended dose is 50 mg to 100 mg a day.

Elderly

If you are an elderly person, your doctor may decide to give you a lower dose, particularly if you have problems with your kidneys.

People with severe kidney problems

If you have severe problems with your kidneys, your doctor may give you a lower dose.


Use in children

This medicine must not be given to children.

350.00 mm

170.00 mm

BACK



If you take more Atenolol than you should

If you take more Atenolol tablets than prescribed by your doctor, talk to a doctor or go to a hospital straight away. Take the medicine pack with you so that the tablets can be identified.

If you forget to take Atenolol

If you forget to take a dose, take it as soon as you remember. However, if it is almost time for the next dose, skip the missed dose. Do not take a double dose to make up for a forgotten dose.

If you stop taking Atenolol

Do not stop taking Atenolol tablets without talking to your doctor. In some cases, you may need to stop taking it gradually.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Allergic reactions:

If you have an allergic reaction, see a doctor straight away. The signs may include raised lumps on your skin (weals), or swelling of your face, lips, mouth, tongue or throat.

Other possible side effects

Common (may affect up to 1 in 10 people)

- You may notice that your pulse rate becomes slower while you are taking the tablets. This is normal, but if you are concerned please tell your doctor about it.
- cold hands and feet
- diarrhoea
- feeling sick (nausea).
- feeling tired

Uncommon (may affect up to 1 in 100 people)

- Disturbed sleep.

Rare (may affect up to 1 in 1,000 people)

- Heart block (which can cause dizziness, abnormal heart beat, tiredness or fainting).
- Numbness or spasm in your fingers which is followed by warmth and pain (Raynaud's disease).
- Nightmares
- Feeling confused.
- Changes in personality (psychoses) or seeing or hearing things that are not there (hallucinations)
- Headache.
- Dizziness (particularly when standing up).
- Tingling of your hands.
- Dry mouth
- Thinning of your hair
- Dry eyes
- Disturbances of vision.
- Skin rash
- Purple marks on your skin
- Being unable to get an erection (impotence)
- Yellowing of your skin or the whites of your eyes (jaundice)
- Reduced numbers of platelets in your blood (this may make you bruise more easily).
- Mood changes.

Very rare (may affect up to 1 in 10,000 people)

- Changes to some of the cells or other parts of your blood. Your doctor may take blood samples every so often to check whether Atenolol has had any effect on your blood.

Not known (cannot be estimated from the available data)

- Lupus-like syndrome (a disease where the immune system produces antibodies that attacks mainly skin and joints)
- depression

Conditions that may get worse

If you have any of the following conditions, they may get worse when you start to take your medicine. This happens rarely affecting less than 1 in 1,000 people.

- Psoriasis (a skin condition).
- Being short of breath or having swollen ankles (if you have heart failure).
- Asthma or breathing problems.
- Poor circulation.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Atenolol

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton & blister after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via waste water or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Atenolol contains

- The active substance is atenolol.
Each tablet contain 25 mg of atenolol.
Each tablet contain 50 mg of atenolol.
Each tablet contain 100 mg of atenolol.

The other ingredients are:

Tablet core: Magnesium carbonate (heavy), maize starch, sodium lauryl sulphate, gelatine, magnesium stearate (E572).
Tablet coating: Hypromellose 2910, Talc, Macrogol 6000, Titanium dioxide (E171).

What Atenolol looks like and contents of the pack

Film-coated tablet.

Atenolol 25 mg film-coated tablets:

White round, biconvex, film coated tablets debossed with 'A25' on one side and having Breakline on other side. [Size: about 6.6mm]

Atenolol 50 mg film-coated tablets:

White round, biconvex, film coated tablets debossed with 'A50' on one side and having Breakline on other side. [Size: about 8.6mm]

Atenolol 100 mg film-coated tablets:

White round, biconvex, film coated tablets debossed with 'A100' on one side and having Breakline on other side. [Size: about 10.5mm]

Atenolol film-coated tablets are available in Clear PVC/PVdC- Aluminium blister packs.

Pack sizes:

25 mg, 50 mg & 100 mg: 28 film-coated tablets

Marketing Authorisation Holder

Milpharm Limited
Ares Block, Odyssey Business Park
West End Road
Ruislip HA4 6QD
United Kingdom

Manufacturer

APL Swift Services (Malta) Limited
HF26, Hal Far Industrial Estate, Hal Far
Birzebbugia, BBG 3000
Malta

or

Milpharm Limited
Ares Block, Odyssey Business Park
West End Road
Ruislip, HA4 6QD
United Kingdom

This leaflet was last revised in 12/2021.