

Package leaflet: Information for the user

Salofalk® 1g/Actuation Rectal Foam mesalazine

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4

What is in this leaflet

1. What Salofalk rectal foam is and what it is used for
2. What you need to know before you use Salofalk rectal foam
3. How to use Salofalk rectal foam
4. Possible side effects
5. How to store Salofalk rectal foam
6. Contents of the pack and other information

1. What Salofalk rectal foam is and what it is used for

Salofalk rectal foam contains the active substance mesalazine, an anti-inflammatory agent used to treat inflammatory bowel disease.

Salofalk rectal foam is used for the treatment of:

- Inflammation of the large intestine (colon) and rectum (back passage) known by doctors as ulcerative colitis.

2. What you need to know before you use Salofalk rectal foam

Do not use Salofalk rectal foam

- If you are or have been told you are **allergic** (hypersensitive) to salicylic acid, to salicylates such as Aspirin® or to any of the other ingredients of Salofalk rectal foam (listed in section 6)
- If you have a serious liver or kidney disease.

Warnings and precautions

Before you start using this medicine you should tell your doctor

- If you have a history of problems with your lungs, particularly if you suffer from **bronchial asthma**.
- If you have a history of **allergy to sulphasalazine**, a substance related to mesalazine.
- If you suffer with problems of your **liver**.
- If you suffer with problems of your **kidney**.
- If you have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after using mesalazine.

Further precautions:

During treatment your doctor may want to keep you under close medical supervision, and you may need to have regular blood and urine tests.

Kidney stones may develop with use of mesalazine. Symptoms may include pain in sides of abdomen and blood in urine. Take care to drink sufficient amount of liquid during treatment with mesalazine.

Serious skin reactions including Stevens-Johnson syndrome, toxic epidermal necrolysis have been reported in association with mesalazine treatment. Stop using mesalazine and seek medical attention immediately if you notice any of the symptoms related to these serious skin reactions described in section 4.

Other medicines and Salofalk rectal foam

Please tell your doctor if you take or use any of the medicines mentioned below, as the effects of these medicines may change (interactions):

- Certain agents that inhibit blood clotting (**medicines for thrombosis or to thin your blood, e.g. warfarin**).
- **Azathioprine, 6-mercaptopurine or thioguanine** (medicines used to treat immune disorders).

Tell your doctor or pharmacist if you are taking or have recently taken or might take any other medicines, including medicines obtained without a prescription. It may still be all right for you to use Salofalk rectal foam and your doctor will be able to decide what is suitable for you.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

You should only use Salofalk rectal foam during pregnancy if your doctor tells you to.

You should also only use Salofalk rectal foam during breast-feeding if your doctor tells you to, as this medicine may pass into breast milk.

Driving and using machines

There are no effects on the ability to drive and use machines.

Salofalk rectal foam contains propylene glycol, sodium metabisulphite and cetostearyl alcohol

This medicine contains 3.44 g propylene glycol in each actuation of Salofalk rectal foam, which may cause skin irritation.

Sodium metabisulphite may rarely cause severe hypersensitivity reactions and bronchospasm (can be experienced in the form of breathing problems).

Cetostearyl alcohol may cause local skin reactions (e.g. contact dermatitis).

3. How to use Salofalk rectal foam

Always use Salofalk rectal foam exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Method of administration:

This medicine may only be used rectally, so it has to be inserted through the anus. It is not intended to be taken by mouth. Do not swallow.

Dosage:

Dose for adults:

The usual dose is 2 spray actuations once a day at bedtime. If you have difficulty retaining this amount of foam, it may also be administered in two separate doses: one at bedtime and the other during the night or early in the morning (after excreting the first dose).


Emptying your bowels before using Salofalk rectal foam produces the best results.

Use in children:


Salofalk rectal foam should not be used in children because there is little experience and only limited documentation for an effect in children.

Preparing to use the foam


Store and use Salofalk rectal foam at room temperature (between 20 and 30°C). Do not refrigerate or freeze. Also see section 5.



Push the applicator firmly onto the spout of the spray can.
Shake the spray can for about 20 seconds to mix the contents.




Before first use, remove the safety lock (plastic flap) from under the pump dome.




Twist the dome on the top of the spray can until the semi-circular gap underneath is in line with the applicator. The spray can is now ready for use.


Using the foam



Place your index finger on the top of pump dome and turn the can upside down. Note that the spray can will only work properly when held with the pump dome pointing down.



Stand with one foot on a stool or chair or lie on your side with the lower leg stretched out and the upper leg bent for balance.
Insert the applicator into your rectum as far as possible. Push the pump dome down fully once and then **slowly** release it – the foam comes out of the spray can when you release the dome. To administer the second spray, push the dome again and release slowly. Keep the applicator in place for 10-15 seconds before taking it out. This ensures the full dose is delivered into the rectum and no foam is spilt.



After administering the foam, remove the applicator and dispose of it with your household waste in the plastic bag provided. Use a new applicator for the next administration.

- Wash your hands and try not to empty your bowels until the next morning.
- If you go to hospital or see another doctor, tell them you are using this medicine.

Use Salofalk rectal foam regularly and consistently to achieve the desired effect.

Duration of treatment

How long you will use the medicine depends upon your condition. Your doctor will decide how long you are to continue the medication.

Mild acute episodes of inflammatory bowel disease (ulcerative colitis) generally subside after 4-6 weeks. If long-term treatment is required, your doctor will prescribe you an oral form of mesalazine, e.g. Salofalk granules.

If you think that the effect of Salofalk rectal foam is too strong or too weak, talk to your doctor.

If you use more Salofalk rectal foam than you should

Contact a doctor if you are in doubt so he or she can decide what to do.

If you use too much Salofalk rectal foam on one occasion, just take your next dose as prescribed. Do not use a smaller amount.

If you forget to use Salofalk rectal foam

Do not take a larger than normal dose of Salofalk rectal foam next time, but continue treatment at the prescribed dosage.

If you stop using Salofalk rectal foam

Do not stop using this product until you have talked to your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, Salofalk rectal foam can cause side effects, although not everybody gets them.

All medicines can cause allergic reactions although serious allergic reactions are very rare. If you get any of the following symptoms after using this medicine, you should contact your doctor immediately:

- **Allergic skin rash**
- **Fever**
- **Breathing difficulties.**

If you experience a marked reduction of your general health, especially, if accompanied by fever, and/or a sore throat and mouth, stop using this foam and contact your doctor immediately.

These symptoms can, very rarely, be due to a fall in the number of white cells in your blood (a condition called agranulocytosis), which may make you more prone to developing a serious infection. A blood test can confirm whether your symptoms are due to an effect of this medicine on your blood.

Stop using mesalazine and seek medical attention immediately if you notice any of the following symptoms:

- reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by fever and flu-like symptoms.

The following side effects have also been reported by patients using mesalazine:

Common: may affect less than 1 in 10 patients

- Abdominal discomfort.

Uncommon: may affect less than 1 in 100 patients

- Anal discomfort, anal irritation and painful urgent need to empty the bowels.

Rare: may affect less than 1 in 1,000 patients

- Abdominal pain, diarrhoea, wind (flatulence), nausea and vomiting
- Headache, dizziness
- Chest pain, breathlessness or swollen limbs because of an effect on your heart
- Increased sensitivity of your skin to sun and ultraviolet light (photosensitivity)

Very rare: may affect less than 1 in 10,000 patients

- Changes in kidney function, sometimes with swollen limbs or flank pain
- Severe abdominal pain, because of acute inflammation of the pancreas
- Shortness of breath, cough, wheezing, lung shadow on x-ray due to allergic and/or inflammatory lung conditions
- Severe diarrhoea and abdominal pain, because of an allergic reaction to this medicine within the bowel
- Skin rash or inflammation
- Muscle and joint pain
- Fever, sore throat or malaise, because of blood count changes
- Jaundice or abdominal pain because of liver and bile flow disorders
- Hair loss and the development of baldness
- Numbness and tingling in the hands and feet (peripheral neuropathy)
- Reversible decrease in semen production

Not known: frequency cannot be estimated from the available data

- Kidney stones and associated kidney pain (see also section 2)

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system:

Ireland

HPRA Pharmacovigilance

Website: www.hpra.ie

United Kingdom

Yellow Card Scheme

Website: www.mhra.gov.uk/yellowcard
or search for MHRA Yellow Card in the
Google Play or Apple App Store

By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Salofalk rectal foam

Keep this medicine out of the sight and reach of children.

Do not use Salofalk rectal foam after the expiry date which is stated on the carton and the spray can. The expiry date refers to the last day of that month.

The contents of the container must be used within 12 weeks after first opening.

Do not store above 30°C.

Do not refrigerate or freeze.

Note: foam may cause staining.

The container is pressurised and contains 3.75% by weight of flammable propellant. Protect from sunlight and temperatures over 50°C. Do not force open, pierce or burn empty containers, even after use. Do not spray near a flame or incandescent material.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Salofalk rectal foam contains

The active substance of Salofalk rectal foam is mesalazine and each spray actuation contains 1 g of mesalazine.

The other ingredients are sodium metabisulphite (E223), disodium edetate, cetostearyl alcohol, polysorbate 60, propylene glycol and propane, n-butane, isobutane as propellants.

What Salofalk rectal foam looks like and contents of the pack

Salofalk rectal foam is a white-greyish to slightly reddish-violet and creamy firm foam.

Salofalk rectal foam is available in packs containing 1 spray can and 14 applicators and as a bundle pack. A bundle pack consists of 4 packs containing 1 spray can and 14 applicators each. Each spray can of Salofalk rectal foam contains 80 grams of foam, which is sufficient for 14 spray actuations (equivalent to 7 doses).

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Dr. Falk Pharma GmbH
Leinenweberstr. 5
79108 Freiburg
Germany
Tel.: +49 (0) 761 / 1514-0
E-mail: zentrale@drfalkpharma.de

Salofalk rectal foam is authorised in the Member States of the EEA under the following name: Austria, Denmark, Finland, Germany, Great Britain, Greece, Ireland, Luxembourg, The Netherlands, Norway, Portugal, Spain and Sweden: Salofalk®

This leaflet was last revised in 01/2021.

Marketing authorisation numbers:

PL 08637/0003

PA573/4/5