



Package leaflet: Information for the user

**Etoricoxib 30 mg film-coated tablets**  
**Etoricoxib 60 mg film-coated tablets**  
**Etoricoxib 90 mg film-coated tablets**  
**Etoricoxib 120 mg film-coated tablets**

Etoricoxib

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Etoricoxib Tablets is and what it is used for
2. What you need to know before you take Etoricoxib Tablets
3. How to take Etoricoxib Tablets
4. Possible side effects
5. How to store Etoricoxib Tablets
6. Contents of the pack and other information

**1. What Etoricoxib Tablets is and what it is used for**

What is Etoricoxib Tablets?

- Etoricoxib Tablets contains the active substance etoricoxib. Etoricoxib is one of a group of medicines called selective COX-2 inhibitors. These belong to a family of medicines called non-steroidal anti-inflammatory drugs (NSAIDs).

**What is Etoricoxib Tablets used for?**

- Etoricoxib Tablets helps to reduce the pain and swelling (inflammation) in the joints and muscles of people 16 years of age and older with osteoarthritis, rheumatoid arthritis, ankylosing spondylitis and gout.
- Etoricoxib Tablets is also used for the short term treatment of moderate pain after dental surgery in people 16 years of age and older.

**What is osteoarthritis?**

Osteoarthritis is a disease of the joints. It results from the gradual breakdown of cartilage that cushions the ends of the bones. This causes swelling (inflammation), pain, tenderness, stiffness and disability.

**What is rheumatoid arthritis?**

Rheumatoid arthritis is a long term inflammatory disease of the joints. It causes pain, stiffness, swelling, and increasing loss of movement in the joints it affects. It may also cause inflammation in other areas of the body.

**What is gout?**

Gout is a disease of sudden, recurring attacks of very painful inflammation and redness in the joints. It is caused by deposits of mineral crystals in the joint.

**What is ankylosing spondylitis?**

Ankylosing spondylitis is an inflammatory disease of the spine and large joints.

**2. What you need to know before you take Etoricoxib Tablets**

**Do not take** Etoricoxib Tablets :

- if you are allergic (hypersensitive) to etoricoxib or any of the other ingredients of this medicine (listed in section 6)
- if you are allergic to non-steroidal anti-inflammatory drugs (NSAIDs), including aspirin and COX-2 inhibitors (see Possible Side Effects, section 4)
- if you have a current stomach ulcer or bleeding in your stomach or intestines
- if you have serious liver disease
- if you have serious kidney disease
- if you are or could be pregnant or are breast-feeding (see 'Pregnancy, breast feeding and fertility')
- if you are under 16 years of age
- if you have inflammatory bowel disease, such as Crohn's Disease, Ulcerative Colitis, or Colitis
- if you have high blood pressure that has not been controlled by treatment (check with your doctor or nurse if you are not sure whether your blood pressure is adequately controlled)
- if your doctor has diagnosed heart problems including heart failure (moderate or severe types), angina (chest pain)
- if you have had a heart attack, bypass surgery, peripheral arterial disease (poor circulation in legs or feet due to narrow or blocked arteries)
- if you have had any kind of stroke (including mini-stroke, transient ischaemic attack or TIA). Etoricoxib Tablets may slightly increase your risk of heart attack and stroke and this is why it should not be used in those who have already had heart problems or stroke.

If you think any of these are relevant to you, do not take the tablets until you have consulted your doctor.

**Warnings and precautions**

Talk to your doctor or pharmacist before taking Etoricoxib Tablets if:

- You have a history of stomach bleeding or ulcers.
- You are dehydrated, for example by a prolonged bout of vomiting or diarrhoea.
- You have swelling due to fluid retention.
- You have a history of heart failure, or any other form of heart disease.
- You have a history of high blood pressure. Etoricoxib Tablets can increase blood pressure in some people, especially in high doses, and your doctor will want to check your blood pressure from time to time.
- You have any history of liver or kidney disease.
- You are being treated for an infection. Etoricoxib Tablets can mask or hide a fever, which is a sign of infection.
- You have diabetes, high cholesterol, or are a smoker. These can increase your risk of heart disease.
- You are a woman trying to become pregnant.
- You are over 65 years of age.

If you are not sure if any of the above apply to you, **talk to your doctor before taking Etoricoxib Tablets** to see if this medicine is suitable for you.

Etoricoxib Tablets works equally well in older and younger adult patients. If you are over 65 years of age, your doctor will want to appropriately keep a check on you. No dosage adjustment is necessary for patients over 65 years of age.

**Children and adolescents**

Do not give this medicine to children and adolescents under 16 years of age.

**Other medicines and Etoricoxib Tablets**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

In particular if you are taking any of the following medicines, your doctor may want to monitor you to check that your medicines are working properly, once you start taking Etoricoxib Tablets :

- medicines that thin your blood (anticoagulants), such as warfarin
- rifampicin (an antibiotic)
- methotrexate (a drug used for suppressing the immune system, and often used in rheumatoid arthritis)
- ciclosporin or tacrolimus (drugs used for suppressing the immune system)
- lithium (a medicine used to treat some types of depression)
- medicines used to help control high blood pressure and heart failure called ACE inhibitors and angiotensin receptor blockers, examples include enalapril and ramipril, and losartan and valsartan
- diuretics (water tablets)
- digoxin (a medicine for heart failure and irregular heart rhythm)
- minoxidil (a drug used to treat high blood pressure)
- salbutamol tablets or oral solution (a medicine for asthma)
- birth control pills (the combination may increase your risk of side effects)
- hormone replacement therapy (the combination may increase your risk of side effects)
- aspirin, the risk of stomach ulcers is greater if you take Etoricoxib Tablets with aspirin.
  - aspirin for prevention of heart attacks or stroke: Etoricoxib Tablets can be taken with **low-dose** aspirin. If you are currently taking low-dose aspirin to prevent heart attacks or stroke, you should not stop taking aspirin until you talk to your doctor
  - aspirin and other non-steroidal anti-inflammatory drugs (NSAIDs): do not take **high dose** aspirin or other anti-inflammatory medicines while taking Etoricoxib Tablets .

**Etoricoxib Tablets with food and drink**

The onset of the effect of Etoricoxib Tablets may be faster when taken without food.

**Pregnancy, breast-feeding and fertility**

*Pregnancy*

Do not take etoricoxib if you are in the last 3 months of pregnancy as it could harm your unborn child or cause problems at delivery. It can cause kidney and heart problems in your unborn baby. It may affect your and your baby's tendency to bleed and cause labour to be later or longer than expected. You should not take Etoricoxib during the first 6 months of pregnancy unless absolutely necessary and advised by your doctor. If you need treatment during this period or while you are trying to get pregnant, the lowest dose for the shortest time possible should be used. If taken for more than a few days from 20 weeks of pregnancy onward, Etoricoxib can cause kidney problems in your unborn baby that may lead to low levels of amniotic fluid that surrounds the baby (oligohydramnios) or narrowing of a blood vessel (ductus arteriosus) in the heart of the baby. If you need treatment for longer than a few days, your doctor may recommend additional monitoring.

*Breast-feeding*

It is not known if etoricoxib is excreted in human milk. If you are breast-feeding, or planning to breast-feed, consult your doctor before taking Etoricoxib Tablets . If you are using Etoricoxib Tablets , you must not breast-feed.

*Fertility*

Etoricoxib Tablets is not recommended in women attempting to become pregnant.

**Driving and using machines**

Dizziness and sleepiness have been reported in some patients taking Etoricoxib Tablets . Do not drive if you experience dizziness or sleepiness.

Do not use any tools or machines if you experience dizziness or sleepiness.

**This medicine contains lactose**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

**Information on sodium content**

This medicine contains less than 1 mmol sodium (30 mg, 60 mg, 90 mg and 120 mg) per tablet is essentially 'sodium-free'.

**3. How to take Etoricoxib Tablets**

Always take this medicine exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Do not take more than the recommended dose for your condition. Your doctor will want to discuss your treatment from time to time. It is important that you use the lowest dose that controls your pain and you should not take Etoricoxib Tablets for longer than necessary. This is because the risk of heart attacks and strokes might increase after prolonged treatment, especially with high doses.

There are different strengths available for this medicinal product and depending on your disease your doctor will prescribe the tablet strength that is appropriate for you.



180 x 480 mm

Front Side

Font Name: Times New Roman

Font Size:

Main Heading - 12 pts (Bold)

Side Heading - 09 pts (Bold)

Body - 09 pts (Regular)

**Product Name: Etoricoxib 30, 60, 90 & 120 mg**

**Component: Pack insert**

**Dimension: 180 x 480 mm**

**Buyer/Country: SPUK**

**NOTE: Pack Insert text layout & dimension shall be changed during product commercialization time**

The recommended dose is:

**Osteoarthritis**

The recommended dose is 30 mg once a day, increase to a maximum of 60 mg once a day if needed.

**Rheumatoid arthritis**

The recommended dose is 60 mg once a day, increased to a maximum of 90 mg once a day if needed.

**Ankylosing spondylitis**

The recommended dose is 60 mg once a day, increased to a maximum of 90 mg once a day if needed.

**Acute pain conditions**

Etoricoxib should be used only for the acute painful period.

**Gout**

The recommended dose is 120 mg once a day which should only be used for the acute painful period, limited to a maximum of 8 days treatment.

**Postoperative dental surgery pain**

The recommended dose is 90 mg once daily, limited to a maximum of 3 days treatment.

**People with liver problems**

- If you have mild liver disease, you should not take more than 60 mg a day.
- If you have **moderate** liver disease, you should not take more than **30 mg a day**.

**Use in children and adolescents**

Etoricoxib tablets should not be taken by children or adolescents under 16 years of age.

**Elderly**

No dose adjustment is necessary for elderly patients. As with other medicines, caution should be exercised in elderly patients.

**Method of administration**

Etoricoxib Tablets is for oral use. Take the tablets once a day. Etoricoxib Tablets can be taken with or without food.

**If you take more Etoricoxib Tablets than you should**

You should never take more tablets than the doctor recommends. If you do take too many etoricoxib tablets, you should seek medical attention immediately.

**If you forget to take Etoricoxib Tablets**

It is important to take Etoricoxib Tablets as your doctor has prescribed. If you miss a dose, just resume your usual schedule the following day. Do not take a double dose to make up for the forgotten tablet.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**If you develop any of these signs you should stop Etoricoxib Tablets and talk to your doctor immediately (see What you need to know before you take Etoricoxib Tablets section 2):**

- shortness of breath, chest pains, or ankle swelling appear or if they get worse
- yellowing of the skin and eyes (jaundice) – these are signs of liver problems
- severe or continual stomach pain or your stools become black
- an allergic reaction- which can include skin problems such as ulcers or blistering, or swelling of the face, lips, tongue, or throat which may cause difficulty in breathing

The frequency of possible side effects listed below is defined using the following convention: Very common (affects more than 1 user in 10)

Common (affects 1 to 10 users in 100)

Uncommon (affects 1 to 10 users in 1,000)

Rare (affects 1 to 10 users in 10,000)

Very rare (affects less than 1 user in 10,000)

The following side effects can occur during treatment with Etoricoxib Tablets:

**Very common**

- stomach pain

**Common**

- dry socket (inflammation and pain after a tooth extraction)
- swelling of the legs and/or feet due to fluid retention (oedema)
- dizziness, headache
- palpitations (fast or irregular heartbeat), irregular heart rhythm (arrhythmia)
- increased blood pressure
- wheezing or shortness of breath (bronchospasms)
- constipation, wind (excessive gas), gastritis (inflammation of the lining of the stomach), heartburn, diarrhoea, indigestion (dyspepsia)/stomach discomfort, nausea, being sick (vomiting), inflammation of the oesophagus, mouth ulcers
- changes in blood tests related to your liver
- bruising
- weakness and fatigue, flu-like illness

**Uncommon**

- gastroenteritis (inflammation of the gastrointestinal tract that involves both the stomach and small intestine/ stomach flu), upper respiratory infection, urinary tract infection
- changes in laboratory values (decreased number of red blood cells, decreased number of white blood cells, platelets decreased)
- hypersensitivity (an allergic reaction including hives which may be serious enough to require immediate medical attention)
- appetite increases or decreases, weight gain
- anxiety, depression, decreases in mental sharpness; seeing, feeling or hearing things that are not there (hallucinations)
- taste alteration, inability to sleep, numbness or tingling, sleepiness
- blurred vision, eye irritation and redness
- ringing in the ears, vertigo (sensation of spinning while remaining still)

- abnormal heart rhythm (atrial fibrillation), fast heart rate, heart failure, feeling of tightness, pressure or heaviness in the chest (angina pectoris), heart attack
- flushing, stroke, mini-stroke (transient ischemic attack), severe increase in blood pressure, inflammation of the blood vessels
- cough, breathlessness, nose bleed
- stomach or bowel bloating, changes in your bowel habits, dry mouth, stomach ulcer, inflammation of the stomach lining that can become serious and may lead to bleeding, irritable bowel syndrome, inflammation of the pancreas
- swelling of the face, skin rash or itchy skin, redness of the skin
- muscle cramp/spasm, muscle pain/stiffness
- high levels of potassium in your blood, changes in blood or urine tests relating to your kidney, serious kidney problems
- chest pain

**Rare**

- angioedema (an allergic reaction with swelling of the face, lips, tongue and/or throat which may cause difficulty in breathing or swallowing, which may be serious enough to require immediate medical attention)/ anaphylactic/anaphylactoid reactions including shock (a serious allergic reaction that requires immediate medical attention)
- confusion, restlessness
- liver problems (hepatitis)
- low blood levels of sodium
- liver failure, yellowing of the skin and/or eyes (jaundice)
- severe skin reactions

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.'

**5. How to store Etoricoxib Tablets**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton. The expiry date refers to the last day of that month.

Blisters: Store in the original package in order to protect from moisture.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

**6. Contents of the pack and other information**

**What Etoricoxib Tablets contains**

- The active substance is etoricoxib. Each film coated tablet contains 30, 60, 90 or 120 mg of etoricoxib.
- The other ingredients are:

*Core* Cellulose Microcrystalline, Calcium Hydrogen Phosphate (Anhydrous), Croscarmellose sodium, Magnesium stearate (veg)

*Coating:* Hypromellose 2910, Lactose monohydrate, Titanium dioxide, Triacetin, Carnauba wax.

The 30mg, 60 mg and 120 mg tablets also contain Iron oxide yellow and Indigo carmine aluminium lake (FD&C Blue#2).

**What Etoricoxib Tablets looks like and contents of the pack**

Etoricoxib tablets are available in four strengths:

*30 mg tablets:* 'Blue-green', 'apple-shaped' biconvex film coated tablets debossed with ET on one side and 30 on the other side.

*60 mg tablets:* 'Dark-green', 'apple-shaped' biconvex film coated tablets debossed with ET on one side and 60 on the other side.

*90 mg tablets:* White, 'apple-shaped' biconvex film coated tablets debossed with ET on one side and 90 on the other side.

*120 mg tablets:* 'Pale-green', 'apple-shaped', biconvex film coated tablets debossed with ET on one side and 120 on the other side.

**30 mg, 60 mg, 90 mg:**

Pack sizes of 28, 98 tablets.

**120 mg:**

Pack sizes of 7, 28 tablets.

Not all pack sizes may be marketed.

**Marketing Authorisation Holder and Manufacturer**

Strides Pharma UK Ltd.  
Unit 4 The Metro Centre,  
Dwight Road, Watford,  
WD18 9SS  
United Kingdom

**This leaflet was last revised in Dec 2023.**



180 x 480 mm

Back Side

Font Name: Times New Roman

Font Size:

Main Heading - 12 pts (Bold)

Side Heading - 09 pts (Bold)

Body - 09 pts (Regular)

**Product Name:** Etoricoxib 30, 60, 90 & 120 mg

**Component:** Pack insert

**Dimension:** 180 x 480 mm

**Buyer/Country:** SPUK

**NOTE: Pack Insert text layout & dimension shall be changed during product commercialization time**