

## Package leaflet: Information for the user

### Ponstan 250 mg Capsules

mefenamic acid

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What Ponstan is and what it is used for
2. What you need to know before you take Ponstan
3. How to take Ponstan
4. Possible side effects
5. How to store Ponstan
6. Contents of the pack and other information

#### **1. What Ponstan is and what it is used for**

Ponstan capsules contain mefenamic acid which is a non-steroidal anti-inflammatory drug (NSAID).

They can help to relieve:

- symptoms of inflammation, such as redness and swelling
- pain and discomfort caused by arthritis, muscular or rheumatic disorders
- headache, muscle ache or toothache
- pain after operations, trauma
- childbirth pain
- fever in children
- painful or heavy periods.

#### **2. What you need to know before you take Ponstan**

**Do not take Ponstan:**

- if you are allergic to mefenamic acid, to any other anti-inflammatory medicines (such as aspirin, ibuprofen, celecoxib), or any of the other ingredients of this medicine (listed in section 6)

- if you have, or have ever had, stomach or intestinal conditions such as peptic ulcer, bleeding in the stomach or severe gastritis
- if you have an inflammatory bowel disease (e.g. ulcerative colitis, Crohn's disease)
- if you have severe heart, liver or kidney problems
- if you have just had heart bypass surgery
- if you are more than 6 months pregnant.

If any of the above apply to you, speak to your doctor or pharmacist.

### **Warnings and precautions**

Talk to your doctor or pharmacist before taking Ponstan:

- if you are taking any other NSAIDs (e.g. ibuprofen, diclofenac)
- if you are taking any other anti-inflammatory medicines including steroids (e.g. prednisolone)
- if you are taking aspirin or medicines that thin the blood (e.g. warfarin, clopidogrel)
- if you are taking antidepressants called selective serotonin re-uptake inhibitors (SSRIs) (e.g. paroxetine)
- if you have kidney or liver problems. Your doctor may check your kidney or liver function before and during treatment
- if you are elderly (see section 3)
- if you are trying to become pregnant (see section on Fertility)
- if you have stomach or digestive tract problems or if you ever had an upset stomach after taking pain killers such as aspirin. Bleeding in the stomach or gut can occur in patients taking Ponstan
- if you have a bleeding disorder or if you are going to have a major operation. Ponstan can affect the clotting of your blood. It can make you bleed more and for longer than usual
- if you have asthma, or a history of asthma, as this medicine may cause breathing difficulties
- if you have a connective tissue disorder, e.g. Systemic Lupus Erythematosus (SLE)
- if you have epilepsy
- if you are dehydrated (thirsty with dry skin, dark urine, dry mouth, headache)
- if you have heart problems, previous stroke or think that you might be at risk of these conditions (e.g. if you have high blood pressure, diabetes or high cholesterol or are a smoker). Additional monitoring may be carried out by your doctor.

Medicines such as Ponstan may be associated with a small increased risk of heart attack or stroke. Any risk is more likely with high doses and prolonged treatment. **Do not exceed** the recommended dose or duration of treatment.

### **Blood tests**

Your doctor may test your blood during treatment.

### **Other medicines and Ponstan**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Some medicines may be affected by Ponstan or they may affect how well Ponstan will work. Tell your doctor or pharmacist if you are taking:

- medicines that can increase the chance of getting ulcers or a bleed in the stomach or gut, such as:
  - o corticosteroids used to treat arthritis and inflammation
  - o medicines such as anti-platelet agents, used to thin the blood (e.g. warfarin, aspirin, clopidogrel)
  - o antidepressants called selective serotonin re-uptake inhibitors (SSRIs) (e.g. paroxetine)
  - o any other anti-inflammatory medicines (e.g. diclofenac, celecoxib)
- aspirin including low doses of aspirin used to prevent your blood from clotting in certain heart conditions
- medicines used for high blood pressure (e.g. atenolol, ramipril, valsartan)
- diuretics (water tablets) or heart medicines (e.g. digoxin, sotalol, diltiazem)
- some diabetic medicines (e.g. glipizide, glibenclamide)
- medicines which suppress the immune system (e.g. ciclosporin, tacrolimus, methotrexate)
- lithium, a medicine used to treat mood swings and some types of depression
- a medicine usually prescribed through hospitals, called mifepristone (taken within the last 12 days)
- quinolone antibiotics (antibiotics used to treat infections)
- aminoglycoside antibiotics, used under medical supervision in hospitals
- zidovudine, a medicine used for HIV
- probenecid, a medicine used in special cases, to protect the kidneys
- medicines which bind to protein in the blood - (check with your pharmacist).

### **Ponstan with alcohol**

Avoid drinking alcohol. It may increase your risk of stomach bleeding.

### **Pregnancy, breast-feeding and fertility**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

### **Pregnancy**

Mefenamic acid will be passed to your unborn baby. It is not known how much it will affect your unborn baby in the first 6 months of pregnancy.

DO NOT take the capsules in the last 3 months of pregnancy as they may delay the onset of labour and prolong its duration. They may also increase the likelihood of bleeding in the mother and in the baby.

If you need to take these capsules, your doctor can help you decide whether or not to take them during the first 6 months of pregnancy.

### **Breast-feeding**

Mefenamic acid passes into breast milk and can affect the baby. You should not take the capsules while breast-feeding unless advised by your doctor.

### **Fertility**

DO NOT take the capsules if you are trying to become pregnant, as they may make it more difficult to get pregnant. You should inform your doctor if you are planning to become pregnant or if you have problems becoming pregnant.

### **Driving and using machines**

Ponstan may cause drowsiness, dizziness, fatigue or affect your vision. If any of these occur do not drive, use machinery, or perform any tasks that may require you to be alert.

### **Ponstan Capsules contain lactose**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

## **3. How to take Ponstan**

Always take this medicine exactly as your doctor has told you. Your doctor will decide on the appropriate dose to suit your condition. Check with your doctor or pharmacist if you are not sure.

- Take the capsules with or immediately after a **meal**.
- Do NOT drink **alcohol** while taking Ponstan.

Alcohol and smoking can irritate the stomach and make some of the side effects worse.

## **Dosage**

### **Adults**

The recommended dose is 2 capsules three times a day.

### **Elderly patients (over 65 years)**

Elderly patients are at a higher risk of side effects and should take the lowest effective dose for the shortest possible time, with additional monitoring carried out by their doctor.

### **Use in children**

It is recommended that children under 12 years of age should be given Mefenamic Acid Suspension (50mg / 5ml).

### **If you take more Ponstan than you should**

If you take more capsules than you should you may harm your stomach, kidneys and you may get seizures (fits). Symptoms of overdose may also include headache, abdominal pain (pain in your stomach) or other abnormal stomach symptoms, nausea (feeling sick), vomiting, rarely diarrhoea, disorientation, excitation, coma (loss of consciousness for a period of time), tiredness, ringing in the ears and fainting.

1. Tell your doctor, pharmacist or nearest hospital casualty department immediately.
2. Take the container and any remaining capsules with you so that people can see what you have taken.
3. Do this even if you feel well.

### **If you forget to take Ponstan**

If you forget to take a dose take it as soon as you remember, but if it is almost time for your next dose, skip the missed dose and continue as usual.

Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them. Do not be alarmed by this list of possible side effects. You may not experience any of them.

**STOP taking** the capsules and **seek medical help immediately** if you have any of the following **allergic reactions**:

- difficulty breathing or swallowing, swelling of the face, lips, tongue or throat
- severe itching of the skin, with a red rash or raised lumps
- blistering of the mouth, eyes, and genital region, and patchy areas of rash, peeling skin

**or any of the following reactions**

- diarrhoea
- passing blood in your stools (faeces/motions)
- passing black tarry stools
- vomiting any blood or dark particles that look like coffee grounds.

**Seek immediate medical attention** if you have any of the following symptoms:

- indigestion or heartburn, abdominal pain (pain in your stomach) or other abnormal stomach symptoms, nausea (feeling sick), vomiting
- any unusual bruising or bleeding, for example nose-bleeds, pinpoint red spots on the skin, unusual purple bruise-like rash on the skin or in the mouth
- signs of anaemia such as feeling tired, breathless, and looking pale
- fever, sore throat, mouth ulcers, repeated infections or infections that will not go away. This may be due to a low level of white blood cells
- seizures (fits)
- signs of low sodium levels such as headache, nausea, vomiting, tiredness, muscle cramps
- sudden headache, stiff neck, fever, sensitivity to bright light, drowsiness and muscle pain, with or without a rash
- fever, rash, nausea, aches and pains, passing more or less urine than usual, passing red urine or passing urine at night. This may be due to changes in your kidneys
- sudden loss or blurring of vision, loss of colour vision, eye pain which worsens with eye movement
- headache, in particular on waking in the morning. This may be due to high blood pressure
- pain behind the ribs radiating towards the back, often worse when lying down, nausea, vomiting, fever. This may be due to inflammation of your pancreas
- yellowing of your skin or eyes, pale faeces and dark urine, unexplained persistent nausea, stomach problems, loss of appetite or unusual tiredness. This may be due to changes in your liver
- worsening of colitis and Crohn's disease
- elevation of one or more liver function tests

- failure of various organs at the same time (multi-organ failure)
- heart failure.

**The side effects listed below have been reported:**

**Not known: frequency cannot be estimated from the available data**

- head-spins (vertigo)
- fatty stools
- sweating
- rapid heartbeat (palpitations)
- mental confusion
- loss of appetite
- constipation or bloating
- blurred vision, eye irritation
- feeling ill (malaise)
- ringing or buzzing in the ears (tinnitus)
- numbness or tingling in hands or feet
- sudden poor blood sugar control if you have diabetes. Your doctor or pharmacist can measure your sugar levels
- asthma or asthma that is worse than usual
- swelling of your hands and feet (around the ankles)
- sore mouth (pain or ulcers on the tongue, cheeks, lips, throat or gums)
- dizziness, drowsiness, feeling lethargic and tired
- signs of low blood pressure such as light-headedness
- reactions to the sun. Your skin may become red, painful and swollen - do not sunbathe, use a sun bed, or expose your skin to artificial UV light
- depression
- inability to sleep
- hallucinations
- nervousness
- ear pain.

Medicines such as Ponstan may be associated with a small increased risk of heart attack or stroke. (See section 2 - end of 'Warnings and precautions').

**Urine tests**

Tell the doctor if you are having urine tests, as your medicine may affect the results.

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Ponstan**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister, carton/bottle/container after EXP. The expiry date refers to the last day of that month.

Do not store above 25°C. Store in the original package.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Ponstan contains**

- The active substance is mefenamic acid. Each capsule contains 250 mg mefenamic acid.
- The other ingredients are:
  - lactose monohydrate, sodium laurilsulfate, gelatin, purified water.
  - The capsule shell contains:  
gelatin, erythrosine (E127), quinoline yellow (E104), titanium dioxide (E171), patent blue V (E131).
  - The printing ink contains either shellac, black iron oxide (E172), propylene glycol (E1520), ammonium hydroxide (E527) **or** shellac, propylene glycol (E1520), strong ammonia solution, potassium hydroxide (E525), black iron oxide (E172).
  - (See end of section 2 for further information on lactose).

### **What Ponstan looks like and contents of the pack**

Ponstan Capsules are ivory opaque and powder blue opaque hard gelatin capsules marked 'PONSTAN 250'.

They are available in blister packs containing 6, 10, 12, 20, 30, 50, 100, or 168 capsules and in containers of 100 or 500 capsules.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder**

Chemidex Pharma Ltd  
Chemidex House  
Egham Business Village  
Crabtree Road



Egham  
Surrey TW20 8RB  
United Kingdom

**Manufacturer**

Lelypharma B.V.  
Zuiveringweg 42  
8243 PZ Lelystad  
the Netherlands

**This leaflet was last revised in March 2020.**

'Ponstan' and 'Chemidex' are trademarks.