

## **Package leaflet: Information for the user**

### **HBVAXPRO® 5 micrograms suspension for injection in pre-filled syringe** Hepatitis B vaccine (recombinant DNA)

**Read all of this leaflet carefully before you or your child is vaccinated because it contains important information.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- If you or your child get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

1. What HBVAXPRO 5 micrograms is and what it is used for
2. What you need to know before you or your child receive HBVAXPRO 5 micrograms
3. How HBVAXPRO 5 micrograms is given
4. Possible side effects
5. How to store HBVAXPRO 5 micrograms
6. Content of the pack and other information

#### **1. What HBVAXPRO 5 micrograms is and what it is used for**

This vaccine is indicated for active immunisation against hepatitis B virus infection caused by all known subtypes in individuals from birth through 15 years of age considered at risk of exposure to hepatitis B virus.

It can be expected that hepatitis D will also be prevented by immunisation with HBVAXPRO as hepatitis D does not occur in the absence of hepatitis B infection.

The vaccine will not prevent infection caused by other agents such as hepatitis A, hepatitis C and hepatitis E and other pathogens known to infect the liver.

#### **2. What you need to know before you or your child receive HBVAXPRO 5 micrograms**

##### **Do not use HBVAXPRO 5 micrograms**

- if you or your child is allergic to hepatitis B surface antigen or to any of the other ingredients of HBVAXPRO (see section 6)
- if you or your child has a severe illness with fever

##### **Warnings and precautions**

The container of this vaccine contains latex rubber. Latex rubber may cause severe allergic reactions.

Talk to your doctor, pharmacist or nurse before you or your child receive HBVAXPRO 5 micrograms.

##### **Other vaccines and HBVAXPRO 5 micrograms**

HBVAXPRO can be administered at the same time as with hepatitis B immunoglobulin, at a separate injection site.

HBVAXPRO can be used to complete a primary immunisation course or as a booster dose in subjects who have previously received another hepatitis B vaccine.

HBVAXPRO may be administered at the same time as with some other vaccines, using separate sites and syringes.

**Tell your doctor, pharmacist or nurse if you or your child is taking or has recently taken any other medicines, including medicines obtained without a prescription.**

### **Pregnancy and breast-feeding**

Caution should be exercised when prescribing the vaccine to pregnant or breast-feeding women. Ask your doctor, pharmacist or nurse for advice before taking any medicine.

### **Driving and using machines**

HBVAXPRO is expected to have no or negligible influence on the ability to drive and use machines.

**HBVAXPRO 5 micrograms contains sodium:** This medicinal product contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

## **3. How HBVAXPRO 5 micrograms is given**

### **Dosage**

The recommended dose for each injection (0.5 mL) is 5 micrograms for individuals from birth through 15 years of age.

A course of vaccination should include at least three injections.

Two immunisation schedules can be recommended:

- two injections with an interval of one month followed by a third injection 6 months after the first administration (0, 1, 6 months).
- if immunity is needed quickly: three injections with an interval of one month and a fourth dose 1 year later (0, 1, 2, 12 months).

In case of a recent exposure to the hepatitis B virus, a first dose of HBVAXPRO together with the appropriate dose of immunoglobulin can be given.

Some local vaccination schedules currently include recommendations for a booster dose. Your doctor, pharmacist or nurse will inform you if a booster dose should be given.

### **Method of administration**

The doctor or nurse will give the vaccine as an injection into muscle. The upper side of the thigh is the preferred site for injection in neonates and infants. The upper arm muscle is the preferred site for injection in children and adolescents.

This vaccine should never be given into a blood vessel.

Exceptionally, the vaccine may be administered subcutaneously in patients with thrombocytopaenia (diminution of blood platelets) or to persons at risk of haemorrhage.

### **If you or your child forget one dose of HBVAXPRO 5 micrograms**

If you or your child miss a scheduled injection, talk to your doctor, pharmacist or nurse. Your doctor or nurse will decide when to give the missed dose.

If you or your child have any further questions on the use of this product, ask your doctor pharmacist or nurse.

## **4. Possible side effects**

Like all medicines, this vaccine can cause side effects, although not everybody gets them.

As with other hepatitis B vaccines, in many instances, the causal relationship of side effects to the vaccine has not been established.

The most common side effects seen are injection-site reactions: soreness, redness and hardening.

Other side effects are reported very rarely:

- Low platelet count, lymph node disease
- Allergic reactions
- Nervous system disorders such as pins and needles, facial paralysis, nerve inflammations including Guillain-Barre Syndrome, inflammation of the nerve of the eye that leads to impaired vision, brain inflammation, exacerbation of multiple sclerosis, multiple sclerosis, convulsions, headache, dizziness and fainting
- Low blood pressure, blood vessel inflammation
- Asthma-like symptoms
- Vomiting, nausea, diarrhoea, abdominal pain
- Skin reactions such as eczema, rash, itching, hives and skin blistering, hair loss
- Joint pain, arthritis, muscle pain, pain in extremity
- Fatigue, fever, vague illness, flu-like symptoms
- Elevation of liver enzymes
- Inflammation of the eye which causes pain and redness

In babies born very prematurely (at or before 28 weeks of gestation) longer gaps than normal between breaths may occur for 2-3 days after vaccination.

## Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store HBVAXPRO 5 micrograms

Keep this vaccine out of the sight and reach of children.

Do not use this vaccine after the expiry date which is stated on the label.

Store in a refrigerator (2°C - 8°C).

Do not freeze.

Store in the original package in order to protect from light.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What HBVAXPRO 5 micrograms contains

The active substance is:

Hepatitis B virus surface antigen, recombinant (HBsAg) \* ..... 5 micrograms  
Adsorbed on amorphous aluminium hydroxyphosphate sulfate (0.25 milligram Al<sup>3+</sup>)<sup>#</sup>

\* produced in *Saccharomyces cerevisiae* (strain 2150-2-3) yeast by recombinant DNA technology.

# Amorphous aluminium hydroxyphosphate sulfate is included in this vaccine as an adsorbant. Adsorbants are substances included in certain vaccines to accelerate improve and/or prolong the protective effects of the vaccine.

The other ingredients are sodium chloride (NaCl), borax and water for injections.

### **What HBVAXPRO 5 micrograms looks like and contents of the pack**

HBVAXPRO 5 micrograms is a suspension for injection in a syringe.

Pack sizes of 1, 10, 20 and 50 pre-filled syringes without needle or with 2 separate needles,

Pack sizes of 1 and 10 pre-filled syringes with 1 separate needle.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

**Marketing Authorisation Holder:** Merck Sharp & Dohme (UK) Limited, 120 Moorgate, London, EC2M 6UR, United Kingdom.

**Manufacturer:** Merck Sharp and Dohme B.V., Waarderweg 39, 2031 BN Haarlem, The Netherlands.

For any information about this vaccine, please contact:

Merck Sharp & Dohme (UK) Limited

Tel: +44 (0) 208 154 8000

Email: [medicalinformationuk@msd.com](mailto:medicalinformationuk@msd.com)

**This leaflet was last revised in October 2023.**

**The following information is intended for medical or health care professionals only:**

#### **Instructions**

The vaccine should be inspected visually prior to administration for any foreign particulate matter and/or abnormal physical appearance. The syringe should be well shaken until a slightly opaque white suspension is obtained.

The needle is attached by twisting in clockwise direction, until the needle fits securely on the syringe.

© 2023 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.

PIL.HBV.5MCG.PFS.23.GB.8486.Art61-3.RCN024417.RCN021777