

**Package leaflet: Information for the user**

## Magnaspartate® 243 mg powder for oral solution magnesium

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.
- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse.

**What is in this leaflet**

1. What MAGNASPARTATE is and what it is used for
2. What you need to know before you take MAGNASPARTATE
3. How to take MAGNASPARTATE
4. Possible side effects
5. How to store MAGNASPARTATE
6. Contents of the pack and other information

**1. What MAGNASPARTATE is and what it is used for**

Magnaspartate is a mineral preparation and contains the active ingredient magnesium aspartate dihydrate equivalent to 243 mg (10 mmol) of magnesium. Magnaspartate is used in the treatment and prevention of magnesium deficiency as diagnosed by a doctor.

You must talk to a doctor if you do not feel better or if you feel worse after taking this medicine.

**2. What you need to know before you take MAGNASPARTATE**

Do not take MAGNASPARTATE:

- if you are allergic to magnesium aspartate dihydrate or any of the other ingredients of this medicine (listed in section 6).
- if you have severe kidney problems
- if you have a very slow heart rhythm (called 'bradycardia')

**Other medicines and MAGNASPARTATE**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This is because Magnaspartate can affect the way some other medicines work. Also, some other medicines can affect the way Magnaspartate works.

In particular, you must allow a time interval of 2-3 hours between the taking MAGNASPARTATE and the following medicines.

- **Cellulose sodium phosphate**, used to prevent kidney stones
- **EDTA**, used in hospital for high calcium levels
- **Flourides**, used for preventing tooth decay
- **Tetracyclines, Nitrofurantoin, Nitroxoline**, antibiotic to treat infections
- **Aminoquinolines**, medicines to treat malaria
- **Quinidine and quinidine derivatives**, medicines to treat heart rhythm disorders
- **Penicillamine**, medicine used to treat severe rheumatoid arthritis and used in Wilson's disease
- **Iron**, to treat anemia
- **Bisphosphonates**, to treat osteoporosis
- **Eltrombopag**, to correct bleeding deficiency

The following medicines may cause magnesium deficiency. Your doctor may decide to adjust your daily dose of magnesium.

- **Aminoglycosides**, antibiotics used to treat infections e.g. gentamicin, streptomycin and neomycin
- **Certain medicines used to treat cancer** such as cisplatin and EGF-receptor antagonists such as cetuximab and erlotinib
- **Ciclosporins**, medicines used to suppress the body's immune system e.g. following an organ transplant
- **Water tablets (Diuretics)**, medicines used to treat blood pressure or heart failure e.g. thiazide, furosemide, bumetanide
- **Proton Pump Inhibitors**, medicines used to treat high acid in the stomach or oesophagus. e.g. omeprazole, pantoprazole
- **Amphotericin B**, antibiotic that fights fungal infections in the body
- **Rapamycin**, medicine used to suppress the body's immune system e.g. following an organ transplant
- **Pentamidine**, medicine to treat infections such as pneumonia
- **Foscarnet**, an antiviral medication

**MAGNASPARTATE with drinks**

Magnaspartate can be taken with water, tea or orange juice.

**Pregnancy, breast-feeding and fertility**

Magnaspartate can be used during pregnancy and during breast-feeding without concerns. Based on long-term experience, no effects of magnesium on male and female fertility are anticipated.

**Driving and using machines**

Magnaspartate has no or negligible influence on the ability to drive and use machines.


**MAGNASPARTATE contains sucrose and sodium.**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine. Contains 2.706 g of sucrose per sachet. This should be taken into account in patients with diabetes mellitus.

In case of frequent/and or permanent use, may be harmful to the teeth. This medicine contains less than 1 mmol sodium (23 mg) per sachet, that is to say essentially 'sodium-free'.

**3. How to take MAGNASPARTATE**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

<b>FOCUS DESIGN</b>	<b>Product:</b> Magnaspartate 243 mg powder for oral solution	<b>Origination:</b> 16/03/20 Update: 20/03/20 Update: 22/04/20 Update: 23/06/20 Update: 29/06/20 Update: 07/07/20	<b>Pharma Code:</b> 1226	<b>Registration:</b>
	<b>Spec Code:</b> 1606837B			
	<b>Dimensions:</b> 170x295mm			<b>Date:</b>
<b>Client:</b> <b>KORA</b> HEALTHCARE	<b>Typofont:</b> Arial 9 point			

12 mm

**The recommended dose is**  
**Adults (over 18 years)** 1-2 sachets daily (equal to 243-486 mg magnesium).

**Use in children and adolescents**  
**Children and Adolescents (10 to 18 years)** 1 sachet daily (equal to 243 mg magnesium)  
**Children (4 to 10 years)** 1 level 5 ml spoon daily (109 mg magnesium) or 1 sachet daily (243 mg magnesium)  
**Children (2 to 4 years)** 1 level 5 ml spoon daily (109 mg magnesium).

**Renal patients:**  
 No dose adjustment is required in mild to moderate renal failure.

**Directions for use:**  
 Dissolve one sachet or one level 5 ml spoon of Magnaspartate in 50-200ml water, tea or orange juice. Stir until the solution in water is cloudy to transparent. In orange juice or tea inactive particles will be visible.  
 Drink immediately or within 24 hours if made with bottled water and stored below 25°C. Discard any remaining content of the sachet.  
 If you are unable to swallow and have a gastric, duodenal or nasal feeding tube, Magnaspartate in 200ml water can be administered through the tube immediately after preparation or within 24 hours if made with bottled water and stored below 25°C.

**If you take more MAGNASPARTATE than you should**  
 No side effects are expected, if your kidneys work properly. Any excessive amount of magnesium will be excreted through your kidneys.  
 If you have taken more of this medicine than directed, or if a person with severe kidney problems or if a child accidentally has taken this medicine, please contact your doctor or local hospital casualty department for judgement of the risk and advice.

**If you forget to take MAGNASPARTATE**  
 Do not take a double dose to make up for a forgotten dose.  
 If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. Possible side effects**  
 Like all medicines, this medicine can cause side effects, although not everybody gets them.  
**Uncommon** (may affect 1 to 10 users in 1,000):  
 Soft stools to diarrhoea at the beginning of treatment (harmless and will usually decrease in frequency as the treatment advances).  
**Very rare** (may affect less than 1 user in 10,000):  
 Fatigue if Magnaspartate is used for a longer period of time.  
 If you get any side effects such as diarrhoea, you should temporarily interrupt the treatment. After the symptoms improve you can restart treatment, with a reduced dosage.

**Reporting of side effects**  
 If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via (see details below).

**IRELAND:**  
 HPRA Pharmacovigilance Section  
 Earlsfort Terrace, IRL - Dublin 2,  
 Tel: +353 1 6764971, Fax: +353 1 6762517  
 Website: www.hpra.ie,  
 e-mail:

**UK**  
 Yellow Card Scheme  
 Website: www.mhra.gov.uk/yellowcard  
 By reporting side effects, you can help provide more information on the safety of this medicine.

**5. How to store MAGNASPARTATE**  
 Do not store above 25°C.  
 When Magnaspartate has been dissolved in water, orange juice or tea, the solution should be consumed immediately or within 24 hours when dissolved in bottled water and stored below 25°C.  
 Keep this medicine out of the sight and reach of children.  
 Do not use this medicine after the expiry date which is stated on the sachet and carton after EXP. The expiry date refers to the last day of that month.  
 Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

**6. Contents of the pack and other information**  
**What MAGNASPARTATE contains**

- The active substance is magnesium. One sachet (6.5 g) contains magnesium aspartate dihydrate 3246 mg; magnesium content: 243 mg.
- The other ingredients are sucrose, citric acid monohydrate, peach/apricot flavour, saccharin sodium, silica, colloidal anhydrous

**What MAGNASPARTATE looks like and contents of the pack**  
 White powder with a peach/apricot-like flavour.  
 Magnaspartate is available in packs of 10 or 20 sachets. Not all pack sizes may be marketed.

**Marketing Authorisation Holder**  
 Kora Corporation Ltd t/a Kora Healthcare  
 Swords Business Park, Swords, Co. Dublin, Ireland

**Manufacturer**  
 Lusomedicamenta Sociedade Tecnica Farmaceutica, S.A.  
 Estrada Consiglieri Pedroso, 69 - B, Queluz, Barcarena, 2730-055, Portugal

This medicinal product is authorised in the Member States of the EEA under the following names:  
 Ireland Magnaspartate 243 mg powder for oral solution  
 UK Magnaspartate 243 mg powder for oral solution

**This leaflet was last revised in July 2020**

12 mm

57 mm

2 mm

2 mm

12 mm

12 mm

 **KORA**  
HEALTHCARE  
1606837B