



7001442

## Package leaflet: Information for the user

# ALDURAZYME® 100 U/ml concentrate for solution for infusion

Laronidase

**Is this leaflet hard to see or read? Phone 0800 035 2525 for help.**

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or your pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet:

1. What Aldurazyme is and what it is used for
2. What you need to know before you are given Aldurazyme
3. How Aldurazyme is given
4. Possible side effects
5. How to store Aldurazyme
6. Contents of the pack and other information

### 1. What Aldurazyme is and what it is used for

Aldurazyme is used to treat patients with MPS I disease (Mucopolysaccharidosis I). It is given to treat the non-neurological manifestations of the disease.

People with MPS I disease have either a low level or no level of an enzyme called  $\alpha$ -L-iduronidase, which breaks down specific substances (glycosaminoglycans) in the body. As a result, these substances do not get broken down and processed by the body as they should. They accumulate in many tissues in the body, which causes the symptoms of MPS I.

Aldurazyme is an artificial enzyme called laronidase. This can replace the natural enzyme which is lacking in MPS I disease.

### 2. What you need to know before you are given Aldurazyme

#### You should not be given Aldurazyme

If you are allergic (hypersensitive) to laronidase or any of the other ingredients of this medicine (listed in section 6).

#### Warnings and precautions

Talk to your doctor before using Aldurazyme.

Contact your doctor immediately if treatment with Aldurazyme causes:

- Allergic reactions, including anaphylaxis (a severe allergy reaction) – see under section 4 “Possible side effects”. Some of these reactions may be life-threatening. Symptoms may include respiratory failure/distress (inability of lungs to work properly), stridor (high-pitched breathing sound) and other disorders due to obstruction of airways, rapid breathing, excessive contraction of the airway muscles causing breathing difficulty (bronchospasm), lack of oxygen in body tissues (hypoxia), low blood pressure, slow heart rate, or itchy rash (urticaria).
- Infusion-associated reactions, i.e., any side effect occurring during the infusion or until the end of the infusion day - see under section 4 “Possible Side Effects” below for symptoms.

If these reactions occur, the Aldurazyme infusion should be stopped immediately and appropriate treatment will be started by your doctor.

These reactions may be particularly severe if you have a pre-existing MPS I-related upper airway obstruction.

You may be given additional medications to help prevent allergic-type reactions, such as antihistamines, medicine to reduce fever (e.g. paracetamol) and/or corticosteroids.

Your doctor will also decide if you can continue receiving Aldurazyme.

#### Other medicines and Aldurazyme

Inform your doctor if you are using medicines containing chloroquine or procaine, due to a possible risk of decreasing the action of Aldurazyme.

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

### Pregnancy, breast-feeding and fertility

There is not enough experience of the use of Aldurazyme in pregnant women. You should not be given Aldurazyme during pregnancy unless clearly necessary.

It is not known whether Aldurazyme appears in breast milk. It is recommended to stop breast-feeding during treatment with Aldurazyme.

No information is available on the effects of Aldurazyme on fertility.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

### Driving and using machines

The effects on the ability to drive and to use machines have not been studied.

### Aldurazyme contains sodium

This medicine contains 30 mg sodium (main component of cooking/table salt) per vial. This is equivalent to 1.5% of the recommended maximum daily dietary intake of sodium for an adult.

### 3. How Aldurazyme is given

#### Instruction for use - dilution and administration

The concentrate for solution for infusion has to be diluted before administration and is for intravenous use (see information for health care professionals).

Administration of Aldurazyme should be carried out in an appropriate clinical setting where resuscitation equipment to manage medical emergencies would be readily available.

#### Home infusion

Your doctor may consider that you can have home infusion of Aldurazyme if it is safe and convenient to do so. If you get a side effect during an infusion of Aldurazyme, your infusion staff member may stop the infusion and start appropriate medical treatment.

#### Dosage

The recommended dosage regimen of Aldurazyme is 100 U/kg body weight given once every week as an intravenous infusion. The initial infusion rate of 2 U/kg/h may be gradually increased every fifteen minutes, if tolerated, to a maximum of 43 U/kg/h. The total volume of the administration should be delivered in approximately 3-4 hours.

Always use this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

#### If you miss an infusion of Aldurazyme

If you have missed an Aldurazyme infusion, please contact your doctor.

#### If you are given more Aldurazyme than needed

If the dose of Aldurazyme given is too high or the infusion is too fast, adverse drug reactions may occur. Receiving an excessively fast infusion of Aldurazyme may cause nausea, abdominal pain, headache, dizziness and difficulty breathing (dyspnoea). In such situations, the infusion should be stopped or the infusion rate slowed down immediately. Your doctor will decide if further intervention is required.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The following information is intended for healthcare professionals only:

Each vial of Aldurazyme is intended for single use only. The concentrate for solution for infusion has to be diluted with sodium chloride 9 mg/ml (0.9%) solution for infusion using aseptic technique. It is recommended that the diluted Aldurazyme solution be administered to patients using an infusion set equipped with an 0.2  $\mu$ m in-line filter.

From a microbiological safety point of view, the product should be used immediately. If not used immediately, in-use storage should not be longer than 24 hours at 2°C - 8°C provided that dilution has taken place under controlled and validated aseptic conditions.

Aldurazyme should not be mixed with other medicinal products in the same infusion.

#### Preparation of the Aldurazyme Infusion (Use Aseptic Technique)

- Determine the number of vials to be diluted based on the individual patient's weight. Remove the required vials from the refrigerator approximately 20 minutes in advance in order to allow them to reach room temperature (below 30°C).
- Before dilution, visually inspect each vial for particulate matter and discoloration. The clear to slightly opalescent and colourless to pale yellow solution should be free of visible particles. Do not use vials exhibiting particles or discoloration.

Side effects were mainly seen while patients were being given the medicine or shortly after (infusion-associated reactions). If you experience any reaction like this, you should **contact your doctor immediately**. The number of these reactions decreased the longer that patients were on Aldurazyme. The majority of these reactions were mild or moderate in intensity. However, severe systemic allergic reaction (anaphylactic reaction) has been observed in patients during or up to 3 hours after Aldurazyme infusions. Some of the symptoms of such a severe allergic reaction were life-threatening and included extreme difficulty breathing, swelling of the throat, low blood pressure, and low oxygen level in the body. A few patients who had a prior history of severe MPS I related upper airway and pulmonary involvement, experienced severe reactions including bronchospasm (airway constriction), respiratory arrest, and swelling of the face. The frequency of bronchospasm and respiratory arrest is unknown. The frequency of severe allergic reaction (anaphylactic reaction) and swelling of the face is considered common and may affect up to 1 in 10 people.

Very common symptoms (may affect more than 1 in 10 people) which were not serious include

- headache,
- nausea,
- abdominal pain,
- rash,
- joint disease,
- joint pain,
- back pain,
- pain in arms or legs,
- flushing,
- fever, chills,
- increased heart rate,
- increased blood pressure,
- reaction at the infusion site such as swelling, redness, build-up of fluid, discomfort, itchy rash, pale colour of the skin, discoloured skin, or sensation of being warm.

Other side effects include the following:

**Common (may affect up to 1 in 10 people)**

- increased body temperature
- tingling
- dizziness
- cough
- difficulty in breathing
- vomiting
- diarrhoea
- rapid swelling under the skin in areas such as the face, throat, arms and legs which can be life-threatening if throat swelling blocks the airway
- hives
- itching
- hair loss
- cold sweat, heavy sweating
- muscle pain
- paleness
- cold hands or feet
- feeling hot, feeling cold
- fatigue
- influenza like illness
- pain at injection site
- restlessness

**Not known (frequency cannot be estimated from the available data)**

- allergic reactions (hypersensitivity)
- abnormally slower heart rate
- increased or abnormally high blood pressure
- voice box swelling
- bluish color of the skin (due to lower levels of oxygen in the blood)
- fast breathing
- redness of the skin

- leakage of the medicine into the surrounding tissue at the site of injection, where it can cause damage
- inability of the lungs to work properly (respiratory failure)
- throat swelling
- high-pitched breathing sound
- obstruction of airways causing difficulty in breathing
- lip swelling
- tongue swelling
- swelling especially of the ankles and feet due to fluid retention
- drug specific antibody, a blood protein produced in response to medicine
- antibody that neutralizes the effect of medicine

#### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the contact details below. By reporting side effects you can help provide more information on the safety of this medicine.

#### **United Kingdom**

Yellow Card Scheme

Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

#### **5. How to store Aldurazyme**

Keep this medicine out of the sight and reach of children.

You should not be given this medicine after the expiry date which is stated on the label after the letters EXP. The expiry date refers to the last day of that month.

#### Unopened vials:

Store in a refrigerator (2°C – 8°C).

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

#### **6. Contents of the pack and other information**

##### **What Aldurazyme contains**

- The active substance is laronidase. One ml of the solution in the vial contains 100 U of laronidase. Each vial of 5 ml contains 500 U of laronidase.
- The other ingredients are sodium chloride, sodium phosphate monobasic monohydrate, sodium phosphate dibasic heptahydrate, polysorbate 80, water for injections.

##### **What Aldurazyme looks like and contents of the pack**

Aldurazyme is supplied as a concentrate for solution for infusion. It is a solution that is clear to slightly opalescent, and colourless to pale yellow.

Pack size: 1, 10 and 25 vials per carton. Not all pack sizes may be marketed.

##### **Marketing Authorisation Holder and Manufacturer**

##### Marketing Authorisation Holder

Sanofi

410 Thames Valley Park Drive

Reading

Berkshire

RG6 1PT

UK

Tel: 0800 035 2525

Email: [uk-medicalinformation@sanofi.com](mailto:uk-medicalinformation@sanofi.com)

##### Manufacturer

EUROAPI UK Limited, 37 Hollands Road, Haverhill, Suffolk CB9 8PU, United Kingdom.

Genzyme Ireland Ltd., IDA Industrial Park, Old Kilmeaden Road, Waterford, Ireland

This leaflet does not contain all the information about your medicine. If you have any questions or are not sure about anything, ask your doctor or pharmacist.

**This leaflet was last revised in May 2024.**

7001442



- Determine the total volume of infusion based on the individual patient's weight, either 100 ml (if bodyweight is less or equal than 20 kg) or 250 ml (if bodyweight is more than 20 kg) of 0.9% sodium chloride intravenous solution.
- Withdraw and discard a volume of sodium chloride 9 mg/ml (0.9%) solution for infusion from the infusion bag equal to the total volume of Aldurazyme to be added.
- Withdraw the required volume from the Aldurazyme vials and combine the withdrawn volumes.
- Add the combined volumes of Aldurazyme to the sodium chloride 9 mg/ml (0.9%) solution for infusion.

- Mix the solution for infusion gently.
- Prior to use visually inspect the solution for particulate matter. Only clear and colourless solutions without visible particles should be used.

Any unused product or waste material should be disposed of in accordance with local requirements.