

**Package leaflet:  
Information for the user**

**Azithromycin 250 mg  
film-coated tablets  
Azithromycin 500 mg  
film-coated tablets**

Azithromycin

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist. This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Azithromycin is and what it is used for
2. What you need to know before you take Azithromycin
3. How to take Azithromycin
4. Possible side effects
5. How to store Azithromycin
6. Contents of the pack and other information

**1. What Azithromycin is and what it is used for**

Azithromycin contains the active substance azithromycin. Azithromycin is an antibiotic that belongs to a group of antibiotics known as macrolides, which block the growth of susceptible bacteria.

Azithromycin is taken for the treatment of the following infections:

Adults and adolescents weighing 45 kg and over

- Infections of the tonsils (tonsillitis) or throat (pharyngitis) caused by streptococcal bacteria
- Bacterial sinus infections (sinusitis)
- Bacterial infections of the middle ear (otitis media)
- Pneumonia (community-acquired pneumonia, not contracted in a hospital)
- Bacterial infections of the skin and underlying tissues
- Early localised Lyme disease (erythema migrans, mainly caused by tick bites)
- Bacterial infections of the gums (periodontitis) or abscess in gums (periodontal abscess)
- Infections of urethra and cervix caused by *Chlamydia trachomatis* bacteria
- Infections of urethra and cervix caused by *Neisseria gonorrhoeae* bacteria. Azithromycin should be used in combination with another antibiotic that is selected by your doctor or pharmacist.
- Chronic inflammation of the prostate caused by *Chlamydia trachomatis* bacteria
- Bacterial infections of genitals with painful sores (chancroid)
- Infections caused by *Mycobacterium avium* complex (MAC) bacteria in people with advanced HIV infection. Azithromycin should be used in combination with another antibiotic called ethambutol.

Azithromycin is also taken for prevention of infections caused by *Mycobacterium avium* complex (MAC) bacteria in people living with HIV infection.

**Adults:**

- Bacterial infections in patients with long-term inflammation of the lungs (chronic bronchitis)
  - Bacterial infection of the womb, fallopian tubes and ovaries (pelvic inflammatory disease) always in combination with another antibiotic(s) that is selected by your doctor or pharmacist.

**2. What you need to know before you take Azithromycin**

**Do not take Azithromycin**  
- if you are allergic to azithromycin, erythromycin, any macrolide or ketolide antibiotic, or any of the other ingredients of this medicine (listed in section 6).

**Warnings and precautions**

Talk to your doctor or pharmacist before taking Azithromycin if you have or have had any of the following conditions:

- heart problems (e.g. problems with your heart rhythm or cardiac insufficiency) or low levels of potassium or magnesium in your blood: these conditions may contribute to serious cardiac side effects of azithromycin
- liver problems: your doctor may need to monitor your liver function or stop the treatment
- severe diarrhoea after administration of any other antibiotics
- localised muscle weakness (myasthenia gravis), as the symptoms of this disease may worsen during treatment
- or if you are taking any ergot derivatives such as ergotamine (used to treat migraine) as these medicines should not be taken together with Azithromycin.

**Stop taking this medicine and contact your doctor immediately (see also "Serious side effects" in section 4):**

- if you feel you are having an allergic reaction (e.g. difficulty in breathing, swelling of the face or throat, rash, blistering).
- if you notice any of the symptoms as described in section 4 related to Stevens-Johnson syndrome, toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms (DRESS) and acute generalised exanthematous pustulosis (AGEP), which have been reported in association with azithromycin treatment.
- if you feel you have an abnormal heartbeat or palpitations, get dizzy or faint when taking Azithromycin.
- if you develop signs of liver problems (e.g. dark urine, loss of appetite or yellowing of the skin or whites of the eyes).
- if you develop severe diarrhoea during or after treatment. Do not take any medicine to treat your diarrhoea without first checking with your doctor. If your diarrhoea continues or reappears within the first weeks after treatment, please also inform your doctor.

Superinfection

Your doctor may observe you for signs of additional bacterial or fungal infections that cannot be treated with Azithromycin (superinfection).

Sexually transmitted infections

Your doctor may test for and exclude a potential infection with syphilis, a sexually transmitted disease that may otherwise progress undetected and be diagnosed delayed. Furthermore, in any case of sexually transmitted bacterial infections your doctor will initiate laboratory follow-up tests to monitor the success of therapy.

**Children and adolescents**

This medicinal product is not recommended if:

- you are less than 18 years old and have been diagnosed with pelvic inflammatory disease

- you are less than 12 years old and you are infected or at risk of being infected with organisms that belong to the *Mycobacterium avium* complex which usually affect people living with HIV who have low defences as its efficacy and safety have not been studied in these cases.

If you weigh less than 45 kg, other medicinal products containing azithromycin exist that may be more convenient for you to take.

**Other medicines and Azithromycin**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Taking Azithromycin at the same time as some other medicines may result in side effects. Therefore, it is particularly important that you tell your doctor if you are using any of the following medicines:

- Atorvastatin and other medicines from the statins group (to lower blood cholesterol and prevent heart disease, including heart attacks and strokes)
- Ciclosporin (to prevent rejection of organ transplants by the body)
- Colchicine (to treat gout and familial Mediterranean fever)
- Dabigatran (to prevent and treat blood clot formation (anticoagulant))
- Digoxin (to treat heart diseases)
- Warfarin or similar medicines used to thin the blood (anticoagulants)
- Medicines that may cause the heart muscle to take longer to contract and relax than usual (QT prolongation), such as the following:
  - Quinidine, procainamide, dofetilide, amiodarone and sotalol (to treat an irregular heartbeat, including a too fast or too slow heartbeat-cardiac arrhythmia)
  - Pimozide (to treat mental illness)
  - Citalopram (to treat depression)
  - Moxifloxacin and levofloxacin (antibiotics)
  - Cisapride (to treat disorders in the gastrointestinal tract)
  - Hydroxychloroquine or chloroquine (to treat autoimmune diseases including rheumatoid arthritis, or to treat or prevent malaria)

**Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy

Your doctor will decide if you should take this medicine during pregnancy, only after making sure that the benefits outweigh the potential risks.

Breast-feeding

Azithromycin passes into breast milk. Your doctor will decide therefore whether you should stop breast-feeding or should avoid treatment with Azithromycin taking into account both the benefit of breast-feeding for your child and the benefit of therapy for you.

**Driving and using machines**

Azithromycin has a moderate influence on the ability to drive and use machines. Azithromycin has been reported to cause dizziness, drowsiness and seizures, as well as problems with seeing and hearing in some people. These possible side effects may have an influence on your ability to drive and use machines.

**Azithromycin contains lactose monohydrate**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Sodium

Azithromycin contains less than 1 mmol (23 mg) of sodium per tablet, that is to say it is essentially 'sodium-free.'

**3. How to take Azithromycin**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The amount of Azithromycin that you need to take each day depends on the bacterial infection that you are being treated for and the specific treatment course that your doctor or pharmacist has instructed you to follow.

Adults and adolescents weighing at least 45 kg

Infection	Treatment course with azithromycin
Infections of the tonsils (tonsillitis) or throat (pharyngitis) caused by streptococcal bacteria	There is a 3-day or a 5-day treatment course for these infections, and the amount of Azithromycin to take each day is described for these treatment courses below.
Bacterial sinus infections (sinusitis)	
Bacterial infections of the middle ear (otitis media)	
Bacterial infections in patients with long-term inflammation of the lungs (chronic bronchitis)*	<b>3-day treatment course</b> 500 mg taken once daily for 3 days.
Pneumonia (community-acquired pneumonia, not contracted in a hospital)#	<b>5-day treatment course</b> 500 mg taken on the first day of treatment and then 250 mg taken once daily for the following 4 days.
Bacterial infections of the skin and underlying tissues	
Bacterial infections of the gums (periodontitis) or abscess in gums (periodontal abscess)	
Early localised Lyme disease (erythema migrans, mainly caused by tick bites)	1000 mg taken on the first day of treatment and then 500 mg taken once daily for the following 9 days.
Infections of urethra and cervix caused by <i>Chlamydia trachomatis</i> bacteria	1000 mg taken as a single dose
Infections of urethra and cervix caused by <i>Neisseria gonorrhoeae</i> bacteria. Azithromycin should be used in combination with another antibiotic that is selected by your doctor or pharmacist.	1000 mg* or 2000 mg* taken as a single dose
Bacterial infection of the womb, fallopian tubes and ovaries (pelvic inflammatory disease). Azithromycin should be used in combination with another antibiotic that is selected by your doctor or pharmacist**	Only if treatment has been started with intravenous azithromycin:  250 mg once daily to complete a 7day course of treatment

P15xxxxx

Black

Chronic inflammation of the prostate caused by <i>Chlamydia trachomatis</i> bacteria	500 mg/day taken on 3 consecutive days per week for a total of 3 weeks
Bacterial infections of genitals with painful sores (chancroid)	1000 mg taken as a single dose
Infections caused by <i>Mycobacterium avium</i> complex (MAC) bacteria in people with advanced HIV infection. Azithromycin should be used in combination with another antibiotic called ethambutol.	<500 mg> or <600 mg> once per day
Prevention of infections caused by <i>Mycobacterium avium</i> complex (MAC) bacteria in people living with HIV infection	<1200 mg> or <1250 mg> once per week

\*only for adult patients  
# for adult patients oral treatment may follow an initial intravenous treatment

**Use in children and adolescents**  
If your weight is less than 45 kg <or you are not able to swallow this medicinal product>, ask your doctor or pharmacist as other medicinal products containing azithromycin are also available that may be more appropriate for you.

#### Method of administration

For oral use.

**Azithromycin 250 mg:** This should be taken by mouth as a single daily dose. Tablets should be swallowed whole with some water, with or without a meal. Taking this medicine just before a meal may help make it easier on your stomach.

**Azithromycin 500 mg:** This should be taken by mouth as a single daily dose. Tablets may be taken with or without a meal. Taking this medicine just before a meal may help make it easier on your stomach. Tablets can be split in two equal halves which can be used to adjust the dose as your doctor or pharmacist has told you.

#### If you take more Azithromycin than you should

If you take more Azithromycin than you should then you may feel unwell. Typical signs of an overdose are vomiting, diarrhoea, abdominal pain and nausea. Tell your doctor or contact your nearest hospital emergency department immediately.

#### If you forget to take Azithromycin

If you forget to take Azithromycin, take it as soon as you can, as long as this is at least 12 hours before the next dose is due. If it is less than 12 hours left to your next dose, skip the missed dose and take the next dose at the usual time. Do not take a double dose to make up for a forgotten dose.

#### If you stop taking Azithromycin

If you stop taking Azithromycin too soon, the infection may return. Take Azithromycin for the full time of treatment, even when you begin to feel better.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

#### Serious side effects

**Stop using Azithromycin and seek medical attention immediately if you notice any of the following symptoms:**

- sudden wheeziness, difficulty in breathing, swelling of eyelids, face or lips, rash or itching especially affecting the whole body (*anaphylactic reaction*, frequency not known)
- rapid or irregular heartbeat (*cardiac arrhythmia* or *torsades de pointes tachycardia*, frequency not known)
- dark urine, loss of appetite or yellowing of the skin or whites of the eyes, which are signs of liver disorders (*hepatic failure* or *hepatic necrosis* (frequency not known), *hepatitis\** (uncommon: may affect up to 1 in 100 people)).
- severe diarrhoea with abdominal cramps, bloody stools and/or fever may mean that you have an infection of the large intestine (*antibiotic-associated colitis*, frequency not known). Do not take medicines against diarrhoea that inhibit the bowel movements (*antiperistaltics*).
- reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by fever and flu-like symptoms (*Stevens-Johnson syndrome\** or *toxic epidermal necrolysis*, frequency not known).
- widespread rash, high body temperature and enlarged lymph nodes (*DRESS syndrome* or *drug hypersensitivity syndrome*, rare (may affect up to 1 in 1,000 people)).
- a red, scaly widespread rash with bumps under the skin and blisters accompanied by fever. The symptoms usually appear at the initiation of treatment (*acute generalised exanthematous pustulosis*, rare (may affect up to 1 in 1,000 people)).

#### Other side effects

**Very common (may affect more than 1 in 10 people)**

- diarrhoea
- abdominal discomfort\*

**Common (may affect up to 1 in 10 people)**

- headache
- being sick (*vomiting*), stomach pain#, feeling sick (*nausea*)#
- changes in blood test results (*lymphocyte count decreased, eosinophil count increased, basophil count increased, monocyte count increased, neutrophil count increased, blood bicarbonate decreased*)

**Uncommon (may affect up to 1 in 100 people)**

- thrush (*candidiasis*) - a fungal infection of the mouth and vagina, other fungal infections
- pneumonia, bacterial infection of the throat, inflammation of the gastrointestinal tract, respiratory disorder, inflammation of the mucous membrane inside the nose, vaginal infection
- changes in the number of white blood cells (*leukopenia, neutropenia, eosinophilia*)
- platelet count increased
- reduction in the proportion of all blood cells in the total blood volume (*hematocrit decreased*)
- allergic reactions, swelling of the hands, feet and face (*angioedema*)
- lack of appetite#
- nervousness, difficulty sleeping (*insomnia*)
- feeling dizzy#, feeling drowsy (somnolence), change in your sense of taste (*dysgeusia*)#, sensation of pins and needles or numbness (*paraesthesia*)#
- impaired vision#
- ear disorder
- spinning sensation (*vertigo*)

- feeling your heartbeat (*palpitations*)
- hot flush
- sudden wheeziness, bleeding from the nose
- constipation, wind#, impaired digestion (*dyspepsia*), inflammation of the lining of the stomach (*gastritis*), difficulty in swallowing (*dysphagia*), swollen belly, dry mouth, belching (*eructation*), mouth ulceration, increased salivation
- rash#, itching#, hives (*urticaria*), dermatitis, dry skin, abnormally increased sweating (*hyperhidrosis*)
- swelling and pain in the joints (*osteoarthritis*), muscle pain, back pain, neck pain
- painful urination (*dysuria*), kidney pain
- menstrual bleeding at irregular intervals (*metrorrhagia*), testicular disorder
- swelling due to fluid retention, especially of the face, ankles and feet (oedema, face oedema, peripheral oedema)
- weakness, tiredness#, general feeling of being unwell, fever
- chest pain, pain
- abnormal laboratory test results (e.g. blood or liver tests)
- post procedural complication

**Rare (may affect up to 1 in 1,000 people)**

- feeling irritated
- liver problems, yellowing of the skin or eyes
- increased sensitivity to sunlight#

**Not known (frequency cannot be estimated from the available data)**

- reduced number of red blood cells due to increased cell breakdown which can cause tiredness and pale skin (*haemolytic anaemia*)
- reduction in number of platelets which can lead to bleeding and bruising (*thrombocytopenia*)
- feeling angry, aggressive, feeling of fear and concern (*anxiety*), acute confusional state (*delirium*), hallucination
- fainting (*syncope*)
- fits (*seizures*)
- reduced sensation to touch, pain and temperature (*hypoesthesia*)#
- feeling hyperactive
- change in your sense of smell (*anosmia, parosmia*)
- total loss of your sense of taste (*ageusia*)
- muscle weakness (*myasthenia gravis*)
- abnormal electrocardiogram (ECG) heart tracing (*QT prolongation*)
- deafness#, reduced hearing# or ringing in your ears (*tinnitus*)#
- low blood pressure
- inflammation of the pancreas causing severe pain in the belly and back (*pancreatitis*)
- your tongue changes colour
- joint pain (*arthralgia*)#
- kidney inflammation (*interstitial nephritis*) and kidney failure

\* These side effects were only seen during azithromycin administration for the prophylaxis and/or therapy of *Mycobacterium avium* complex infections in people living with HIV with insufficient recovery of the immune system.

# These side effects were more common during azithromycin administration for the prophylaxis and/or therapy of *Mycobacterium avium* complex infections in people living with HIV with insufficient recovery of the immune system.

#### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via Yellow Card Scheme Website:

[www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Azithromycin

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

The medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

What Azithromycin contains

- The active substance is azithromycin dihydrate. Each film-coated tablet contains 250 mg of azithromycin (as dihydrate). Each film-coated tablet contains 500 mg of azithromycin (as dihydrate). The other ingredients are

**Tablet core:** Calcium hydrogen phosphate, starch, pregelatinized (maize starch), croscarmellose sodium, sodium lauryl sulfate, magnesium stearate.

**Tablet coating:** Lactose monohydrate, hypromellose, titanium dioxide (E 171), triacetin.

#### What Azithromycin looks like and contents of the pack

Film-coated tablet.

##### Azithromycin 250 mg film-coated tablets:

White to off-white, oblong shaped, film coated biconvex tablets, debossed with "66" on one side and "D" on other side. The size is 13.5 mm x 6.6 mm

##### Azithromycin 500 mg film-coated tablets:

White to off-white, oval shaped, film coated biconvex tablets debossed with "6" and "7" on either side of the score-line on one side and "D" on other side. The size is 17.1 mm x 8.5 mm

The tablet can be divided into equal doses. Azithromycin film-coated tablets are available in clear PVC-Aluminium blister packs.

**Blister packs:** 2,3,4,6 and 12 film-coated tablets

Not all pack sizes may be marketed.

#### Marketing Authorisation Holder and Manufacturer

Milpharm Limited  
Ares Block, Odyssey  
Business Park  
West End Road  
Ruislip HA4 6QD  
United Kingdom

#### Manufacturer

APL Swift Services (Malta) Limited  
Hf26, Hal Far Industrial Estate, Hal Far  
Birzebbugia, BBG 3000  
Malta

or

Milpharm Limited  
Ares Block, Odyssey  
Business Park  
West End Road  
Ruislip HA4 6QD  
United Kingdom

**This leaflet was last revised in 05/2026.**