

Package leaflet: Information for the user

Benepali 25 mg solution for injection in pre-filled syringe etanercept

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- Your doctor will also give you a Patient Card, which contains important safety information that you need to be aware of before and during treatment with Benepali.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you or a child in your care. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours or those of the child you are caring for.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Benepali is and what it is used for
2. What you need to know before you use Benepali
3. How to use Benepali
4. Possible side effects
5. How to store Benepali
6. Contents of the pack and other information
7. Instructions for use (see overleaf)

1. What Benepali is and what it is used for

Benepali contains the active substance etanercept.

Benepali is a medicine that is made from two human proteins. It blocks the activity of another protein in the body that causes inflammation. Benepali works by reducing the inflammation associated with certain diseases.

In adults (aged 18 and over), Benepali can be used for:

- moderate or severe **rheumatoid arthritis**;
- **psoriatic arthritis**;
- severe **axial spondyloarthritis** including **ankylosing spondylitis**;
- moderate or severe **plaque psoriasis**.

In each case Benepali is used, usually when other widely used treatments have not worked well enough or are not suitable for you.

For **rheumatoid arthritis**, Benepali is usually used in combination with methotrexate, although it may also be used alone if treatment with methotrexate is unsuitable for you. Whether used alone or in combination with methotrexate, Benepali can slow down the damage to your joints caused by the rheumatoid arthritis and improve your ability to do normal daily activities.

For **psoriatic arthritis** patients with multiple joint involvement, Benepali can improve your ability to do normal daily activities.

For patients with **multiple symmetrical painful or swollen joints** (e.g., hands, wrists and feet), Benepali can slow down the structural damage to those joints caused by the disease.

Benepali is also prescribed for the treatment of the following diseases in children and adolescents.

- For the following types of juvenile idiopathic arthritis when treatment with methotrexate has not worked well enough or is not suitable for them:
 - Polyarthritis (rheumatoid factor positive or negative) and extended oligoarthritis in patients from the age of 2 years and weighing 62.5 kg or more.
 - Psoriatic arthritis in patients from the age of 12 years and weighing 62.5 kg or more.
- For enthesitis-related arthritis in patients from the age of 12 years and weighing 62.5 kg or more when other widely used treatments have not worked well enough or are not suitable for them.
- Severe plaque psoriasis in patients from the age of 6 years and weighing 62.5 kg or more who have had an inadequate response to (or are unable to take) phototherapies or other systemic therapies.

2. What you need to know before you use Benepali

Do not use Benepali

- if you or the child you are caring for, are **allergic to etanercept** or any of the other **ingredients of this medicine** (listed in section 6). If you or the child experience allergic reactions such as chest tightness, wheezing, dizziness or rash, do not inject more Benepali, and contact your doctor immediately.
- if you or the child have, or are at risk of developing a **serious blood infection** called sepsis. If you are not sure, please contact your doctor.
- if you or the child have an **infection of any kind**. If you are not sure, please talk to your doctor.

Warnings and precautions

Talk to your doctor before taking Benepali.

- **Allergic reactions:** If you or the child experience allergic reactions such as chest tightness, wheezing, dizziness or rash, do not inject more Benepali, and contact your doctor immediately.
- **Infections/surgery:** If you or the child develop a new infection, or are about to have any major surgery, your doctor may wish to monitor the treatment with Benepali.
- **Infections/diabetes:** Tell your doctor if you or the child have a history of recurrent infections or suffer from diabetes or other conditions that increase the risk of infection.
- **Infections/monitoring:** Tell your doctor of any recent travel outside the European region. If you or the child develop symptoms of an infection such as fever, chills or cough, notify your doctor immediately. Your doctor may decide to continue to monitor you or the child for the presence of infections after you stop using Benepali.
- **Tuberculosis:** As cases of tuberculosis have been reported in patients treated with Benepali, your doctor will check for signs and symptoms of tuberculosis before starting Benepali. This may include a thorough medical history, a chest X-ray and a tuberculin test. The conduct of these tests should be recorded on the Patient Card. It is very important that you tell your doctor if you or the child have ever had tuberculosis, or have been in close contact with someone who has had tuberculosis. If symptoms of tuberculosis (such as persistent cough, weight loss, listlessness, mild fever), or any other infection appear during or after therapy, tell your doctor immediately.
- **Hepatitis B:** Tell your doctor if you or the child have or have ever had hepatitis B. Your doctor should test for the presence of hepatitis B infection before you or the child begin treatment with Benepali. Treatment with Benepali may result in reactivation of hepatitis B in patients who have previously been infected with the hepatitis B virus. If this occurs, you should stop using Benepali.

- **Hepatitis C:** Tell your doctor if you or the child have hepatitis C. Your doctor may wish to monitor the treatment with Benepali in case the infection worsens.
- **Blood disorders:** Seek medical advice immediately if you or the child have any signs or symptoms such as persistent fever, sore throat, bruising, bleeding or paleness. Such symptoms may point to the existence of potentially life-threatening blood disorders, which may require discontinuation of Benepali.
- **Nervous system and eye disorders:** Tell your doctor if you or the child have multiple sclerosis, optic neuritis (inflammation of the nerves of the eyes) or transverse myelitis (inflammation of the spinal cord). Your doctor will determine if Benepali is an appropriate treatment.
- **Congestive heart failure:** Tell your doctor if you or the child have a history of congestive heart failure, because Benepali needs to be used with caution under these circumstances.
- **Cancer:** Tell your doctor if you have or have ever had lymphoma (a type of blood cancer) or any other cancer before you are given Benepali. Patients with severe rheumatoid arthritis, who have had the disease for a long time, may be at higher than average risk of developing lymphoma. Children and adults taking Benepali may have an increased risk of developing lymphoma or another cancer. Some children and teenage patients who have received etanercept or other medicines that work the same way as etanercept have developed cancers, including unusual types, which sometimes resulted in death. Some patients receiving Benepali have developed skin cancers. Tell your doctor if you or the child develop any change in the appearance of the skin or growths on the skin.
- **Chickenpox:** Tell your doctor if you or the child are exposed to chickenpox when using Benepali. Your doctor will determine if preventive treatment for chickenpox is appropriate.
- **Alcohol abuse:** Benepali should not be used for the treatment of hepatitis related to alcohol abuse. Please tell your doctor if you or the child in your care have a history of alcohol abuse.
- **Wegener's granulomatosis:** Benepali is not recommended for the treatment of Wegener's granulomatosis, a rare inflammatory disease. If you or the child in your care have Wegener's granulomatosis, talk to your doctor.
- **Anti-diabetic medicines:** Tell your doctor if you or the child have diabetes or are taking medicines to treat diabetes. Your doctor may decide if you or the child need less anti-diabetic medicine while taking Benepali.
- **Vaccinations:** Some vaccines, such as oral polio vaccine, should not be given during Benepali therapy. Please consult your doctor before you or the child receive any vaccines.

Children and adolescents

Benepali is not indicated for use in children and adolescents who weigh less than 62.5 kg.

- **Vaccinations:** If possible, children should be up to date with all vaccinations before using Benepali. Some vaccines, such as oral polio vaccine, should not be given while using Benepali. Please consult your doctor before you or the child receive any vaccines.
- **Inflammatory bowel disease (IBD):** There have been cases of IBD in patients with juvenile idiopathic arthritis (JIA) treated with etanercept. Tell the doctor if the child develops any abdominal cramps and pain, diarrhoea, weight loss or blood in the stool.

Benepali should not normally be used in children with polyarthritis or extended oligoarthritis below the age of 2 years or weighing less than 62.5 kg, or in children with enthesitis-related arthritis or psoriatic arthritis below the age of 12 years or weighing less than 62.5 kg, or in children with psoriasis below the age of 6 years or weighing less than 62.5 kg.

Other medicines and Benepali

Tell your doctor or pharmacist if you or the child are taking, have recently taken or might take any other medicines (including sulfasalazine).

You or the child should **not use** Benepali with medicines that contain the active substance anakinra or abatacept.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Women of childbearing potential should be advised to use appropriate contraception to avoid becoming pregnant during Benepali therapy and for three weeks after discontinuation of therapy.

Benepali should only be used during pregnancy if clearly needed.

If you received Benepali during pregnancy, your baby may have a higher risk of getting an infection. In addition, one study found more birth defects when the mother had received etanercept in pregnancy, compared with mothers who had not received etanercept or other similar medicines (TNF-antagonists), but there was no particular kind of birth defect reported. Another study found no increased risk of birth defects when the mother had received etanercept in pregnancy. Your doctor will help you to decide whether the benefits of treatment outweigh the potential risk to your baby. It is important that you tell the baby's doctors and other healthcare professionals about the use of Benepali during pregnancy before the baby receives any vaccine (for more information see section 2, "Vaccinations").

Women using Benepali should not breast-feed, since Benepali passes into human breast milk.

Driving and using machines

No information is available if the use of Benepali affects the ability to drive or use machines.

Benepali contains sodium

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

3. How to use Benepali

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

If you feel that the effect of Benepali is too strong or too weak, talk to your doctor or pharmacist.

Use in adult patients (aged 18 years or over)

Rheumatoid arthritis, psoriatic arthritis, and axial spondyloarthritis including ankylosing spondylitis

The usual dose is 25 mg given twice a week or 50 mg once a week as an injection under the skin. However, your doctor may determine an alternative frequency at which to inject Benepali.

Plaque psoriasis

The usual dose is 25 mg twice a week or 50 mg once a week.

Alternatively, 50 mg may be given twice a week for up to 12 weeks, followed by 25 mg twice a week or 50 mg once a week.

Your doctor will decide how long you should use Benepali and whether retreatment is needed based on your response. If Benepali has no effect on your condition after 12 weeks, your doctor may tell you to stop using this medicine.

Use in children and adolescents

Dose and frequency of dosing for the child or adolescent will depend on body weight and disease. Your doctor will determine the correct dose for the child and will prescribe an appropriate strength of etanercept.

Paediatric patients weighing 62.5 kg or more can be dosed 25 mg given twice a week or 50 mg given once a week using a fixed-dose pre-filled syringe or pre-filled pen.

Other etanercept products with appropriate dosage forms for children are available.

For polyarthritis or extended oligoarthritis in patients from the age of 2 years and weighing 62.5 kg or more, or enthesitis-related arthritis or psoriatic arthritis in patients from the age of 12 years and weighing 62.5 kg or more, the usual dose is 25 mg given twice a week or 50 mg given once a week.

For psoriasis in patients from the age of 6 years and weighing 62.5 kg or more, the usual dose is 50 mg and should be given once weekly. If Benepali has no effect on the child's condition after 12 weeks, your doctor may tell you to stop using this medicine.

The doctor will provide you with detailed directions for preparing and measuring the appropriate dose.

Method and route of administration

Benepali is administered by an injection under the skin (subcutaneous use).

Detailed instructions on how to inject Benepali are provided in section 7 "Instructions for use".

Do not mix the Benepali solution with any other medicine.

To help you remember, it may be helpful to write in a diary which day(s) of the week Benepali should be used.

If you use more Benepali than you should

If you have used more Benepali than you should (either by injecting too much on a single occasion or by using it too frequently), **talk to a doctor or pharmacist immediately**. Always have the outer carton of the medicine with you, even if it is empty.

If you forget to use Benepali

If you forget a dose, you should inject it as soon as you remember, unless the next scheduled dose is the next day; in which case you should skip the missed dose. Then continue to inject the medicine on the usual day(s). If you do not remember until the day that the next injection is due, do not take a double dose (two doses on the same day) to make up for a forgotten dose.

If you stop using Benepali

Your symptoms may return upon discontinuation.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Allergic reactions

If any of the following happen, do not inject more Benepali. **Tell your doctor immediately, or go to the casualty department at your nearest hospital.**

- Trouble swallowing or breathing
- Swelling of the face, throat, hands, or feet
- Feeling nervous or anxious, throbbing sensations, sudden reddening of the skin and/or a warm feeling
- Severe rash, itching, or hives (elevated patches of red or pale skin that often itch)

Serious allergic reactions are rare. However, any of the above symptoms may indicate an allergic reaction to Benepali, so you should seek immediate medical attention.

Serious side effects

If you notice any of the following, you or the child may need urgent medical attention.

- Signs of **serious infections** (including pneumonia, deep skin infections, joint infections and blood infection), such as high fever that may be accompanied by cough, shortness of breath, chills, weakness, or a hot, red, tender, sore area on the skin or joints;
- Signs of **blood disorders**, such as bleeding, bruising, or paleness;
- Signs of **nerve disorders**, such as numbness or tingling, changes in vision, eye pain, or onset of weakness in an arm or leg;
- Signs of **heart failure** or **worsening heart failure**, such as fatigue or shortness of breath with activity, swelling in the ankles, a feeling of fullness in the neck or abdomen, night-time shortness of breath or coughing, bluish colour of the nails or the lips;
- Signs of **cancers**: Cancers may affect any part of the body including the skin and blood, and possible signs will depend on the type and location of the cancer. These signs may include weight loss, fever, swelling (with or without pain), persistent cough, presence of lumps or growths on the skin;
- Signs of **autoimmune reactions** (where antibodies are made that may harm normal tissues in the body) such as pain, itching, weakness, and abnormal breathing, thinking, sensation, or vision;
- Signs of lupus or lupus-like syndrome, such as weight changes, persistent rash, fever, joint or muscle pain, or fatigue;
- Signs of **inflammation of the blood vessels** such as pain, fever, redness or warmth of the skin, or itching.

These are rare or uncommon side effects, but are serious conditions (some of which may rarely be fatal). If any of the above occurs, tell your doctor immediately, or visit the casualty department at your nearest hospital.

Other side effects

The known side effects of Benepali include the following in groups of decreasing frequency:

- **Very common** (may affect more than 1 in 10 people)
Infections (including colds, sinusitis, bronchitis, urinary tract infections and skin infections); injection site reactions (including bleeding, bruising, redness, itching, pain, and swelling). Reactions at the injection site (these do not occur as often after the first month of treatment). Some patients have developed a reaction at an injection site that was used before.
- **Common** (may affect up to 1 in 10 people)
Allergic reactions; fever; rash; itching; antibodies directed against normal tissue (autoantibody formation).
- **Uncommon** (may affect up to 1 in 100 people)
Serious infections (including pneumonia, deep skin infections, joint infections, blood infection, and infections at various sites); worsening of congestive heart failure; low red blood cell count, low white blood cell count, low neutrophil (a type of white blood cell) count; low blood platelet count; skin cancer (excluding melanoma); localised swelling of the skin (angioedema); hives (elevated patches of red or pale skin that often itch); eye inflammation; psoriasis (new or worsening); inflammation of the blood vessels affecting multiple organs; elevated liver blood tests (in patients also receiving methotrexate treatment, the frequency of elevated liver blood tests is common); abdominal cramps and pain, diarrhoea, weight loss or blood in the stool (signs of bowel problems).
- **Rare** (may affect up to 1 in 1,000 people)
Serious allergic reactions (including severe localised swelling of the skin and wheezing); lymphoma (a type of blood cancer); leukaemia (cancer affecting the blood and bone marrow); melanoma (a type of skin cancer); combined low platelet, red, and white blood cell count; nervous system disorders (with severe muscle weakness and signs and symptoms similar to

those of multiple sclerosis or inflammation of the nerves of the eyes or spinal cord); tuberculosis; new onset congestive heart failure; seizures; lupus or lupus-like syndrome (symptoms may include persistent rash, fever, joint pain, and tiredness); skin rash, which may lead to severe blistering and peeling of the skin; inflammation of the liver caused by the body's own immune system (autoimmune hepatitis; in patients also receiving methotrexate treatment, the frequency is uncommon); immune disorder that can affect the lungs, skin and lymph nodes (sarcoidosis); inflammation or scarring of the lungs (in patients also receiving methotrexate treatment, the frequency of inflammation or scarring of the lungs is uncommon); lichenoid reactions (itchy reddish-purple skin rash and/or threadlike white-grey lines on mucous membranes), opportunistic infections (which include tuberculosis and other infections that occur when resistance to disease is lowered), erythema multiforme (inflammatory skin rash), cutaneous vasculitis (inflammation of blood vessels in the skin); damage to nerves, including Guillain-Barré syndrome (a serious condition which can affect breathing and damage body organs).

- **Very rare** (may affect up to 1 in 10,000 people)
Failure of the bone marrow to produce crucial blood cells; toxic epidermal necrolysis (a life-threatening skin condition).
- **Not known** (frequency cannot be estimated from available data)
Merkel cell carcinoma (a type of skin cancer); Kaposi's sarcoma (a rare cancer related to infection with human herpes virus 8. Kaposi's sarcoma most commonly appears as purple lesions on the skin); excessive activation of white blood cells associated with inflammation (macrophage activation syndrome); recurrence of hepatitis B (a liver infection); worsening of a condition called dermatomyositis (muscle inflammation and weakness with an accompanying skin rash); listeria (a bacterial infection).

Side effects in children and adolescents

The side effects and their frequencies seen in children and adolescents are similar to those described above.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme.

Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Benepali

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and the label of the pre-filled syringe after EXP. The expiry date refers to the last day of that month.

Store in a refrigerator (2°C - 8°C). Do not freeze.

Keep the pre-filled syringes in the outer carton in order to protect from light.

After taking a syringe from the refrigerator, **wait approximately 30 minutes to allow the Benepali solution in the syringe to reach room temperature**. Do not warm in any other way. Immediate use is then recommended.

Benepali may be stored outside of the refrigerator at temperatures up to a maximum of 25°C for a

single period of up to four weeks; after which, it should not be refrigerated again. Benepali should be discarded if not used within four weeks after removal from the refrigerator. It is recommended that you record the date that Benepali is removed from the refrigerator and the date after which Benepali should be discarded (no more than 4 weeks following the removal from the refrigerator).

Inspect the solution in the syringe. It should be clear to slightly opalescent, colourless or pale yellow, and may contain small white or almost transparent particles of protein. This appearance is normal for Benepali. Do not use this medicine if you notice the solution is discoloured, cloudy, or if particles other than those described above are present. If you are concerned with the appearance of the solution, then contact your pharmacist for assistance.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Benepali contains

- The active substance is etanercept. Each pre-filled syringe contains 25 mg of etanercept.
- The other ingredients are sucrose, sodium chloride, sodium dihydrogen phosphate monohydrate, disodium hydrogen phosphate heptahydrate, and water for injections (see section 2 “Benepali contains sodium”).

What Benepali looks like and contents of the pack

Benepali is supplied as a pre-filled syringe containing a clear to slightly opalescent, colourless or pale yellow solution for injection (injection).

Benepali is available in packs containing 4 pre-filled syringes, multipacks comprising 2 cartons, each containing 4 pre-filled syringes and multipacks comprising 6 cartons, each containing 4 pre-filled syringes. Not all pack sizes may be marketed.

Marketing Authorisation Holder

Samsung Bioepis NL B.V.
Olof Palmestraat 10
2616 LR Delft
The Netherlands

Manufacturer

FUJIFILM Diosynth Biotechnologies Denmark ApS
Biotek Allé 1
Hillerød, 3400
Denmark

Biogen Netherlands B.V.
Prins Mauritslaan 13
1171 LP Badhoevedorp
The Netherlands

Samsung Bioepis NL B.V.
Olof Palmestraat 10
2616 LR Delft
The Netherlands

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

Biogen Belgium NV/S.A
Tél/Tel: + 32 (0)2 808 5947

България

Ewopharma AG Representative Office
Тел.: + 359 249 176 81

Česká republika

Biogen (Czech Republic) s.r.o.
Tel: + 420 228 884 152

Danmark

Biogen (Denmark) A/S
Tlf: + 45 78 79 37 53

Deutschland

Biogen GmbH
Tel: + 49 (0)30 223 864 72

Eesti

Ewopharma AG Eesti filiaal
Tel: + 372 6 68 30 56

Ελλάδα

Genesis Pharma S.A.
Τηλ: + 30 211 176 8555

España

Biogen Spain, S.L.
Tel: + 34 931 790 519

France

Biogen France SAS
Tél: + 33 (0)1 776 968 14

Hrvatska

Ewopharma d.o.o
Tel: + 385 (0)1 777 64 37

Ireland

Biogen Idec (Ireland) Ltd.
Tel: +353 (0)1 513 33 33

Ísland

Icepharma hf.
Sími: + 354 800 9836

Italia

Biogen Italia s.r.l.
Tel: + 39 (0)6 899 701 50

Κύπρος

Genesis Pharma (Cyprus) Ltd
Τηλ: + 357 22 00 04 93

Lietuva

Ewopharma AG Atstovybė
Tel: +370 52 07 91 38

Luxembourg/Luxemburg

Biogen Belgium NV/SA
Tél/Tel: +35 227 772 038

Magyarország

Biogen Hungary Kft.
Tel.: + 36 1 848 04 64

Malta

Pharma.MT Ltd
Tel: + 356 27 78 15 79

Nederland

Biogen Netherlands B.V.
Tel: + 31 (0)20 808 02 70

Norge

Biogen Norway AS
Tlf: + 47 21 93 95 87

Österreich

Biogen Austria GmbH
Tel: + 43 (0)1 267 51 42

Polska

Biogen Poland Sp. z o.o.
Tel.: + 48 22 116 86 94

Portugal

Biogen Portugal Sociedade Farmacêutica,
Unipessoal, Lda
Tel: + 351 308 800 792

România

Ewopharma AG Representative Office
Tel: + 40 377 881 045

Slovenija

Biogen Pharma d.o.o.
Tel: + 386 (0)1 888 81 07

Slovenská republika

Biogen Slovakia s.r.o.
Tel: + 421 (0)2 333 257 10

Suomi/Finland

Biogen Finland Oy
Puh/Tel: + 358 (0)9 427 041 08

Sverige

Biogen Sweden AB
Tel: +46 (0)8 525 038 36

Latvija

Ewopharma AG pārstāvniecība
Tel: + 371 66 16 40 32

United Kingdom (Northern Ireland)

Biogen Idec (Ireland) Ltd.
Tel: +44 (0)20 360 886 22

This leaflet was last revised in 03/2021

Detailed information on this medicine is available on the European Medicines Agency web site:
<http://www.ema.europa.eu>.

7. Instructions for use


Read the instructions for use before you start using Benepali and each time you get a refill of your prescription. There may be new information.

- **Do not** try to give yourself the injection unless your doctor or nurse has shown you how to give the injection.

A single-use pre-filled syringe contains one 25 mg dose of Benepali.

Find a well-lit, clean surface and gather all the equipment you need:

- **A new Benepali pre-filled syringe**



- **Do not** shake the pre-filled syringe.

Not included in pack:

- **1 alcohol swab, gauze pad and plaster**



- **Sharps disposal container**



A. Before you start

1. Inspect the pre-filled syringe:

Check the expiry date on the pre-filled syringe label.

- **Do not** use the pre-filled syringe past the expiration date.
- **Do not** use the pre-filled syringe if it has been dropped onto a hard surface. Components inside the pre-filled syringe may be broken.
- **Do not** use the pre-filled syringe if the needle cover is missing or not securely attached.

2. Inspect the solution:

Look at the medicine in the pre-filled syringe.

The medicine should be clear to slightly opalescent, colourless or pale yellow, and may contain small white or almost transparent particles of protein.

- **Do not** use the solution if it is discoloured, cloudy, or if particles other than those described above are present.


3. Allow the medicine to reach room temperature:

Remove one pre-filled syringe from the refrigerator and leave at room temperature for at least 30 minutes before injecting.

This is important to make the medicine easier and more comfortable to inject.

- **Do not** remove the needle cover until you are ready to inject.
- **Do not** use heat sources, such as a microwave or hot water, to warm Benepali.

4. Choose an injection site:



The Benepali pre-filled syringe is for a subcutaneous injection. It should be injected into the thigh, abdomen, or back of the upper arm (see image on the left).


Rotate the site for each injection.

If you are injecting into the abdomen, choose a site that is at least 5 cm away from the belly button.

- **Do not** inject into areas that are red, hard, bruised, or tender.
- **Do not** inject into scars or stretch marks.
- If you have psoriasis, **do not** inject into any raised, thick, red, or scaly skin patches, or lesions.


B. Injection steps

Step 1:



Wash your hands with soap and water.

Step 2:




Wipe the skin at the injection site with an alcohol swab.

See 'Choose an injection site' for guidance with choosing an injection site.

- **Do not** touch this area again before giving the injection.

Step 3:




Pull the needle cover straight off and dispose of it in the bin or sharps container.

See 'Choose an injection site' for guidance with choosing an injection site.

- **Do not** twist or bend the needle cover while removing it, as this may damage the needle.
- **Do not** touch the plunger rod while removing the needle cover.
- **Never recap the needle.**

Step 4:




Gently pinch a fold of skin at the cleaned injection site.

Hold the pre-filled syringe at 45 degrees to the skin. With a quick, dart-like motion, insert the needle fully into the skin.


You can let go of the skin that you are pinching after the needle is completely inserted.

Step 5:



Slowly push down the plunger to inject all of the Benepali solution.

Step 6:



When the syringe is empty, remove the needle from the skin at the same angle as inserted.

- **Never** recap the needle. Recapping could lead to a needle stick injury.

Disposal:



Dispose of the entire syringe in an approved sharps container.

Check with your healthcare provider for instructions on how to properly dispose a filled sharps container. Sharps disposal containers may be purchased at your local pharmacy.

- **Do not** throw the sharps container in household bin.
- **Do not** recycle.
- **Do not** reuse the Benepali pre-filled syringe.
- **Always keep the container out of the sight and reach of children.**

C. Injection site care

If there is bleeding at the injection site, press a gauze pad over the injection site.

- **Do not** rub the injection site.

If needed, cover the injection site with a plaster.