



PACKAGE LEAFLET  
Information for the user

# IBULEVE™ GEL

ibuprofen 5% w/w

## Read all of this leaflet carefully before using this product.

Keep this leaflet. You may need to read it again.

Ask your doctor or pharmacist if you need more information or advice.

You must contact a doctor if your symptoms worsen or do not improve after a few weeks.

If any of the side effects get serious or if you notice any side effects not listed in this leaflet please tell your doctor or pharmacist.

### In this leaflet:

1. What Ibuleve is and what it is used for
2. Before you use Ibuleve
3. How to use Ibuleve
4. Possible side effects
5. How to store Ibuleve
6. Further information

### 1. WHAT IBULEVE IS AND WHAT IT IS USED FOR

- Ibuleve is an anti-inflammatory painkiller applied to, and absorbed through, the skin.
- It is for the treatment of the following conditions involving the musculoskeletal system:
  - backache
  - rheumatic or muscular pain
  - sprains
  - strains
  - neuralgia.
- It is also for pain relief in common arthritic conditions.
- Ibuleve is recommended for use by **adults, the elderly and children over the age of 12 years**. Children under the age of 12 may also use it if recommended by their doctor.
- The **active ingredient** in this product is ibuprofen. This is one of a group of medicines known as **non-steroidal anti-inflammatory drugs (NSAIDs)**.
- Ibuprofen works by:
  - relieving pain
  - reducing swelling and inflammation.

### 2. BEFORE YOU USE IBULEVE

#### Do not use Ibuleve:

- if you are **allergic (hypersensitive)** to ibuprofen or any of the other ingredients of Ibuleve listed in Section 6;
- if you are **asthmatic**, or suffer from **rhinitis** (allergic runny nose) or **urticaria** (hives) **and** have ever had a bad reaction to aspirin, ibuprofen or other NSAIDs in the past;
- if you are **pregnant or breast-feeding**;
- on broken, damaged, infected or diseased skin.

**Before applying this product for the first time**, make sure it is suitable for you to use:

Because Ibuleve is delivered through the skin, directly over the painful area, there is less risk of the complications that sometimes occur when ibuprofen (or a similar anti-inflammatory painkiller) is taken by mouth. However, in rare cases you may be at risk:

- if you have a stomach ulcer (also called a peptic or gastric ulcer);

- if you have ever had kidney problems;
- if you have ever had asthma;
- if you have ever had a bad reaction to aspirin or ibuprofen taken by mouth.

**If any of the above apply to you, only use this product on advice from your doctor or pharmacist.**

**Take special care** when using this product:

- Use it only on the skin.
- Do not use it on children under 12 years old unless advised by a doctor.
- Do not apply it to broken or irritated skin.
- Keep the gel away from the eyes, nose and mouth.

#### Using other medicines

- Interaction between Ibuleve and blood pressure lowering drugs and anticoagulants (medicines that stop blood clotting) is possible, in theory, although very unlikely. If you would like more advice about this, speak to your doctor or pharmacist.
- If you are also taking aspirin or other NSAIDs by mouth, discuss this beforehand with your doctor or pharmacist because these may increase the risk of undesirable effects.
- Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines including other medicines obtained without prescription.

#### Pregnancy and breast-feeding

You should not use Ibuleve if you are pregnant or breast-feeding.

#### Driving and using machinery

Using this product is not known to affect your ability to drive or use machinery.

#### Important information about one of the ingredients of Ibuleve

The product contains a small amount of propylene glycol. This ingredient may cause skin irritation in a small number of people who are intolerant to it. If your doctor has told you that you have an intolerance to propylene glycol, please talk to your doctor before using this product.

### 3. HOW TO USE IBULEVE

#### Before using this tube of gel for the first time, open it as follows:

- Unscrew the cap, turn it upside down and then use the sharp point of the cap to pierce the membrane in the neck of the tube.

**To use the gel** (for adults, the elderly and children over 12 years old):

- Use the gel up to three times a day, or as often as advised by your doctor.

- Lightly apply the gel to intact skin over the painful area.
- Apply only enough gel to thinly cover the affected area, then massage gently until absorbed.
- Wash hands after use, unless treating them.
- Carry on using the gel in this way until your condition gets better (you may find you need to use it for a few weeks, and your doctor may want you to continue using it for longer than this).
- If your symptoms worsen or continue for more than a few weeks, discuss this with your doctor before continuing treatment.

**If the gel comes into contact with broken skin or gets into the eyes, nose or mouth**

- The product may cause irritation if it comes into contact with broken skin or gets into the eyes, nose or mouth. If this happens, rinse the affected areas with plenty of water. If rinsing one eye, take care to avoid washing product into the other eye. If irritation persists, tell your doctor or pharmacist.

**If the gel is accidentally swallowed**

- Symptoms may include headache, vomiting, drowsiness and low blood pressure.
- If you experience any of these symptoms contact a doctor or hospital straight away.

**If you forget to use this product**

Do not worry if you occasionally forget to use this product, just carry on using it when you remember. If you have any further questions on the use of this product, ask your doctor or pharmacist.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, Ibuleve can cause side effects, although not everybody gets them.

**Occasionally**, mild skin rashes, itching or irritation can sometimes occur at the site of application. If this is unacceptable, or persists, stop using the product and tell your doctor or pharmacist.

**Very rarely**, the following side effects can happen with ibuprofen, although these are extremely uncommon with products such as Ibuleve that are applied to the skin.

**If you experience any of the following, stop using Ibuleve immediately and tell your doctor:**

- **Allergic reactions** (particularly in people who have a history of asthma or allergic problems), such as:
  - unexplained runny nose and watery eyes, or, in more serious cases asthma or aggravated asthma involving breathing difficulties, wheezing or chest tightness;
  - generalised allergic skin reactions involving itch, swelling, inflammation, redness and perhaps blistering and light sensitivity;
  - other more serious generalised allergic reactions possibly involving unexplained nausea and vomiting, swollen eyes, face or tongue, difficulty swallowing, dizziness or light-headedness. Unconsciousness could perhaps occur in the most serious cases.
- **Kidney problems** (particularly in people who have a history of kidney disease), such as:
  - decreased urine volume;
  - loss of appetite / weight loss;
  - swelling to abdomen.

- **Problems with the digestive system** (particularly in people who have a history of stomach ulcers etc), such as:
  - stomach pain;
  - heartburn / indigestion.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard).

By reporting side effects you can help provide more information on the safety of this medicine.

**5. HOW TO STORE IBULEVE**

- Keep it out of the reach and sight of children.
- Always replace the cap tightly after use.
- Do not store the product above 25°C.
- Do not use after the expiry date shown on the fold of the tube and the carton. The expiry date refers to the last day of that month.
- Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

**6. FURTHER INFORMATION ABOUT IBULEVE**

**What Ibuleve contains:**

The **active ingredient** is **ibuprofen** (5% w/w). The **other ingredients** are industrial methylated spirit, carbomers, propylene glycol, diethylamine and purified water.

**What Ibuleve looks like and contents of the pack**

- The product is a clear, colourless gel that contains no fragrance.
- The gel is available in tubes containing 30g.

**The Marketing Authorisation holder is** Diomed Developments Limited, Tatmore Place, Gosmore, Hitchin, Herts, SG4 7QR, UK.

**The Manufacturer is** DDD Limited, 94 Rickmansworth Road, Watford, Herts, WD18 7JJ, UK.

**The Distributor is** DDD Limited, 94 Rickmansworth Road, Watford, Herts, WD18 7JJ, UK.

**This leaflet was last revised in** May 2015.

To listen to or request a copy of this leaflet in Braille, large print or audio, please call free of charge: 0800 198 5000 (UK only).

Please be ready to give the following information: Ibuleve 00173/0060.

This is a service provided by the Royal National Institute of Blind People (RNIB).