



Package leaflet: Information for the user

Olanzapine 2.5 mg tablets
Olanzapine 5 mg tablets
Olanzapine 7.5 mg tablets
Olanzapine 10 mg tablets
Olanzapine 15 mg tablets
Olanzapine 20 mg tablets

olanzapine

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Olanzapine is and what it is used for
2. What you need to know before you take Olanzapine
3. How to take Olanzapine
4. Possible side effects
5. How to store Olanzapine
6. Contents of the pack and other information

1. WHAT OLANZAPINE IS AND WHAT IT IS USED FOR

Olanzapine tablets contain the active substance olanzapine. Olanzapine belongs to a group of medicines called antipsychotics and is used to treat the following conditions

- Schizophrenia a disease with symptoms such as hearing, seeing or sensing things which are not there, mistaken beliefs, unusual suspiciousness, and becoming withdrawn. People with this disease may also feel depressed, anxious or tense.
- Moderate to severe manic episodes, a condition with symptoms of excitement or euphoria.

Olanzapine has been shown to prevent recurrence of these symptoms in patients with bipolar disorder whose manic episode has responded to olanzapine treatment.

2. What you need to know before you take OLANZAPINE

Do not take Olanzapine

- If you are allergic (hypersensitive) to olanzapine or any of the other ingredients of this medicine (listed in section 6). An allergic reaction may be recognised as a rash, itching, a swollen face, swollen lips or shortness of breath. If this has happened to you, tell your doctor.
- If you have been previously diagnosed with eye problems such as certain kinds of glaucoma (increased pressure in the eye).

Warnings and precautions

Talk to your doctor or pharmacist before you take Olanzapine.

- The use of Olanzapine in elderly patients with dementia is not recommended as it may have serious side effects.
- Medicines of this type may cause unusual movements mainly of the face or tongue. If this happens after you have been given Olanzapine tell your doctor.
- Very rarely, medicines of this type cause a combination of fever, faster breathing, sweating, muscle stiffness and drowsiness or sleepiness. If this happens, contact your doctor at once.
- Weight gain has been seen in patients taking Olanzapine. You and your doctor should check your weight regularly. Consider referral to a dietician or help with a diet plan if necessary.
- High blood sugar and high levels of fat (triglycerides and cholesterol) have been seen in patients taking Olanzapine. Your doctor should do blood tests to check blood sugar and certain fat levels before you start taking Olanzapine and regularly during treatment.
- Tell the doctor if you or someone else in your family has a history of blood clots, as medicines like these have been associated with the formation of blood clots.

If you suffer from any of the following illnesses tell your doctor as soon as possible:

- Diabetes
- Heart disease
- Liver or kidney disease
- Parkinson's disease
- Seizures
- Prostate problems
- A blocked intestine (Paralytic ileus)
- Blood disorders
- Stroke or "mini" stroke (temporary symptoms of stroke)
- If you know that you may have salt depletion as a result of prolonged severe diarrhoea and vomiting (being sick) or usage of diuretics (water tablets)

If you suffer from dementia, you or your carer/relative should tell your doctor if you have ever had a stroke or "mini" stroke.

As a routine precaution, if you are over 65 years your blood pressure may be monitored by your doctor.

Children and adolescents

Olanzapine is not for patients who are under 18 years.

Other medicines and Olanzapine

Only take other medicines while you are on Olanzapine if your doctor tells you that you can. You might feel drowsy if Olanzapine is taken in combination with antidepressants or medicines taken for anxiety or to help you sleep (tranquillisers).

Tell your doctor if you are taking have recently taken or might take any other medicines,

In particular, tell your doctor if you are taking:

- medicines for Parkinson's disease.
- carbamazepine (an anti-epileptic and mood stabiliser), fluvoxamine (an antidepressant) or ciprofloxacin (an antibiotic) - it may be necessary to change your Olanzapine dose.

Olanzapine with alcohol

Do not drink any alcohol if you have been given Olanzapine as together with alcohol it may make you feel drowsy.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.. You should not be given this medicine when breast-feeding, as small amounts of Olanzapine can pass into breast milk.

The following symptoms may occur in newborn babies, of mothers that have used olanzapine in the last trimester (last three months of their pregnancy): shaking, muscle stiffness and/or weakness, sleepiness, agitation, breathing problems, and difficulty in feeding. If your baby develops any of these symptoms you may need to contact your doctor.

Driving and using machines

There is a risk of feeling drowsy when you are given Olanzapine. If this happens do not drive or operate any tools or machines. Tell your doctor.

Olanzapine contains lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take OLANZAPINE

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Your doctor will tell you how many Olanzapine tablets to take and how long you should continue to take them. The daily dose of Olanzapine is between 5 mg and 20 mg. Consult your doctor if your symptoms return but do not stop taking Olanzapine unless your doctor tells you to.

You should take your Olanzapine tablets once a day following the advice of your doctor. Try to take your tablets at the same time each day. It does not matter whether you take them with or without food. Olanzapine tablets are for oral use. You should swallow the Olanzapine tablets whole with water.

If you take more Olanzapine than you should

Patients who have taken more Olanzapine than they should, have experienced the following symptoms: rapid beating of the heart, agitation/aggressiveness, problems with speech, unusual movements (especially of the face or tongue) and reduced level of consciousness. Other symptoms may be: acute confusion, seizures (epilepsy), coma, a combination of fever, faster breathing, sweating, muscle stiffness and drowsiness or sleepiness, slowing of the breathing rate, aspiration, high blood pressure or low blood pressure, abnormal rhythms of the heart. Contact your doctor or hospital straight away if you experience any of the above symptoms.. Show the doctor your pack of tablets.

If you forget to take Olanzapine

Take your tablets as soon as you remember. Do not take two doses in one day.

If you stop taking Olanzapine

Do not stop taking your tablets just because you feel better. It is important that you carry on taking Olanzapine for as long as your doctor tells you.

If you suddenly stop taking Olanzapine, symptoms such as sweating, unable to sleep, tremor, anxiety or nausea and vomiting might occur. Your doctor may suggest you to reduce the dose gradually before stopping treatment.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Tell your doctor immediately if you have:

- unusual movement (a common side effect that may affect up to 1 in 10 people) mainly of the face or tongue;
- blood clots in the veins (an uncommon side effect that may affect up to 1 in 100 people) especially in the legs (symptoms include swelling, pain, and redness in the leg), which may travel through blood vessels to the lungs causing chest pain and difficulty in breathing. If you notice any of these symptoms seek medical advice immediately;

P15XXXXXX

- a combination of fever, faster breathing, sweating, muscle stiffness and drowsiness or sleepiness (the frequency of this side effect cannot be estimated from the available data).

Very common side effects: (may affect more than 1 in 10 people) include weight gain; sleepiness; and increases in levels of prolactin in the blood

In the early stages of treatment, some people may feel dizzy or faint (with a slow heart rate), especially when getting up from a lying or sitting position. This will usually pass on its own but if it does not, tell your doctor.

Common side effects: (may affect up to 1 in 10 people) include

- changes in the levels of some blood cells, circulating fats and early in treatment, temporary increases in liver enzymes..
- increases in the level of sugars in the blood and urine.
- increases in levels of uric acid and creatine phosphokinase in the blood
- feeling more hungry.
- dizziness.
- restlessness.
- tremor.
- unusual movements (dyskinesias)
- constipation.
- dry mouth.
- rash.
- loss of strength.
- extreme tiredness.
- water retention leading to swelling of the hands, ankles or feet,
- fever
- joint pain and sexual dysfunctions such as decreased libido in males and females or erectile dysfunction in males.

Uncommon side effects: (may affect up to 1 in 100 people) include

- Hypersensitivity (e.g. swelling in the mouth and throat, itching, rash)
- Diabetes or the worsening of diabetes, occasionally associated with ketoacidosis (ketones in the blood and urine) or coma;
- Seizures, usually associated with a history of seizures (epilepsy);
- Muscle stiffness or spasms (including eye movements);
- Restless legs syndrome
- Problems with speech
- stuttering
- slow heart rate.
- sensitivity to sunlight.
- Bleeding from the nose;
- Abdominal distension;
- drooling;
- Memory loss or forgetfulness
- urinary incontinence, lack of ability to urinate.
- hair loss.
- absence or decrease in menstrual periods.
- changes in breasts in males and females such as an abnormal production of breast milk or abnormal growth.

Rare side effects (may affect up to 1 in 1000 people)

- Lowering of normal body temperature
- Abnormal rhythms of the heart
- Sudden unexplained death
- Inflammation of the pancreas causing severe Stomach pain, fever and sickness
- Liver disease appearing as yellowing of the skin and white parts of the eyes
- Muscle disease presenting as unexplained aches and pains
- Prolonged and/or painful erection.

Very rare side effects include serious allergic reactions such as Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS). DRESS appears initially as flu-like symptoms with a rash on the face and then with an extended rash, high temperature, enlarged lymph nodes, increased levels of liver enzymes seen on blood tests and an increase in a type of white blood cells (eosinophilia)

While taking olanzapine, elderly patients with dementia may suffer from stroke, pneumonia, urinary incontinence, falls, extreme tiredness, visual hallucinations, a rise in body temperature, redness of the skin and have trouble walking. Some fatal cases have been reported in this particular group of patients.

In patients with Parkinson's disease Olanzapine may worsen the symptoms.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly. via Yellow Card Scheme
Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store OLANZAPINE

Keep this medicine out of sight and reach of children.

Store below 25°C.

Do not use this medicine after the expiry date, which is stated on the outer carton, blister and bottle label.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Olanzapine contains

- The active substance is olanzapine. Each Olanzapine tablet contains either 2.5 mg, 5 mg, 7.5 mg, 10 mg, 15 mg or 20 mg of the active substance. The exact amount is shown on your Olanzapine tablet pack.
- The other ingredients are - lactose monohydrate, crospovidone (type B), hydroxypropyl cellulose and magnesium stearate.

What Olanzapine looks like and contents of the pack

Olanzapine 2.5 mg Tablets:

Yellow coloured, circular (5.2 mm in diameter), biconvex uncoated tablets, debossed with 'C' on one side and '45' on the other side

Olanzapine 5 mg Tablets:

Yellow coloured, circular (6.5 mm in diameter), biconvex uncoated tablets, debossed with 'C' on one side and '46' on the other side

Olanzapine 7.5 mg Tablets:

Yellow coloured, circular (7.5 mm in diameter), biconvex uncoated tablets, debossed with 'C' on one side and '47' on the other side

Olanzapine 10 mg Tablets:

Yellow coloured, circular (8.0 mm in diameter), biconvex uncoated tablets, debossed with 'C' on one side and '48' on the other side

Olanzapine 15 mg Tablets:

Yellow coloured, circular (9.0 mm in diameter), biconvex uncoated tablets, debossed with 'C' on one side and '49' on the other side

Olanzapine 20 mg Tablets:

Yellow coloured, circular (10.5 mm in diameter), biconvex uncoated tablets, debossed with 'C' on one side and '50' on the other side

Olanzapine tablets are available in: PVC/Polyamide/Aluminium/PVC/Aluminium blister pack:

7, 14, 28, 30, 35, 50, 56, 70, 96, 98 & 100 tablets.

HDPE bottle and polypropylene closure with silica gel desiccant:

30 & 1000 tablets

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Milpharm Limited
Ares, Odyssey Business Park
West End Road
South Ruislip HA4 6QD
United Kingdom

Manufacturer

APL Swift Services (Malta) Limited
HF 26, Hal Far Industrial Estate, Hal Far
Brizebbugia BBG 3000
Malta

or

Orion Corporation, Orion Pharma
Orionintie 1,
FI-02200 Espoo,
Finland

This medicinal product is authorised in the Member States of the EEA under the following names:

Belgium	Olanzapine AB 2,5mg/ 5mg/ 7,5mg/ 10mg/ 15mg/ 20mg tabletten
Cyprus	Olanzapine Aurobindo 5 mg/ 10 mg δισκία
Denmark	Olanzapin "Orion"
Finland	Olanzapin Orion 2,5mg/ 5mg/ 7,5mg/ 10mg/ 15mg/ 20mg tabletit
France	OLANZAPINE ARROW 5mg/ 7,5mg/ 10mg, comprimé
Germany	Olanzapin Aurobindo 2,5mg/ 5mg/ 7,5mg/ 10mg/ 15mg/ 20 mg Tabletten
Italy	Olanzapina Aurobindo
Malta	Olanzapine Aurobindo 2.5mg/ 5mg/ 7.5mg/ 10mg/ 15mg/ 20mg tablets
Netherlands	Olanzapine Aurobindo 2,5mg/ 5mg/ 10mg/ 15mg/ 20mg, tabletten
Poland	Sizin/Olanzapina Aurobindo
Portugal	Olanzapina Aurobindo
Romania	Olanzapina Aurobindo 5mg/ 10mg/ 15mg/ 20 mg comprimate
Spain	OLANZAPINA AUROBINDO 2,5mg/ 5mg/ 7,5mg/ 10mg, comprimidos EFG
Sweden	Olanzapin Orion 2,5mg/ 5mg/ 7,5mg/ 10mg/ 15mg/ 20mg tablettér
United Kingdom	Olanzapine 2.5mg/ 5mg/ 7.5mg/ 10mg/ 15mg/ 20mg tablets

This leaflet was last revised in 03/2020.

P15XXXXXX