

# Zolmitriptan 2.5mg Film-coated Tablets

## Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet:

- 1 What Zolmitriptan tablets are and what they are used for**
- 2 What you need to know before you take Zolmitriptan tablets**
- 3 How to take Zolmitriptan tablets**
- 4 Possible side effects**
- 5 How to store Zolmitriptan tablets**
- 6 Contents of pack and other information**

## 1 What Zolmitriptan tablets are and what they are used for

Zolmitriptan tablets contains zolmitriptan and belongs to a group of medicines called triptans.

Zolmitriptan is used to treat migraine headache.

Migraine symptoms may be caused by the widening of blood vessels in the head. Zolmitriptan is thought to reduce the widening of these blood vessels. This helps to take away the headache and other symptoms of a migraine attack, such as feeling or being sick (nausea or vomiting) and being sensitive to light and sound. Zolmitriptan works only when a migraine attack has started. It will not stop you from getting an attack.

## 2 What you need to know before you take Zolmitriptan tablets

### Do not take Zolmitriptan tablets if you:

- are **allergic** to zolmitriptan or any of the other ingredients of this medicine (listed in section 6)
- have **high blood pressure**
- have ever had **heart** problems, including a heart attack, angina (chest pain caused by exercise or effort), Prinzmetal's angina (chest pain which happens at rest) or have experienced heart related symptoms such as shortness of breath or pressure over the chest
- have **circulatory problems** (restricted blood flow in your legs or arms)
- have had a **stroke** or short-lasting symptoms similar to stroke (transient ischaemic attack or TIA)
- have **severe kidney** problems
- are at the same time taking some **other medicines for migraine** (e.g. ergotamine or ergot-type medicines like dihydroergotamine and methysergide) or other triptan medicines for migraine. See section below: 'Other medicines and Zolmitriptan tablets' for further information.

If you are not sure if any of these apply to you, talk to your doctor or pharmacist.

### Warnings and precautions

**Talk to your doctor or pharmacist before taking Zolmitriptan tablets if you:**

- are at risk of getting **ischaemic heart disease** (poor blood flow in the arteries of the heart). Your risk is greater if you smoke, have high blood pressure, high levels of cholesterol, diabetes or if anyone in your family has ischaemic heart disease
- have been told that you have **Wolff-Parkinson-White Syndrome** (a type of abnormal heart beat)
- have ever had **liver** problems
- have **headaches** which are not like your usual migraine headache
- are taking any other **medicine for treatment of depression** (see 'Other medicines and Zolmitriptan tablets' later in this section).

When Zolmitriptan is taken at the same time as medicines of the type SSRI or SNRI, which are used for the treatment of depression, there is a risk of development of so called serotonin syndrome. The symptoms can be severe and include shivering, over-reactive reflexes, nausea, fever, sweating, delirium, mental confusion and coma. In case you are taking this combination your doctor should monitor you carefully, especially in the beginning of treatment, when doses are increased or if another serotonergic medication is added. If you experience any of these symptoms contact a doctor as soon as possible.

As with other migraine treatments, using too much zolmitriptan can cause daily headaches or can make your migraine headaches worse. Ask your doctor if you think that this is the case for you. You may need to stop taking zolmitriptan to correct the problem.

If you go into hospital tell the medical staff you are taking Zolmitriptan.

### Children and adolescents

Zolmitriptan is not recommended for people aged under 18 years.

### Older people

Zolmitriptan is not recommended for people over 65.

### Other medicines and Zolmitriptan tablets

**Tell your doctor or pharmacist** if you are taking, have recently taken or might take any other medicines. In particular, tell your doctor if you are taking any of the following medicines:

#### Medicines for migraine

- if you take other triptans than zolmitriptan leave 24 hours before taking Zolmitriptan, and after taking Zolmitriptan leave 24 hours before taking any other triptans.
- if you take medicines containing ergotamine or ergot-type medicines (such as dihydroergotamine or methysergide), leave 24 hours before taking Zolmitriptan after taking Zolmitriptan leave 6 hours before taking ergotamine or ergot-type medicines.

#### Medicines for depression (see also section 'Warnings and precautions' above)

- moclobemide or fluvoxamine
- medicines called SSRIs (selective serotonin reuptake inhibitors)
- medicines called SNRIs (serotonin norepinephrine reuptake inhibitors) such as venlafaxine, duloxetine

#### Other medicines

- cimetidine (for indigestion or stomach ulcers)
- a quinolone antibiotic (such as ciprofloxacin)

If you are using herbal remedies containing St John's Wort (*Hypericum perforatum*), side effects of Zolmitriptan may be more likely to happen.

### Zolmitriptan tablets with food and drink

You can take Zolmitriptan with or without food. It does not affect the way that Zolmitriptan works.

### Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

It is not known if taking Zolmitriptan during pregnancy is harmful.

Do not breast-feed within 24 hours of taking Zolmitriptan.

### Driving and using machines

During a migraine attack your reactions may be slower than usual. Bear this in mind when you drive or use any tools or machines.

Zolmitriptan is unlikely to affect driving or using tools or machines. However, it is best to wait to see how Zolmitriptan affects you before you try these activities.

### Zolmitriptan tablets contain lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### 3 How to take Zolmitriptan tablets

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

You can take Zolmitriptan as soon as a migraine headache starts. You can also take it once an attack is underway. The recommended dose is one 2.5 mg tablet (2.5 mg zolmitriptan) or two 2.5 mg tablets (5 mg zolmitriptan). You can take another 2.5 mg tablet (2.5 mg zolmitriptan) or two 2.5 mg tablets (5 mg zolmitriptan) if the migraine is still present after two hours or if it returns within 24 hours.

If the tablets did not give you enough help with your migraine, tell your doctor. Your doctor may raise the dose to 5 mg zolmitriptan or change your treatment.

Do not use more than the dose prescribed for you.

Do not use more than two doses in one day. If you have been prescribed one 2.5 mg tablet (2.5 mg zolmitriptan), the maximum daily dose is two 2.5 mg tablets (5 mg zolmitriptan). If you have been prescribed two 2.5 mg tablets (5 mg zolmitriptan), the maximum daily dose is four 2.5 mg tablets (10 mg zolmitriptan). Swallow your tablets with a drink of water

#### If you take more Zolmitriptan tablets than you should

If you have taken more Zolmitriptan than prescribed by your doctor, tell your doctor or go to the nearest hospital straight away. Take the Zolmitriptan medicine with you. When too many tablets are taken the symptoms may possibly include sedation.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### 4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Some of the symptoms below could be part of the migraine attack itself.

#### If you notice any of the following serious side effects, stop taking Zolmitriptan tablets and contact a doctor straight away:

*Rare (may affect up to 1 in 1,000 people)*

- Allergic reactions including itchy rash (urticaria) and swelling of the face, lips, mouth, tongue and throat.

*Very rares (may affect up to 1 in 10,000 people)*

- Angina (pain in the chest, often brought on by exercise), heart attack or spasm of the blood vessels of the heart. The signs include chest pain and shortness of breath.
- Spasm of the blood vessels of the gut, which can cause damage to your gut. The signs include stomach pain or bloody diarrhoea.

Other possible side effects include:

*Common (may affect up to 1 in 10 people):*

- Abnormal sensations such as tingling in your fingers and toes or skin that is sensitive to touch.
- Feeling sleepy, dizzy or warm
- Headache
- Uneven heart beat
- Feeling sick, vomiting
- Stomach pain
- Dry mouth
- Difficulty swallowing
- Muscle weakness or muscle pain
- Feeling weak
- Heaviness, tightness, pain or pressure in throat, neck, arms and legs, or chest

*Uncommon (may affect up to 1 in 100 people):*

- Very fast heart beat
- Slightly higher blood pressure
- Increase in the amount of water you pass (urine) or in how often you need to pass water

*Very rare (may affect up to 1 in 10,000 people):*

- Sudden urgent need to pass water (urine).

#### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the internet at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard). Alternatively you can call Freephone 0800 100 3352 (available from 10am to 2pm Mondays to Fridays) or fill in a paper form available from your local pharmacy.

By reporting side effects you can help provide more information on the safety of this medicine.

### 5 How to store Zolmitriptan tablets

Keep out of the sight and reach of children.

Do not use Zolmitriptan after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

### 6 Contents of the pack and other information

#### What Zolmitriptan tablets contains

- The active substance is zolmitriptan. Zolmitriptan film-coated tablets contain 2.5mg of zolmitriptan.
- The other ingredients are: tablet core: lactose (anhydrous), microcrystalline cellulose, sodium starch glycolate (Type A), magnesium stearate; tablet coating: hypromellose (E464), titanium dioxide (E171), polydextrose, talc, maltodextrin, medium chain triglycerides, iron oxide yellow (E172) and iron oxide black (E172).

#### What Zolmitriptan tablets looks like and contents of the pack

Zolmitriptan 2.5mg film-coated tablets are light yellow, round, biconvex tablets with 'ZL 2.5' debossed on one side.

Zolmitriptan 2.5mg comes in blister packs containing 6 or 12 film-coated tablets.

#### Marketing Authorisation Holder

Actavis Group PTC ehf  
Reykjavikurvegur 76-78  
220 Hafnarfjörður  
Iceland

#### Manufacturer

Actavis Ltd  
BLB016 Bulebel Industrial Estate  
Zejtun ZTN3000  
Malta

This leaflet was last revised in – November 2016.