

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### Buttercup Bronchostop Cough Pastilles

Thyme herb extract

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or qualified healthcare practitioner has told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist or qualified healthcare practitioner if you need more information or advice.
- If you get any side effects, talk to your pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor or qualified healthcare practitioner if you do not feel better or if you feel worse after 7 days.

#### **What is in this leaflet:**

1. What Buttercup Bronchostop Cough Pastilles are and what they are used for
2. What you need to know before you take Buttercup Bronchostop Cough Pastilles
3. How to take Buttercup Bronchostop Cough Pastilles
4. Possible side effects
5. How to store Buttercup Bronchostop Cough Pastilles
6. Contents of the pack and other information

#### **1. WHAT BUTTERCUP BRONCHOSTOP COUGH PASTILLES ARE AND WHAT THEY ARE USED FOR**

Buttercup Bronchostop Cough Pastilles are a traditional herbal medicinal product used for the relief of coughs, such as chesty coughs and dry, tickly, irritating coughs and catarrh, based on traditional use only.

#### **2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE BUTTERCUP BRONCHOSTOP COUGH PASTILLES**

##### **Do not take Buttercup Bronchostop Cough Pastilles**

- If you are allergic to Thyme or other members of the Lamiaceae family (e.g. Basil, Rosemary, Sage), or to any of the other ingredients of this medicine (listed in section 6).
- If you are pregnant or breast-feeding.

##### **Warnings and precautions**

###### **Children**

Do not give this medicine to children under 12 years.

### **Other medicines and Buttercup Bronchostop Cough Pastilles**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

### **Pregnancy and breast-feeding**

Do not take this medicine if you are pregnant or breast-feeding.

### **Driving and using machines**

No studies on the effects on the ability to drive and to use machines have been performed. Buttercup Bronchostop Cough Pastilles contain no ingredients that are known to cause drowsiness.

### **Buttercup Bronchostop Cough Pastilles contain fructose and sorbitol**

This medicine contains 523 mg sorbitol and 300 mg fructose in each pastille, corresponding to approximately 0.1 carbohydrate unit. This should be considered in patients with diabetes mellitus. Sorbitol is a source of fructose. If your doctor has told you that you (or your child) have an intolerance to some sugars or if you have been diagnosed with hereditary fructose intolerance (HFI), a rare genetic disorder in which a person cannot break down fructose, talk to your doctor before you (or your child) take or receive this medicine.

### **Buttercup Bronchostop Cough Pastilles contain propylene-glycol (E1520)**

The medicine contains 5.53 mg of propylene-glycol in each pastille.

### **Buttercup Bronchostop Cough Pastilles contain benzyl-alcohol (E1519)**

This medicine contains 0.0018 mg benzyl-alcohol (E1519) in each pastille. Benzyl-alcohol (E1519) may cause allergic reactions.

### **Buttercup Bronchostop Cough Pastilles contain sodium**

This medicine contains less than 1 mmol sodium (23 mg) per pastille, that is to say essentially 'sodium-free'.

## **3. HOW TO TAKE BUTTERCUP BRONCHOSTOP COUGH PASTILLES**

Always take this medicine exactly as described in this leaflet or as your doctor, pharmacist or qualified healthcare practitioner has told you. Check with your doctor, pharmacist or qualified healthcare practitioner if you are not sure.

For oral use (allow to dissolve in the mouth through sucking).

### **Dosage**

#### **Adults, the elderly and children over 12 years:**

1 – 2 pastilles to be taken every 4 hours, 4 times a day. If required, up to a maximum of 12 pastilles can be taken per day.

**Do not give to children under 12 years.**

**DO NOT EXCEED THE STATED DOSE.**

You must talk to a doctor or qualified healthcare practitioner if you do not feel better or if you feel worse after 7 days.

If you are short of breath, have a high temperature (fever), your phlegm is yellow-green or brown in colour (purulent sputum), you must consult your doctor or qualified healthcare practitioner.

**If you take more Buttercup Bronchostop Cough Pastilles than you should**

Talk to a doctor straight away, or contact your nearest hospital casualty department. Take the carton and this leaflet with you.

**If you forget to take Buttercup Bronchostop Cough Pastilles**

Take the next dose when it is due. Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Allergic reactions including Quincke's oedema and anaphylactic shock have been reported.

**Stop taking the pastilles and tell your doctor immediately if:**

- you get any swelling of the face, lips tongue or throat
- you have any difficulty in swallowing or breathing (including shortness of breath)
- you get a skin rash after taking the pastilles

The above symptoms may mean you are having a very rare, serious allergic reaction to this medicine.

**Other possible side effects**

If any of the following side effects get serious or concerns you, or you notice any other side effect, stop taking the pastilles.

- Stomach disorders (*frequency unknown*)

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

**5. HOW TO STORE BUTTERCUP BRONCHOSTOP COUGH PASTILLES**

Do not store above 25°C.

Store the blister in the original package in order to protect from light.

**Keep out of the sight and reach of children.**

Do not use this medicine after the expiry date which is stated on the carton. The expiry date refers to the last day of that month.

Do not throw away medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures help protect the environment.

## 6. CONTENTS OF THE PACK AND OTHER INFORMATION

### What Buttercup Bronchostop Cough Pastilles contain

Each pastille contains the active ingredient 59.5 mg of extract (as dry extract) from Thyme herb (*Thymus vulgaris* L. and *Thymus zygis* L., herb ) (DER 7-13:1) Extraction solvent: water

Each pastille also contains 300 mg of fructose and 523 mg of sorbitol

### The other ingredients are:

Maltodextrin, Arabic gum, Fructose, Sorbitol liquid non-crystallising (E420), Citric acid anhydrous, Saccharin sodium, Aronia (chokeberry) flavour (which contains propylene-glycol (E1520)), Fruit of forest (berry) flavour (propylene-glycol (E1520) and benzyl-alcohol (E1519)), Paraffin light liquid, Beeswax white, Purified water

### What Buttercup Bronchostop Cough Pastilles look like and contents of the package

Buttercup Bronchostop Cough Pastilles are hexagonal, brown pastilles.

Buttercup Bronchostop Cough Pastilles are available in blister packs containing 10, 20 or 40 pastilles. Not all pack sizes may be marketed.

### THR Holder and Manufacturer

Kwizda Pharma GmbH, Effingergasse 21, A-1160 Vienna, Austria

Distributed by Omega Pharma Ltd., 32 Vauxhall Bridge Road, London, SW1V 2SA, UK.

THR 16467/0002



**Is this leaflet hard to see or read? Phone 01748 828860 for help.**

**This leaflet was revised in September 2020**