



LEAFLET FACE

92.0 mm	<p>4- Possible side effects If your baby experiences any side effect such as allergic reaction, after using this product, stop using it and seek medical advice. Reporting of side effects If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects, you can help provide more information on the safety of this medicine. NZ: You can also report side effects directly via https://nzphvc.otago.ac.nz/report/</p> <p>5- How to store this medicine KEEP OUT OF THE SIGHT AND REACH OF CHILDREN. Store below 25°C. Use within 14 days after first opening. Use before the expiry date shown on the front of the bottle.</p>	<p>6- Further information Woodward's Gripe Water - Alcohol Free & Sugar Free is an oral solution containing terpeneless dill seed oil 2.3mg/5mL, and sodium hydrogen carbonate 52.5mg/5mL as active ingredients and also maltitol liquid, glycerol, sodium salts of esters of p-hydroxybenzoic acid (E215, E217 and E219), disodium edetate, propylene glycol and purified water. MA Holder: Reckitt Benckiser Healthcare (UK) Ltd, 103-105 Bath Road Slough SL1 3UH Manufacturer : FAMAR A.V.E, 49th National Road Athens-Lamia, Avlona Attiki, 19011 Greece Or Reckitt Benckiser Healthcare (UK) Limited, Dansom Lane, Hull, HU8 7DS, United Kingdom Last leaflet update 09/2023</p>	<p>Read the leaflet carefully before use and for full list of ingredients. Store below 25°C. Use within 14 days of first opening the bottle.</p> <p>MA Holder: Reckitt Benckiser Healthcare (UK) Ltd, 103-105 Bath Road Slough SL1 3UH</p> <p>PL 00063/0698</p> <p>Reckitt Benckiser NZ, Auckland Toll Free Contact Number: 0800 40 30 30</p>	
	Page 8,9	Page 10,11	Page 12	Page 1
60.00 mm	60.00 mm	30.00 mm	33.00 mm	

Pharmacode Reads This Way
First Bar
Last Bar

LEAFLET REVERSE

92.0 mm	<p>Patient Information Woodward's Gripe Water - Alcohol Free & Sugar Free Terpeneless dill seed oil 2.3mg/5mL, Sodium hydrogen carbonate 52.5mg/5mL</p> <p>1- What is this medicine used for This medicine is an antacid and antiflatulent product. It is used to relieve gripping pain or waves of severe abdominal pain (infant colic) in babies, due to trapped air in the tummy. It can be given to babies from 1 to 12 months.</p> <p>2- Before you use this medicine Do not give to a baby: - who is allergic to any of the ingredients in this product (see section 6) - who has kidney problems. Important information about some of the ingredients This product is sugar free but contains maltitol, which may affect blood sugar levels. Therefore if your baby has an inherited intolerance to some sugars, check</p>	<p>with your doctor before giving this medicine. May have a mild laxative effect. Calorific value 2.3 kcal/g maltitol. This medicine contains hydroxybenzoate derivatives (E215, E217 and E219) which may cause allergic reactions (possibly delayed). This medicine contains 15.73 mg sodium (main component of cooking/table salt) in each 5ml. This is equivalent to 0.8 % of the recommended maximum daily dietary intake of sodium for an adult. This medicine contains 200 mg propylene glycol in each 5 ml. If your child is less than 5 years old, talk to your doctor or pharmacist before giving them this medicine, in particular if they use other medicines that contain propylene glycol or alcohol. If you suffer from a liver or kidney disease, do not take this medicine unless recommended by your doctor. Your doctor may carry out extra checks while you are taking this medicine.</p>	<p>Maximum Daily Dose: Sodium: 188.76 mg (8.21 mmol) Propylene glycol for child aged 6 - 12 months**: 2400 mg Propylene glycol for child aged 1 - 6 months***: 1200 mg Maltitol liquid: 11.4 g **(343 mg/kg/day based on weight of 7 kg for a 6 month old)*** (300 mg/kg/day based on weight of 4 kg for a 1 month old)</p> <p>3- How to use this medicine This medicine is only for babies from 1 to 12 months old. 1-6 months old: one 5mL spoonful 6-12 months old: two 5mL spoonfuls Give during or after a feed. Do not use more than 6 times in 24 hours. Allow two hours before administering other medicinal products, where possible. If symptoms persist, contact your doctor. If you give your baby too much of this medicine, speak to a doctor or pharmacist at once.</p>
	Page 2,3	Page 4,5	Page 6,7
63.00 mm	60.00 mm	60.00 mm	