

## Package leaflet: Information for the patient

---

### Dalacin® T Topical Lotion

clindamycin phosphate

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or your pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What Dalacin T Topical Lotion is and what it is used for
2. What you need to know before you use Dalacin T Topical Lotion
3. How to use Dalacin T Topical Lotion
4. Possible side effects
5. How to store Dalacin T Topical Lotion
6. Contents of the pack and other information

### 1. What Dalacin T Topical Lotion is and what it is used for

Dalacin T Topical Lotion contains the active substance clindamycin phosphate. It is an anti-infective used on the skin for the treatment of acne in adults.

### 2. What you need to know before you use Dalacin T Topical Lotion

#### Do not use Dalacin T Topical Lotion

- if you are allergic to clindamycin, lincomycin or any of the other ingredients of this medicine (listed in section 6).
- if you have a history of inflammatory bowel disease or antibiotic-associated colitis.

#### Warnings and precautions

Talk to your doctor or pharmacist before using Dalacin T Topical Lotion if you have diarrhoea or usually get diarrhoea when you take antibiotics.

If you develop severe or prolonged or bloody diarrhoea during or after using Dalacin T Topical Lotion **tell your doctor immediately** since it may be necessary to interrupt the treatment. This may be a sign of bowel inflammation (*pseudomembranous colitis*) which can occur following treatment with antibiotics.

Take care when using Dalacin T Topical Lotion around your mouth as it has an unpleasant taste.

#### Other medicines and Dalacin T Topical Lotion

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Tell your doctor if you are using any skin treatments that contain benzoyl peroxide. These medicines should not be used at the same time as Dalacin T Topical Lotion.

If you are going to hospital for a surgical procedure, tell your doctor and/or anaesthetist if you are taking neuromuscular blocking agents (drugs that cause paralysis of muscle).

### **Pregnancy and breast-feeding**

#### **Pregnancy**

Dalacin T Topical Lotion should be used during pregnancy only if clearly needed.

#### **Breast-feeding**

Tell your doctor if you will be breast-feeding while using Dalacin T Topical Lotion as the active substance in this medicine may be passed into breast milk. Your doctor will decide if Dalacin T Topical Lotion is suitable for you. If you continue to take Dalacin T, you should stop breast-feeding.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

### **Driving and using machines**

No effects on the ability to drive or use machines have been seen with Dalacin T Topical Lotion.

### **Dalacin T Topical Lotion contains cetostearyl alcohol and methyl parahydroxybenzoate**

Cetostearyl alcohol may cause local skin reactions (e.g. contact dermatitis). Methyl parahydroxybenzoate may cause allergic reactions (possibly delayed).

## **3. How to use Dalacin T Topical Lotion**

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Usually, you should use this medicine twice a day. Shake the bottle well before use and wash the affected area before applying the lotion. Apply a thin film of the lotion to the affected area by gently rubbing it into your skin. You may use a cotton pad for this, or you can use your fingers.

Wash and dry your hands before and after using the lotion. The Dalacin T Lotion bottle has a dispensing cap. Always replace the cap after use.

### **If you forget to apply Dalacin T Topical Lotion**

Do not worry if you miss one or two applications; just carry on as soon as you remember.

### **If you stop using Dalacin T Topical Lotion**

Do not stop using Dalacin T Topical Lotion as soon as your acne starts to get better. Ask your doctor when you should stop the treatment.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them. **Tell your doctor immediately** if you develop severe, persistent or bloody diarrhoea (which may be associated with stomach pain or fever). This is an uncommon side effect which may occur after treatment with antibiotics and can be a sign of serious bowel inflammation or pseudomembranous colitis.

**Side effects that have been seen with Dalacin T Topical Lotion are:**

**Very common: may affect more than 1 in 10 people**

- Skin irritation (may cause symptoms such as burning, itching, peeling), hives, dryness of skin
- Hives
- A disease of the sebaceous glands resulting in an oily coating, crusts, or scales on the skin (seborrhoea)

**Uncommon: may affect up to 1 in 100 people**

- Gastrointestinal disorder (may cause symptoms such as diarrhoea, and nausea)

**Frequency not known, but may also occur**

- Infection of the hair follicles in the skin (also known as folliculitis),
- Stinging of the eye,
- Pain in the eye,
- Stomach ache,
- Inflammation of the skin caused by contact with irritants or allergens

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for the MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Dalacin T Topical Lotion**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the bottle and carton after EXP. The expiry date refers to the last day of that month.

Do not store above 25°C.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

## **6. Contents of the pack and other information**

### **What Dalacin T Topical Lotion contains**

The active substance is clindamycin phosphate. Each ml of Dalacin T lotion contains 10 mg of clindamycin.

The other ingredients are

glycerol, sodium lauroyl sarcosinate, stearic acid, tegin, cetostearyl alcohol, isostearyl alcohol, methyl parahydroxybenzoate and water. See section 2

“Dalacin T Topical Lotion contains cetostearyl alcohol and methyl parahydroxybenzoate”.

**What Dalacin T Topical Lotion looks like and contents of the pack**

Dalacin T Topical Lotion is a white to off-white aqueous emulsion in a plastic bottle with a dispensing cap. Bottles contain 30 ml or 60 ml of Dalacin T Topical Lotion.

Not all pack sizes may be marketed.

**Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder:

Pfizer Limited  
Ramsgate Road  
Sandwich  
Kent  
CT13 9NJ  
United Kingdom.

**Manufacturer:**

Pfizer Service Company BVBA  
Hoge Wei  
1930 Zaventem  
Belgium.

**Company Contact Address:**

For further information on your medicine contact Medical Information at Pfizer Limited in Walton Oaks, Tadworth, Surrey. Tel: 01304 616161.

This leaflet was last revised in 02/2021.

Ref: DA 24\_1