



Package leaflet: Information for the user

HALEON

Piriton

Children's Allergy Syrup

Chlorphenamine Maleate

Read all of this leaflet carefully before you start using this medicine because it contains important information for you. Always use this medicine exactly as described in this leaflet or as your pharmacist has told you. This medicine is available without prescription, but you still need to use Piriton Children's Allergy Syrup carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse.

In this leaflet:

1. What Piriton Children's Allergy Syrup is and what it is used for
2. What you need to know before you take Piriton Children's Allergy Syrup
3. How to take Piriton Children's Allergy Syrup
4. Possible side effects
5. How to store Piriton Children's Allergy Syrup
6. Contents of the pack and other information

1. What Piriton Children's Allergy Syrup is and what it is used for

Piriton Children's Allergy Syrup is used to treat the allergic symptoms of hay fever and other allergies. The active ingredient is chlorphenamine maleate, an antihistamine which can help to relieve the symptoms of some allergies and itchy skin rashes. It can be used to treat the itchiness, redness, swelling, tenderness and irritation that can be caused by:

- hay fever and other allergies e.g. pet, house dust mite and mould spore allergies
- nettle rash and hives
- skin allergies and dermatitis
- prickly heat and heat rash
- reactions to food, food additives or medicines
- insect bites and stings
- the itchy rash of chickenpox.

2. What you need to know before you take Piriton Children's Allergy Syrup



Do not take Piriton Children's Allergy Syrup

- If you have ever had an allergic reaction to antihistamines or any of the ingredients (listed in Section 6)
- If you have taken monoamine oxidase inhibitors (MAOIs) prescribed for depression in the last two weeks
- If you are taking other medicines containing antihistamines, including products for the relief of colds and coughs.



Take special care with Piriton Children's Allergy Syrup

- Talk to your doctor before you take this syrup if you have very high blood pressure, heart disease, epilepsy, glaucoma, enlarged prostate, liver disease, kidney disease, bronchitis, asthma, bronchiectasis or chronic lung disorders (difficulty in breathing and cough that won't go away).
- Avoid drinking alcohol with this medicine.
- Do not drive or operate machinery if the syrup makes you feel drowsy, dizzy or if you experience blurred vision.
- Piriton Children's Allergy Syrup contains 6.3% v/v ethanol (alcohol), similar to 6.3 ml beer or 2.6 ml wine per 5 ml spoonful. This could be harmful if you suffer from alcoholism. This should also be considered if you are pregnant or breast feeding, in children and high-risk groups such as patients with liver disease or epilepsy.
- Clean your teeth regularly. The syrup contains sucrose and long term use could increase the risk of tooth decay.
- If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before using Piriton Children's Allergy Syrup.
- Piriton Children's Allergy Syrup contains 2.36 g of liquid sucrose per 5 ml. This should be taken into account in patients with diabetes mellitus.



If you are taking other medicines

Talk to your doctor or pharmacist before taking this syrup if you are taking any prescribed medicines; particularly phenytoin (for epilepsy) or medicines for anxiety or to help you sleep.



Pregnancy and breast feeding

Talk to your doctor before taking the Piriton Children's Allergy Syrup if you are pregnant or breast feeding.

1





3. How to take Piriton Children's Allergy Syrup



Use the spoon provided

Age	Dosage	Maximum in 24 hours
1-2 years	One 2.5 ml spoonful twice daily	Two 2.5 ml spoonfuls (5 ml)
2-6 years	One 2.5 ml spoonful every 4-6 hours	Six 2.5 ml spoonfuls (15 ml)
6-12 years	One 5 ml spoonful every 4-6 hours	Six 5 ml spoonfuls (30 ml)
Adults and children over 12 years	Two 5 ml spoonfuls (10 ml) every 4-6 hours	Twelve 5 ml spoonfuls (60 ml)
Elderly	Two 5 ml spoonfuls (10 ml) every 4-6 hours	Six 5 ml spoonfuls (30 ml)

Elderly: Talk to your doctor or pharmacist before you take this syrup as you may be more likely to get side effects including confusion and you may need to take a lower daily dose.



- Do not take more than the recommended dose.
- The minimum time interval between doses should be 4 hours.
- Do not use continuously for more than 14 days without consulting a doctor.
- Do not give to children under 12 months.

If you or a child take too much syrup

Contact your doctor or casualty department. Do not drive if you have taken too much.

If you forget to take the syrup

Take a dose as soon as you remember, unless it is nearly time to take the next one. Never take two doses together.

If your symptoms persist, see your doctor.

4. Possible side effects

Like all medicines, Piriton Children's Allergy Syrup can have side effects, but not everybody gets them. Children and older people are more prone to side effects.

Stop using this product and consult your doctor if you experience:

Allergic reaction which can be severe. Signs include:

- Raised and itchy rash (hives)
- Swelling, sometimes of the face or mouth causing difficulty in breathing
- Collapse

These reactions are rare.

Possible side effects:

Very Common (may affect more than 1 in 10 people)

- drowsiness and sedation

Common (may affect up to 1 in 10 people)

- disturbance in concentrating, un-coordination, dizziness, headache
- blurred vision
- feeling or being sick, dry mouth
- fatigue

Other side effects may include: allergic reactions, swelling of the skin, loss of appetite, confusion, excitation, irritability, nightmares, low mood, low blood pressure, thickening of phlegm, vomiting, abdominal pain, diarrhoea, indigestion or upset stomach, itchy skin, rash, sensitivity to the sun, muscle twitching, muscle weakness, difficulty passing water, chest tightness, ringing in the ears, changes in heart rhythm, low blood count, bruising, hepatitis (severe stomach pain, nausea or vomiting and loss of appetite and/or jaundice (yellow colouring of the skin or the whites of the eyes)).

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Piriton Children's Allergy Syrup

Keep out of the sight and reach of children.

Do not use this medicine after the 'EXP' date shown on the pack.

Store below 25°C. Protect from light.

6. Contents of the pack and other information

Active ingredient Each 5 ml dose contains Chlorphenamine Maleate 2 mg.

Other ingredients Liquid sucrose, glycerol, ethanol, tingle flavour, peppermint oil, water and, as a preservative, a mixture of methyl, ethyl and propyl hydroxybenzoates (E 218, E 214 and E 216).

Methyl, ethyl and propyl hydroxybenzoates (E 218, E 214 and E 216) may cause allergic reactions (possibly delayed).

The bottle contains 150 ml.

The marketing authorisation holder is Haleon UK Trading Limited, The Heights, Weybridge, KT13 0NY, U.K. and all enquiries should be sent to this address.

The manufacturer is Omega Pharma Manufacturing, GmbH & Co. KG, Benzstr. 25, 71083 Herrenberg, Germany.

This leaflet was last revised in September 2023.

Trade Marks are owned by or licensed to the Haleon group of companies.

