

PACKAGE LEAFLET - INFORMATION FOR THE USER

CO-DYDRAMOL 10mg/500mg TABLETS
Dihydrocodeine tartrate and Paracetamol

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

This medicine contains dihydrocodeine which is an opioid, which can cause addiction. You can get withdrawal symptoms if you stop taking it suddenly.

Keep this leaflet. You may need to read it again. If you have further questions, please ask your doctor or pharmacist. This medicine has been prescribed for you only. Do not pass this medicine on to others. It may harm them, even if their symptoms are the same as yours. If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

What is in this leaflet:

- | | |
|--|---|
| 1. What co-dydramol is and what it is used for | 4. Possible side effects |
| 2. What you need to know before you take co-dydramol | 5. How to store co-dydramol |
| 3. How to take co-dydramol | 6. Contents of the pack and other information |

1. WHAT CO-DYDRAMOL IS AND WHAT IT IS USED FOR

The name of your medicine is Co-dydramol Tablets (called co-dydramol throughout this leaflet). This medicine has been prescribed for you for the relief of mild to moderate pain. Co-dydramol contains two different medicines called paracetamol and dihydrocodeine. Dihydrocodeine belongs to a class of medicines called opioids, which are 'pain relievers'. This medicine has been prescribed to you and should not be given to anyone else. Opioids can cause addiction and you may get withdrawal symptoms if you stop taking it suddenly. Your prescriber should have explained how long you will be taking it for and when it is appropriate to stop, how to do this safely.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE CO-DYDRAMOL**Important things you should know about co-dydramol**

- Do not take for longer than your doctor tells you to.
- This medicine contains paracetamol. Do not take anything else containing paracetamol while taking this medicine.
- Taking a painkiller for headaches too often or for too long can make them worse.

Taking this medicine regularly, particularly for a long time, can lead to addiction. Your prescriber should have explained how long you will be taking it for and when it is appropriate to stop, how to do this safely.

Rarely, increasing the dose of this medicine can make you more sensitive to pain. If this happens, you need to speak to your prescriber about your treatment.

Addiction can cause withdrawal symptoms when you stop taking this medicine. Withdrawal symptoms can include restlessness, difficulty sleeping, irritability, agitation, anxiety, feeling your heartbeat (palpitations), increased blood pressure, feeling or being sick, diarrhoea, loss of appetite, shaking, shivering or sweating. Your prescriber will discuss with you how to gradually reduce your dose before stopping the medicine. It is important that you do not stop taking the medicine suddenly as you will be more likely to experience withdrawal symptoms.

Opioids should only be used by those they are prescribed for. Do not give your medicine to anyone else. Taking higher doses or more frequent doses of opioid, may increase the risk of addiction.

Overuse and misuse can lead to overdose and/or death.

Warnings and precautions**Do not take co-dydramol and tell your doctor if:**

- You are taking medicines to treat depression called MAOIs (monoamine oxidase inhibitors) or have taken them in the last 2 weeks. MAOIs are medicines such as moclobemide, phenelzine, tranylcypromine.
- You are allergic to the active substances or any of the other ingredients in your medicine (listed in Section 6). Signs of an allergic reaction include a rash and breathing problems. There can also be swelling of the legs, arms, face, throat or tongue.
- You have breathing problems.
- You are having an asthma attack.
- You have liver disease.
- You are an alcoholic.

Do not take co-dydramol if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking co-dydramol.

Children

Co-dydramol must not be given to children under 12 years of age.

Take special care and check with your doctor before taking co-dydramol if:

- You are or have ever been addicted to opioids, alcohol, prescription medicines, or illegal drugs.
- You have previously suffered from withdrawal symptoms such as agitation, anxiety, shaking or sweating when you have stopped taking alcohol or drugs.
- You feel you need to take more co-dydramol to get the same level of pain relief, this may mean you are becoming addicted to it. Speak to your prescriber who will discuss your treatment and may change your dose or switch you to an alternative pain reliever.
- You have kidney problems.
- You have asthma.
- You have problems with your thyroid gland (hypothyroidism).
- You are in the last three months of pregnancy (see Pregnancy and breast-feeding section).

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking this medicine.

Other medicines and co-dydramol

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes medicines obtained without a prescription, including herbal medicines. This is because co-dydramol and some other medicines can affect the way each other work.

Concomitant use of co-dydramol and sedative medicines such as benzodiazepines or related drugs increases the risk of drowsiness, difficulties in breathing (respiratory depression), coma and may be life-threatening. Because of this, concomitant use should only be considered when other treatment options are not possible.

However if your doctor does prescribe co-dydramol together with sedative medicines the dose and duration of concomitant treatment should be limited by your doctor.

Please tell your doctor about all sedative medicines you are taking, and follow your doctor's dose recommendation closely. It could be helpful to inform friends or relatives to be aware of the signs and symptoms stated above. Contact your doctor when experiencing such symptoms.

Tell your doctor or pharmacist if you are taking any of the following medicines, or have taken them in the past two weeks:

- Medicines which make you drowsy or sleepy (CNS depressants or a benzodiazepine) such as sleeping tablets, sedatives, tranquilisers, hypnotics and medicines used to treat anxiety or anaesthetics.
- Antidepressant medicines such as imipramine, amitriptyline, tranylcypromine, dosulepin, mirtazapine or chlorpromazine.

Tell your doctor or pharmacist if you are taking:

- Flucloxacillin (an antibiotic), due to a serious risk of blood and fluid abnormality (high anion gap metabolic acidosis) that must have urgent treatment and which may occur particularly in case of severe renal impairment, sepsis (when bacteria and their toxins circulate in the blood leading to organ damage), malnutrition, chronic alcoholism, and if the maximum daily doses of paracetamol are used.

Tell your doctor or pharmacist if you are taking any other medicines:

- Medicines used to thin the blood such as warfarin.
- Chloramphenicol - an antibiotic used for infections.

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking co-dydramol.

Co-dydramol with alcohol

You should not drink alcohol while you are taking these tablets.

Pregnancy and breast-feeding

- Do not take co-dydramol if you are pregnant or think you might be pregnant unless you have discussed this with your prescriber and the benefits of treatment are considered to outweigh the potential harm to the baby. If necessary, co-dydramol can be used during pregnancy. You should use the lowest possible dose that reduces your pain and/or your fever and use it for the shortest time possible. Contact your doctor or midwife if the pain and/or fever are not reduced or if you need to take the medicine more often.
- If you use co-dydramol during pregnancy, your baby may become dependent and experience withdrawal symptoms after the birth which may need to be treated.
- Do not take co-dydramol while you are breastfeeding as dihydrocodeine passes into breast milk and will affect your baby.

Driving and using machines

Co-dydramol can affect your ability to drive as it may make you sleepy or dizzy.

- Do not drive whilst taking this medicine until you know how it affects you

- It is an offence to drive if this medicine affects your ability to drive.
 - However, you would not be committing an offence if:
 - The medicine has been prescribed to treat a medical or dental problem and
 - You have taken it according to the instructions given by the prescriber or in the information provided with the medicine and
 - It was not affecting your ability to drive safely.
- Talk to your doctor or pharmacist if you are not sure whether it is safe for you to drive whilst taking this medicine.

3. HOW TO TAKE CO-DYDRAMOL

Your prescriber should have discussed with you, how long the course of tablets will last. They will arrange a plan for stopping treatment. This will outline how to gradually reduce the dose and stop taking the medicine.

Always take co-dydramol exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

- Do not take more than the recommended dose.
- Do not take for longer than your doctor tells you to.
- Swallow the tablets whole with a glass of water.

Adults: 1 to 2 tablets every 4 to 6 hours when necessary up to a maximum of 8 tablets in 24 hours. Elderly people may be prescribed a lower dose.

Use in children aged 16-18 years: 1 to 2 tablets every 6 hours when necessary up to a maximum of 8 tablets in 24 hours.

Use in children aged 12 to 15 years: 1 tablet every 4 to 6 hours when necessary to a maximum of 4 tablets in 24 hours.

Co-dydramol should not be given to children under 12 years of age.

If you take more co-dydramol than you should

- Talk to a doctor at once if you take too much of this medicine even if you feel well. This is because too much paracetamol can cause delayed, serious liver damage.
- Remember to take any remaining tablets and the pack with you. This is so the doctor knows what you have taken.
- Symptoms of an overdose include feeling or being sick, loss of appetite, stomach pain or liver damage, coma, clammy skin, fits, confusion, drowsiness, tiredness, low blood pressure, pinpoint pupils, slow heartbeat or breathing rate.

If you forget to take co-dydramol

If you forget to take a dose at the right time, take it as soon as you remember. However do not take a double dose to make up for a forgotten dose. Remember to leave at least 4 hours between doses.

If you stop taking co-dydramol

Do not suddenly stop taking this medicine. If you want to stop taking this medicine, discuss this with your prescriber first. They will tell you how to do this, usually by reducing the dose gradually so that any unpleasant withdrawal effects are kept to a minimum. Withdrawal symptoms such as restlessness, difficulty sleeping, irritability, agitation, anxiety, feeling your heartbeat (palpitations), increased blood pressure, feeling or being sick, diarrhoea, shaking, shivering or sweating may occur if you suddenly stop taking this medicine.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them. The following side effects may happen with this medicine:

Important side-effects you should know about co-dydramol

- Taking a painkiller for headaches too often or for too long can make them worse.

Unknown frequency: dependence and addiction (see section "How do I know if I am addicted?").

Stop taking co-dydramol and see a doctor or go to a hospital straight away if:

- You get swelling of the hands, feet, ankles, face, lips or throat which may cause difficulty in swallowing or breathing. You could also notice an itchy, lumpy rash (hives) or nettle rash (urticaria). This may mean you are having an **allergic reaction** to co-dydramol.
- You get serious skin reactions. Very rare cases have been reported.

Talk to your doctor straight away if you notice any of the following serious side effects:

- Severe stomach pain, which may reach through to your back. This could be a sign of inflammation of the pancreas (pancreatitis). This is a very rare side effect.

Tell your doctor or pharmacist if any of the following side effects gets serious or lasts longer than a few days:

- Constipation, feeling sick (nausea), being sick (vomiting)
- Tummy (abdominal) pain
- Headache, dizziness, light-headedness, drowsiness, confusion
- Becoming dependent on dihydrocodeine
- You get infections or bruise more easily than usual. This could be because of a blood problem.
- Redness or blood spots on the skin (purpura)

Drug Withdrawal

When you stop taking co-dydramol, you may experience drug withdrawal symptoms, which include restlessness, difficulty sleeping, irritability, agitation, anxiety, feeling your heartbeat (palpitations), increased blood pressure, feeling or being sick, diarrhoea, shaking, shivering or sweating.

How do I know if I am addicted?

If you notice any of the following signs whilst taking co-dydramol, it could be a sign that you have become addicted.

- You need to take the medicine for longer than advised by your prescriber.
- You feel you need to use more than the recommended dose.
- You are using the medicine for reasons other than prescribed.
- When you stop taking the medicine you feel unwell, and you feel better once taking the medicine again.

If you notice any of these signs, it is important you talk to your prescriber.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. HOW TO STORE CO-DYDRAMOL

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date shown on the pack. The expiry date refers to the last day of that month.

Store your medicine in the original packaging in order to protect from moisture.

Do not throw away medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION**What Co-dydramol Tablets contain**

- The active substances are 500mg of paracetamol and 10mg of dihydrocodeine tartrate. The other ingredients are pregelatinised starch, maize starch, povidone, potassium sorbate, microcrystalline cellulose, stearic acid, purified talc and magnesium stearate.

What Co-dydramol Tablets look like and contents of the pack

Flat bevelled edge white tablets, marked S/8 on one side. They come in PVC blister strips.

Pack sizes of 30, 60 and 100 tablets.

Not all pack sizes may be marketed.

The Marketing Authorisation Holder is:

Zentiva Pharma UK Limited, 12 New Fetter Lane, London, EC4A 1JP, UK

The Manufacturer is:

Zentiva Pharma UK Limited, Office 136-152, Austen House, Station View Units A-J, Guildford, GU1 4AR, UK or

Zentiva Pharma UK Limited, First Floor, Andrews House, College Road, Guildford, GU1 4QB, United Kingdom

This leaflet was updated in January 2023

