

#### 4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them. The following side effects may happen with this medicine:

**Stop taking paracetamol and see a doctor or go to a hospital straight away if:**

- You get swelling of the hands, feet, ankles, face, lips or throat which may cause difficulty in swallowing or breathing. You could also notice an itchy, lumpy rash (hives) or nettle rash (urticaria)

This may mean you are having an **allergic reaction** to paracetamol

- You get serious skin reactions. Very rare cases have been reported.

**Tell your doctor or pharmacist if any of the following side effects gets serious or lasts longer than a few days:**

- You get infections or bruise more easily than usual. This could be because of a blood problem (such as agranulocytosis, neutropenia or thrombocytopenia). This side effect has only happened in a few people taking paracetamol.

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

#### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

#### 5. HOW TO STORE PARACETAMOL

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date shown on the pack. The expiry date refers to the last day of that month.
- Store your medicine in the original packaging in order to protect from moisture.
- Do not store above 25°C.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

#### 6. CONTENTS OF THE PACK AND OTHER INFORMATION

##### What Paracetamol 500mg Soluble Tablets contain

- The active substance is paracetamol. Each tablet contains 500mg of paracetamol.
- The other ingredients are sorbitol, saccharin sodium, sodium lauryl sulphate, citric acid, sodium carbonate, sodium bicarbonate, povidone, and dimeticone.

##### Contents of pack

Paracetamol 500mg Soluble Tablets come in cartons of 12, 16, 24, 32, 48, 60 and 100. Not all pack sizes may be sold.

##### The Marketing Authorisation Holder is

Zentiva Pharma UK Limited, 12 New Fetter Lane, London, EC4A 1JP, United Kingdom

##### The Manufacturer is

A. Nattermann & Cie. GmbH, Nattermannallee 1, 50829 Cologne, Germany

This leaflet was last updated in April 2021

'Zentiva' is a registered trademark. © 2021 Zentiva

#### PACKAGE LEAFLET – INFORMATION FOR THE USER

## PARACETAMOL 500MG SOLUBLE TABLETS

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again
- If you have any further questions, ask your doctor or pharmacist
- Do not pass this medicine on to others. It may harm them, even if their signs of illness are the same as yours
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

#### What is in this leaflet:

1. What paracetamol is and what it is used for
2. What you need to know before you take paracetamol
3. How to take paracetamol
4. Possible side effects
5. How to store paracetamol
6. Contents of the pack and other information

#### 1. WHAT PARACETAMOL IS AND WHAT IT IS USED FOR

The name of your medicine is Paracetamol 500mg Soluble Tablets (called paracetamol throughout this leaflet). This medicine contains paracetamol. It belongs to a group of medicines called analgesics (painkillers) and is used to treat pain (including headache, toothache and period pain) and cold or flu symptoms.

#### 2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE PARACETAMOL

##### Do not take paracetamol and tell your doctor if:

- You are allergic (hypersensitive) to the active substance or any of the other ingredients in your medicine (listed in Section 6)  
Signs of an allergic reaction include a rash and breathing problems. There can also be swelling of the legs, arms, face, throat or tongue

##### Children

Paracetamol 500mg Soluble Tablets must not be given to children under 10 years of age

Do not take paracetamol if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking paracetamol.

##### Warnings and precautions

**Take special care and check with your doctor before taking paracetamol if:**

- You have severe kidney or liver problems
- You have a liver problem caused by alcohol

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking this medicine.

##### Other medicines and paracetamol

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes medicines obtained without a prescription, including herbal medicines. This is because paracetamol can affect the way some other medicines work. Also, some other medicines can affect the way paracetamol works.

##### While taking paracetamol you should not take any other medicines which contain paracetamol.

This includes some painkillers, cough and cold remedies. It also includes a wide range of other medicines available from your doctor and more widely in shops.

**Tell your doctor if you are taking any of the following medicines:**

- Medicines used to thin the blood such as warfarin
- Metoclopramide or domperidone - used to stop you feeling sick (nausea) or being sick (vomiting)
- Colestyramine - for lowering blood cholesterol levels

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking paracetamol.

**Paracetamol with alcohol**

You should not drink alcohol while you are taking these tablets. Taking alcohol with paracetamol can increase your chances of getting side effects.

**Pregnancy and breast-feeding**

Talk to your doctor before taking these tablets if:

- You are pregnant, think you may be pregnant or plan to get pregnant
- You are breast-feeding or planning to breast-feed

If necessary, paracetamol can be used during pregnancy. You should use the lowest possible dose that reduces your pain and/or your fever and use it for the shortest time possible. Contact your doctor or midwife if the pain and/or fever are not reduced or if you need to take the medicine more often.

**Paracetamol tablets contain:**

- **Sodium:** This medicine contains 388 mg sodium (main component of cooking/table salt) in each soluble tablet. This is equivalent to 19.4 % of the recommend maximum daily dietary intake of sodium for an adult.
- **Sorbitol:** This medicine contains 50 mg sorbitol in each soluble tablet. Sorbitol is a source of fructose. If your doctor has told you that you (or your child) have an intolerance to some sugars or if you have been diagnosed with hereditary fructose intolerance (HFI), a rare genetic disorder in which a person cannot break down fructose, talk to you doctor before you (or your child) take or receive this medicine.

**3. HOW TO TAKE PARACETAMOL**

Always take paracetamol exactly as it says in this leaflet. Check with your doctor or pharmacist if you are not sure.

- **Do not take more than the recommended dose**
- **If you need to use this medicine for more than three days at a time, see your doctor**

**Adults and children over 16**

- The usual dose of paracetamol is 2 effervescent tablets
- Dissolve the effervescent tablets in a full glass of water before taking
- Wait at least 4 hours before taking another dose
- Do not take more than 4 doses in any 24-hour period

**Use in children**

Paracetamol 500mg Soluble Tablets should not be given to children under 10 years of age.

**Children aged 10 to 15 years**

Take one tablet every four to six hours when necessary to a maximum of four doses in 24 hours.

**If you take more paracetamol than you should**

- Talk to a doctor at once if you take too much of this medicine even if you feel well. This is because too much paracetamol can cause delayed, serious liver damage.
- Remember to take any remaining tablets and the pack with you. This is so the doctor knows what you have taken

**If you forget to take paracetamol**

- If you forget to take a dose at the right time, take it as soon as you remember. However, do not take a double dose to make up for a forgotten dose. Remember to leave at least 4 hours between doses.