

Package leaflet: Information for the user

Oxycodone Hydrochloride 5mg/5ml Oral Solution
(referred to as Oxycodone 5mg/5ml Solution in this leaflet)

This medicine contains oxycodone which is an opioid, which can cause addiction. You can get withdrawal symptoms if you stop taking it suddenly.

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, please ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Oxycodone 5mg/5ml Solution is and what it is used for
2. What you need to know before you take Oxycodone 5mg/5ml Solution
3. How to take Oxycodone 5mg/5ml Solution
4. Possible side effects
5. How to store Oxycodone 5mg/5ml Solution
6. Contents of the pack and other information

1. What Oxycodone 5mg/5ml Solution is and what it is used for

This medicine has been prescribed for you for the relief of moderate to severe pain. It contains oxycodone which belongs to a group of medicines called opioids, which are ‘pain relievers’. This medicine has been prescribed for you and should not be given to anyone else. Opioids can cause addiction and you may get withdrawal symptoms if you stop taking it suddenly. Your prescriber should have explained how long you will be taking it for and when it is appropriate to stop, how to do this safely.

2. What you need to know before you take Oxycodone 5mg/5ml Solution

Do not take Oxycodone 5mg/5ml Solution if you:

- are allergic (hypersensitive) to oxycodone, or any of the other ingredients of the medicine (listed in section 6)
- have breathing problems, such as severe chronic obstructive lung disease, severe bronchial asthma or severe respiratory depression. Your doctor will have told you if you have any of these conditions. Symptoms may include breathlessness, coughing or breathing more slowly or weakly than expected
- have a condition where the small bowel does not work properly (paralytic ileus), your stomach empties more slowly than it should (delayed gastric emptying) or you have severe pain in your abdomen
- have a heart problem after long-term lung disease (cor pulmonale)
- have increased carbon dioxide levels in the blood. Symptoms may include dizziness, drowsiness, fatigue, shortness of breath and headache
- have moderate to severe liver problems. If you have other long-term liver problems you should only take this medicine if recommended by your doctor
- have ongoing problems with constipation
- are under 18 years of age.

Warnings and precautions

Talk to your doctor before using Oxycodone 5mg/5ml Solution if you:

- or anyone in your family are or have ever been addicted to opioids, alcohol, prescription medicines or illegal drugs
- have previously suffered from withdrawal symptoms such as agitation, anxiety, shaking or sweating, when you have stopped taking alcohol or drugs
- feel you need to take more Oxycodone 5mg/5ml Solution to get the same level of pain relief, this may mean you are becoming tolerant to the effects of this medicine or are becoming addicted to it. Speak to your prescriber who will discuss your treatment and may change your dose or switch you to an alternative pain reliever
- are elderly or weakened
- have an under-active thyroid gland (hypothyroidism), as you may need a lower dose
- have myxoedema (a thyroid disorder with dryness, coldness and swelling (‘puffiness’) of the skin affecting the face and limbs
- know you are suffering from a brain injury or tumour, or you have a head injury, severe headache or feel sick as this may indicate that the pressure in your skull is increased
- have low blood pressure (hypotension)
- have low blood volume (hypovolaemia); this can happen with severe external or internal bleeding, severe burns, excessive sweating, severe diarrhoea or vomiting
- feel very lightheaded or faint
- have a mental disorder as a result of an infection (toxic psychosis)
- have inflammation of the pancreas (which causes severe pain in the abdomen and back)
- have problems with your gall bladder or bile duct
- have inflammatory bowel disease
- have an enlarged prostate gland, which causes difficulty in passing urine (in men)
- have poor adrenal gland function (your adrenal gland is not working properly which may cause symptoms including weakness, weight loss, dizziness, feeling or being sick), e.g. Addison’s disease
- have severely impaired lung function. Symptoms may include breathlessness and coughing
- have long term pain unrelated to cancer
- have a condition where your breathing stops for short periods whilst you are asleep, known as sleep apnoea
- have kidney or liver problems.

Taking this medicine regularly, particularly for a long time, can lead to addiction and may result in life-threatening overdose. If you have concern that you may become dependent on Oxycodone 5mg/5ml Solution, it is important that you consult your doctor. Your doctor should have explained how long you will be using it for and when it is appropriate to stop, how to do this safely.

Rarely, increasing the dose of this medicine can make you more sensitive to pain. If this happens, you need to speak to your doctor about your treatment.

Addiction can cause withdrawal symptoms when you stop taking this medicine. Withdrawal symptoms can include restlessness, difficulty sleeping, irritability, agitation, anxiety, feeling your heartbeat (palpitations), increased blood pressure, feeling or being sick, diarrhoea, loss of appetite, shaking, shivering or sweating. Your doctor will discuss with you how to gradually reduce your dose before stopping the medicine. It is important that you do not stop taking the medicine suddenly as you will be more likely to experience withdrawal symptoms.

Opioids should only be used by those they are prescribed for. Do not give your medicine to anyone else. Taking higher doses or more frequent doses of opioid may increase the risk of addiction. Overuse and misuse can lead to overdose and/or death.

Tolerance, dependence and addiction

This medicine contains oxycodone, which is an opioid. It can cause dependence and/or addiction.

This medicine contains oxycodone which is an opioid medicine. Repeated use of opioid painkillers can result in the drug being less effective (you become accustomed to it, known as tolerance).

Repeated use of Oxycodone 5mg/5ml Solution can also lead to dependence abuse, and addiction which may result in life-threatening overdose. The risk of these side effects can increase with a higher dose and longer duration of use.

If you have concerns that you may become dependent on Oxycodone 5mg/5ml Solution it is important that you consult your doctor. Your doctor should have explained how long you will be using it for and when it is appropriate to stop, how to do this safely.

Dependence or addiction can make you feel that you are no longer in control of how much medicine you need to take or how often you

need to take it. You might feel that you need to carry on taking your medicine, even when it doesn’t help to relieve your pain.

Rarely, increasing the dose of this medicine can make you more sensitive to pain. If this happens, you need to speak to your doctor about your treatment.

The risk of becoming dependent or addicted varies from person to person. You may have a greater risk of becoming dependent or addicted on Oxycodone if:

- You or anyone in your family have ever abused or been dependent on alcohol, prescription medicines or illegal drugs (“addiction”).
 - You are a smoker.
 - You have ever had problems with your mood (depression, anxiety or a personality disorder) or have been treated by a psychiatrist for other mental illnesses.
- If you notice any of the following signs whilst taking Oxycodone, it could be a sign that you have become dependent or addicted.
- You need to take the medicine for longer than advised by your doctor
 - You need to take more than the recommended dose
 - You are using the medicine for reasons other than prescribed, for instance, ‘to stay calm’ or ‘help you sleep’
 - You have made repeated, unsuccessful attempts to quit or control the use of the medicine
 - When you stop taking the medicine you feel unwell, and you feel better once taking the medicine again (‘withdrawal effects’).

If you notice any of these signs, it is important to talk to your doctor.

Sleep-related breathing disorders

Oxycodone 5mg/5ml Solution can cause sleep-related breathing disorders such as sleep apnoea (breathing pauses during sleep) and sleep related hypoxemia (low oxygen level in the blood). The symptoms can include breathing pauses during sleep, night awakening due to shortness of breath, difficulties to maintain sleep or excessive drowsiness during the day. If you or another person observe these symptoms, contact your doctor. A dose reduction may be considered by your doctor.

If you are going to have an operation, please tell the doctor at the hospital that you are taking this medicine.

You may experience hormonal changes while taking this medicine. Your doctor may want to monitor these changes.

Withdrawal

Addiction can cause withdrawal symptoms when you stop taking this medicine. Withdrawal symptoms can include restlessness, difficulty sleeping, irritability, agitation, anxiety, feeling your heartbeat (palpitations), increased blood pressure, feeling or being sick, diarrhoea, loss of appetite, shaking, shivering or sweating. If you notice any of these signs, speak to your doctor to discuss the best treatment pathway for you, including when it is appropriate to stop and how to stop safely (See section 3, If you stop taking Oxycodone 5mg/5ml Solution).

It is important that you do not stop taking the medicine suddenly as you will be more likely to experience withdrawal symptoms.

Opioids should only be used by those they are prescribed for. Do not give your medicine to anyone else. Taking higher doses or more frequent doses of opioid may increase the risk of addiction. Overuse and misuse can lead to overdose and/or death.

Do not inject Oxycodone 5mg/5ml Solution. This can cause serious side effects including tissue death at the site of injection, infection, inflammation of the lungs and damage to the heart which may be fatal.

Other medicines and Oxycodone 5mg/5ml Solution

Taking Oxycodone 5mg/5ml Solution at the same time as other medicines that slow down the central nervous system can cause slow or difficulty breathing (respiratory depression), severe sleepiness, loss of consciousness and death. These medicines include:

- other medicines used to treat pain known as opioids (such as codeine or morphine)
- medicines used to treat epilepsy (gabapentinoids) such as pregabalin
- medicines used to treat anxiety
- medicines used to make you feel sleepy (such as benzodiazepines)
- medicines used to treat psychiatric or mental disorders (such as phenothiazines)
- anaesthetics
- muscle relaxants
- medicines used to treat high blood pressure
- a type of medicine used to treat depression known as monoamine oxidase inhibitors (MAOIs), such as tranylcypromine, phenelzine and isocarboxazid. You should not take Oxycodone 5mg/5ml Solution if you are currently taking this type of medicine, or have taken this medicine in the last two weeks.

Because of this, your doctor will only prescribe Oxycodone 5mg/5ml Solution where there are no other treatment options, and only in small doses for short periods of time. If you or your friends, family or caregivers notice that you are having difficulty breathing or that you have become very sleepy or lost consciousness you (or they) should inform your doctor **immediately**.

Taking Oxycodone 5mg/ml Solution with medicines used to treat depression known as Selective Serotonin Re-uptake Inhibitors (SSRIs) or Serotonin Norepinephrine Re-uptake Inhibitors (SNRIs) can cause a condition known as serotonin toxicity. The symptoms of this include agitation, seeing or hearing things that aren’t real (hallucinations), loss of consciousness, a fast heartbeat, blood pressure changes, increased body temperature, muscle twitching, lack of coordination, stiffness, feeling or being sick, or diarrhoea. If you are taking SSRI or SNRI medicines such as citalopram, duloxetine, escitalopram, fluoxetine, fluvoxamine, paroxetine, sertraline or venlafaxine your doctor may reduce your dose of Oxycodone 5mg/5ml Solution.

Tell your doctor or pharmacist if you are taking:

- a type of medicine used to treat depression known as tricyclic antidepressants, such as amitriptyline, clomipramine, imipramine, lofepramine or nortriptyline
- medicines used to treat allergies, such as cetirizine, fexofenadine or chlorphenamine
- medicines to help you sleep or stay calm (for example hypnotics or sedatives, including benzodiazepines)
- medicines to treat depression (such as paroxetine)
- medicines to treat psychiatric or mental disorders (such as phenothiazines or neuroleptic drugs)
- other strong analgesics (‘painkillers’)
- muscle relaxants
- medicines to treat high blood pressure
- quinidine (a medicine to treat an irregular heart beat)
- cimetidine (a medicine for stomach ulcers, indigestion or heartburn)
- antifungal medicines (such as ketoconazole, voriconazole, itraconazole and posaconazole)
- antibiotics (such as clarithromycin, erythromycin or telithromycin)
- medicines known as ‘protease inhibitors’ to treat HIV (e.g. boceprevir, ritonavir, indinavir, nelfinavir or saquinavir)
- rifampicin (to treat tuberculosis)
- carbamazepine (a medicine to treat seizures, fits or convulsions and certain pain conditions)
- medicines used to treat seizures, fits or convulsions such as carbamazepine and phenytoin
- a herbal remedy used to treat depression known as St. John’s Wort (also known as *Hypericum perforatum*)
- medicines to treat Parkinson’s disease.

Taking Oxycodone 5mg/5ml Solution with food, drink and alcohol

Drinking alcohol during your treatment with this medicine may make you sleepy or increase the risk of serious side effects such as shallow breathing with a risk of stopping breathing, and loss of consciousness. It is recommended not to drink alcohol while you’re taking Oxycodone 5mg/5ml Solution liquid.

You should avoid drinking grapefruit juice during your treatment with this medicine.

Pregnancy and breast-feeding

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy

Do not take Oxycodone 5mg/5ml Solution if you are pregnant or think you might be pregnant unless you have discussed this with your doctor and the benefits of treatment are considered to outweigh the potential harm to the baby. If you take Oxycodone 5mg/5ml Solution during pregnancy your baby may become dependent and experience withdrawal symptoms after the birth which may need to be treated.

Breast-feeding

Do not take Oxycodone 5mg/5ml Solution while you are breastfeeding as oxycodone passes into breast milk and will affect your baby.

Driving and using machines

This medicine may cause a number of side effects such as drowsiness which could affect your ability to drive or use machinery (see section 4 for a full list of side effects). These are usually more noticeable when you first start taking this medicine, or when changing to a higher dose. If you are affected you should not drive or use machinery. This medicine can affect your ability to drive as it may make you sleepy or dizzy.

- Do not drive while taking this medicine until you know how it affects you
- It is an offence to drive while you have this medicine in your body over a specified limit unless you have a defence (called the ‘statutory defence’)
- However you would not be committing an offence if:
 - The medicine has been prescribed to treat a medical or dental problem and
 - You have taken it according to the instructions given by the prescriber or in the information provided with the medicine and
 - It was not affecting your ability to drive safely.

Talk to your doctor or pharmacist if you are not sure whether it is safe for you to drive while taking this medicine.

Oxycodone 5mg/5ml Solution contains sodium

This medicine contains 4.25mg sodium (main component of cooking/ table salt) in each 5ml. This is equivalent to 0.21% of the recommended maximum daily dietary intake of sodium for an adult.

Oxycodone 5mg/5ml Solution contains sodium benzoate (E211)

This medicine contains 1mg sodium benzoate in each ml, which is equivalent to 5mg/5ml.

Sodium benzoate may increase jaundice (yellowing of the skin and eyes) in newborn babies (up to 4 weeks old).

3. How to take Oxycodone 5mg/5ml Solution

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Your doctor should have discussed with you how long the course of Oxycodone 5mg/ml Solution will last. They will arrange a plan for stopping treatment. This will outline how to gradually reduce the dose and stop taking the medicine.

Before starting treatment and regularly during treatment, your doctor will discuss with you what you may expect from using Oxycodone 5mg/5ml Solution, when and how long you need to take it, when to contact your doctor, and when you need to stop it (see also if you stop using Oxycodone 5mg/5ml Solution).

Adults (over 18 years of age)

The usual starting dose is 5ml every four to six hours. However, your doctor will prescribe the dose required to treat your pain. If you find that you are still in pain whilst taking this medicine discuss this with your doctor.

Children

Children and adolescents under 18 years of age should not take the medicine.

Patients with kidney or liver problems

Please tell your doctor if you suffer from kidney or liver problems as they may prescribe a lower dose depending on your condition.

Do not exceed the dose recommended by your doctor. You should check with your doctor or pharmacist if you are not sure.

Do not drink Oxycodone 5mg/5ml Solution directly from the bottle. Measure out the required dose using the spoon, cup or syringe you have been provided. Drinking directly from the bottle increases the risk of overdose.

You must only take this medicine by mouth. This medicine should never be injected as this may lead to serious side effects, which may be fatal.

If you take more Oxycodone 5mg/5ml Solution than you should or if someone accidentally swallows your medicine

Call your doctor or hospital immediately. People who have taken an overdose may feel very sleepy, sick or dizzy or have hallucinations. They may also have breathing difficulties leading to unconsciousness or even death and may need emergency treatment in hospital. An overdose may result in a brain disorder (known as toxic leukoencephalopathy). When seeking medical attention make sure that you take this leaflet and any remaining medicine with you to show the doctor.

If you forget to take Oxycodone 5mg/5ml Solution

If you remember within 4 hours of the time your medicine was due, take it straight away. Take your next dose at your normal time. If you are more than 4 hours late, please call your doctor or pharmacist for advice. Do not take two doses within four hours. Do not take a double dose to make up for a forgotten dose.

If you stop taking Oxycodone 5mg/5ml Solution

Do not suddenly stop taking this medicine. If you want to stop taking this medicine, discuss this with your doctor first. They will tell you how to do this, usually by reducing the dose gradually so that any unpleasant withdrawal effects are kept to a minimum. Withdrawal symptoms such as restlessness, difficulty sleeping, irritability, agitation, anxiety, feeling your heartbeat (palpitations), increased blood pressure, feeling or being sick, diarrhoea, shaking, shivering or sweating may occur if you suddenly stop taking this medicine.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

All medicines can cause allergic reactions, although serious allergic reactions are rare. **Tell your doctor immediately** if you get any sudden wheeziness, difficulties in breathing, swelling of the eyelids, face or lips, rash or itching especially those covering your whole body. These may be signs of a serious allergic reaction.

The most serious side effect is a condition where you breathe more slowly or weakly than usual (respiratory depression) and can lead to severe sleepiness and loss of consciousness. This side effect may affect up to 1 in 100 people and is more likely to occur when taking certain other medicines (see section 2 ‘Other medicines and Oxycodone 5mg/5ml Solution’). **Tell your doctor immediately** if this happens to you. You may wish to ask your friends, family or caregivers to monitor you for these signs and symptoms.

Drug withdrawal

When you stop taking Oxycodone 5mg/5ml Solution you may experience drug withdrawal symptoms, which include restlessness, difficulty sleeping, irritability, agitation, anxiety, feeling your heartbeat (palpitations), increased blood pressure, feeling or being sick, diarrhoea, shaking, shivering or sweating.

How do I know if I am addicted?

- If you notice any of the following signs whilst taking Oxycodone 5mg/5ml Solution it could be a sign that you have become addicted.
- You need to take the medicine for longer than advised by your prescriber
 - You feel you need to use more than the recommended dose

- You are using the medicine for reasons other than prescribed
 - When you stop taking the medicine you feel unwell, and you feel better once taking the medicine again.
- If you notice any of these signs it is important you talk to your doctor.

Very common side effects

(May affect more than 1 in 10 people)

- Constipation (your doctor can prescribe a laxative to overcome this problem)
- Feeling or being sick (this should normally wear off after a few days, however your doctor can prescribe an anti-sickness medicine if it continues to be a problem)
- Drowsiness (this is most likely when you start taking your medicine or when your dose is increased, but it should wear off after a few days)
- Dizziness
- Headache
- Itchy skin.

Common side effects

(May affect up to 1 in 10 people)

- Dry mouth, loss of appetite, indigestion, abdominal pain or discomfort, diarrhoea
- Confusion, depression, a feeling of unusual weakness, shaking, lack of energy, tiredness, anxiety, nervousness, difficulty in sleeping, abnormal thoughts or dreams
- Difficulty in breathing or wheezing, shortness of breath, decreased cough reflex
- Rash
- Sweating.

Uncommon side effects

(May affect up to 1 in 100 people)

- Withdrawal symptoms (see section ‘Drug withdrawal’)
- Difficulty in swallowing, belching, hiccups, wind, a condition where the bowel does not work properly (ileus), inflammation of the stomach, changes in taste or unpleasant taste
- A feeling of dizziness or spinning (vertigo), hallucinations, mood changes, unpleasant or uncomfortable mood, a feeling of extreme happiness, disorientation, restlessness, agitation, generally feeling unwell, loss of memory, difficulty in speaking, reduced sensitivity to pain or touch, tingling or numbness, seizures, fits or convulsions, blurred or impaired vision, fainting, unusual muscle stiffness or slackness, involuntary muscle contractions or spasms
- Difficulty or pain in passing urine, impotence, decreased sexual drive, low levels of sex hormones in the blood (‘hypogonadism’, seen in a blood test)
- Fast, irregular heartbeat, palpitations, a feeling of lightheadedness. flushing of the skin
- Dehydration, thirst, chills, swelling of the hands, ankles or feet
- Dry skin, severe flaking or peeling of the skin
- Redness of the face, reduction in size of the pupils in the eye, high temperature
- Colicky abdominal pain or discomfort
- A worsening of liver function tests (seen in a blood test).

Rare side effects

(May affect up to 1 in 1,000 people)

- Low blood pressure or feeling faint, especially on standing up
- A raised, itchy rash (hives).

Frequency not known

(Frequency cannot be estimated from the available data)

- Dependence and addiction (see section ‘How do I know if I am addicted?’)
- Withdrawal symptoms (see ‘Drug withdrawal’ in section 2 of the leaflet)
- A need to take increasingly higher doses of this medicine to obtain the same level of pain relief (tolerance)
- An increased sensitivity to pain
- Aggression
- Tooth decay
- Absence of menstrual periods
- A blockage in the flow of bile from the liver (cholestasis). This can cause itchy skin, yellow skin, very dark urine and very pale stools
- Sleep apnoea (breathing pauses during sleep).

Long term use of Oxycodone 5mg/5ml Solution during pregnancy may cause life-threatening withdrawal symptoms in the newborn. Symptoms to look for in the baby include irritability, hyperactivity and abnormal sleep pattern, high pitched cry, shaking, being sick, diarrhoea and not putting on weight.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme, website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Oxycodone 5mg/5ml Solution

Keep this medicine out of the sight and reach of children. Accidental overdose by a child is dangerous and may be fatal. Store this medicine in a locked safe and secure storage space, where other people cannot access it. It can cause serious harm and be fatal to people when it has not been prescribed for them.

- Do not store above 25°C
- Store in the original package in order to protect from light
- Do not use after the expiry date (shown as Exp. On the packaging). The expiry date refers to the last day of the month, your doctor or nurse will check for this
- Once opened this medicine must be used within 1 month
- Consult a pharmacist if signs of deterioration of the product are observed.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Oxycodone 5mg/5ml Solution contains

The active ingredient is oxycodone hydrochloride. Each 5ml contains oxycodone hydrochloride 5mg (equivalent to 4.5mg oxycodone base).

The other ingredients are: citric acid monohydrate, sodium citrate, sodium benzoate (E211), sodium saccharin, hypromellose (methocel-E15 premium LV) and purified water.

What Oxycodone 5mg/5ml Solution looks like and the contents of the pack

This medicine is a clear colourless solution. Each bottle contains 250ml of solution.

Marketing Authorisation Holder:

Wockhardt UK Ltd, Ash Road North, Wrexham, LL13 9UF, UK

Manufacturer:

CP Pharmaceuticals Ltd, Ash Road North, Wrexham, LL13 9UF, UK

Other formats:

To listen to or request a copy of this leaflet in Braille, large print or audio please call, free of charge: 0800 198 5000 (UK Only). Please be ready to give the following information:

Product name	Reference number
Oxycodone Hydrochloride 5mg/5ml Oral Solution	29831/0458

This is a service provided by the Royal National Institute of Blind People. **This leaflet was last revised in 02/2025**