

PACKAGE LEAFLET

Package leaflet: Information for the user

Ziclag 30mg Prolonged-Release Tablet

Gliclazide

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again
- If you have any further questions, ask your doctor or pharmacist
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Ziclag Tablets are and what they are used for
2. What you need to know before you take Ziclag Tablets
3. How to take Ziclag Tablets
4. Possible side effects
5. How to store Ziclag Tablets
6. Contents of the pack and other information

1. What Ziclag Tablets are and what they are used for

Gliclazide-Lupin 30mg prolonged-release tablet.

Each tablet contains 30 mg Gliclazide.

Pharmacotherapeutic group: Sulfonamides, urea derivatives, ATC code: A10BB09

Ziclag is a medicine which **reduces blood sugar levels** (oral antidiabetic medicine belonging to the sulphonylurea group).

Ziclag 30mg prolonged release tablets are used to keep blood sugar at the correct level in adults with **diabetes** (type 2 diabetes mellitus) when it is not controlled by dietary measures, physical exercise and weight loss alone.

2. What you need to know before you take Ziclag Tablets

Do not take Ziclag 30mg prolonged release tablets:

- if you are **allergic** (hypersensitive) to **gliclazide** or any of the other ingredients in this medicine (see section 6, "Further Information") or to any other medicines of the same group (sulphonylureas or sulphonamides).
- if you have insulin-dependent (Type 1) **diabetes**;
- if you have ketone bodies and sugar in your urine (this may mean you have **diabetic keto-acidosis**), diabetic pre-coma or coma
- if you have **severe kidney or liver disease**;
- if you are taking any medicines used to treat **fungal infections** (miconazole – see Section "Taking other medicines")
- if you are **breast-feeding** (see section "Pregnancy and breastfeeding")

Warnings and Precautions

Talk to your doctor or pharmacist before taking Gliclazide 30 mg prolonged release tablets.

You should observe the treatment plan prescribed by your doctor to achieve proper blood sugar levels. This means, apart from regular tablet intake, you observe the dietary regimen, have physical exercise and, where necessary, reduce weight.

During gliclazide treatment regular monitoring of your blood (and possibly urine) sugar level and also your glycated haemoglobin (HbA1c) is necessary.

In the first few weeks of treatment the risk of having reduced blood sugar levels (hypoglycaemia) may be increased. So particularly close medical monitoring is necessary.

Low blood sugar (Hypoglycaemia) may occur:

- if you take meals irregularly or skip meals altogether,
- if you are fasting,
- if you are malnourished,
- if you change your diet,
- if you increase your physical activity and carbohydrate intake does not match this increase,
- if you drink alcohol, especially in combination with skipped meals,
- if you take other medicines or natural remedies at the same time,
- if you take too high doses of gliclazide,
- if you suffer from particular hormone-induced disorders (functional disorders of the thyroid gland, of the pituitary gland or adrenal cortex),
- if your kidney function or liver function is severely decrease

If you have low blood sugar you may have the following symptoms:

headache, intense hunger, nausea, vomiting, weariness, sleep disorders, restlessness, aggressiveness, poor concentration, reduced alertness and reaction time, depression, confusion, speech or visual disorders, tremor, sensory disturbances, dizziness, and helplessness.

The following signs and symptoms may also occur: sweating, clammy skin, anxiety, fast or irregular heartbeat, high blood pressure, sudden strong pain in the chest that may radiate into nearby areas (angina pectoris).

If blood sugar levels continue to drop you may suffer from considerable confusion (delirium), develop convulsions, lose self-control, your breathing may be shallow, and your heart beat slowed down, you may become unconscious.

In most cases the symptoms of low blood sugar vanish very quickly when you consume some form of sugar, e.g. glucose tablets, sugar cubes, sweet juice, sweetened tea. You should therefore always carry some form of sugar with you (glucose tablets, sugar cubes). Remember that artificial sweeteners are not effective. Please contact your doctor or the nearest hospital if taking sugar does not help or if the symptoms recur.

Symptoms of low blood sugar may be absent, less obvious or develop very slowly or you are not aware in time that your blood sugar level has dropped. This may happen if you are an elderly patient taking certain medicines (e.g. those acting on the central nervous system and beta blockers). If you are in stress-situations (e.g. accidents, surgical operations, fever etc.) your doctor may temporarily switch you to insulin therapy.

Symptoms of high blood sugar (hyperglycaemia) may occur when gliclazide has not yet sufficiently reduced the blood sugar, when you have not complied with the treatment plan prescribed by your doctor or in special stress situations. These may include thirst, frequent urination, dry mouth, dry itchy skin, skin infections and reduced performance.

If these symptoms occur, you must contact your doctor or pharmacist.

If you have a family history of or know you have the hereditary condition glucose-6-phosphate dehydrogenase (G6PD) deficiency (abnormality of red blood cells), lowering of the haemoglobin

level and breakdown of red blood cells (haemolytic anaemia) can occur. Contact your doctor before taking this medicinal product.

Children and adolescents

Gliclazide 30 mg prolonged release tablets is not recommended for use in children due to a lack of data.

Other medicines and Gliclazide 30 mg prolonged release tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

The blood sugar lowering **effect of Gliclazide may be strengthened** and signs of **low blood sugar levels** may occur when one of the following medicines is taken:

- other medicines used to treat high blood sugar (oral anti-diabetics, GLP-1 receptor inhibitors or insulin)
- antibiotics (e.g. sulfonamides, clarithromycin),
- medicines used to treat high blood pressure or heart failure (beta blockers, ACE inhibitors such as captopril or enalapril)
- medicines to treat fungal infections (miconazole, fluconazole)
- medicines to treat ulcers in the stomach or duodenum (H2 receptor antagonists, medicines to treat depression (monoamine oxidase inhibitors)
- **Pain killers** or **anti-rheumatics**, such as ibuprofen or phenylbutazone
- medicines containing **alcohol**

If you take one of the following medicines, the **effect of Gliclazide may be weakened** and **raised blood sugar levels may occur**:

- medicines to treat disorders of the central **nervous system** (chlorpromazine)
- medicines reducing **inflammation** (corticosteroids)
- medicines to treat **asthma or used labour** (intravenous salbutamol, ritodrine, terbutaline)
- medicines to treat breast disorders, heavy menstrual bleeding and endometriosis (danazol)
- St John's Wort – *Hypericum perforatum* – preparations.

Blood glucose disturbance (low blood sugar and high blood sugar) can occur when a medicine belonging to a class of antibiotic called fluroquinolones is taken at the same time than Gliclazide 30mg Prolonged release tablets, especially in elderly patients.

Gliclazide may increase the effects of medicines used to **reduce blood clotting (warfarin)**.

Consult your doctor before you start taking another medicine. If you go to the hospital tell the medical staff, you are taking Ziclagse prolonged release tablets.

Ziclagse 30mg prolonged release tablets with food, drink and alcohol:

Ziclagse Tablets can be taken with food and non-alcoholic drinks. Drinking alcohol is not recommended as it can alter the control of your diabetes in an unpredictable manner.

Pregnancy and breast-feeding:

Pregnancy:

Ziclagse 30mg prolonged release tablets are **not recommended for use during pregnancy**. If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby while taking this medicine.

Breast-feeding:

You must not take Ziclagse 30mg prolonged release tablets while you are breast-feeding.

Driving and Using Machines:

Your ability to concentrate or react may be impaired if your blood sugar is too low (hypoglycaemia), or too high (hyperglycaemia) or if you develop visual problems as a result of such conditions. Bear in mind that you could endanger yourself or others (i.e. when driving a car or using machines).

Please **ask your doctor whether you can drive** a car if you:

- have frequent episodes of low blood sugar (hypoglycaemia);
- have few or no warning signals of low blood sugar (hypoglycaemia).

3. How to take Ziclag Tablets

Always take Ziclag 30mg prolonged release tablets exactly as your doctor or pharmacist has told you. You should check with your doctor or pharmacist if you are not sure.

Dose

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The dose is determined by the doctor, depending on your blood and possibly urine sugar levels.

Change in external factors (e.g. weight reduction, change in life style, stress) or improvements in the blood sugar control may require changed gliclazide doses.

The recommended dose is one to four tablets (maximum 120 mg) in a single intake at breakfast time. This depends on the response to treatment.

If a combination therapy of Gliclazide 30 mg prolonged release tablets with metformin, an alpha glucosidase inhibitor, a thiazolidinedione, a dipeptidyl peptidase-4 inhibitor, a GLP-1 receptor agonist or insulin is initiated your doctor will determine the proper dose of each medicine individually for you.

If you notice that your blood sugar levels are high although you are taking the medicine as prescribed, you should contact your doctor or pharmacist.

Route and method of administration

Swallow the tablet(s) whole. Do not chew them. Take your tablets with a glass of water at **breakfast** time (and preferably at the same time each day). You **must always eat a meal after** taking your tablet(s).

If you take more Ziclag Tablets than you should:

If you have taken too many tablets than you should, **contact your doctor or the nearest hospital immediately.**

The signs of overdose are those of low blood sugar (hypoglycaemia) described in section 2. The symptoms of can be helped by taking sugar (4 to 6 lumps) or sugary drinks straight away, followed by a substantial snack or meal. If the patient is unconscious immediately inform a doctor and call the emergency services. The same should be done if somebody, e.g. a child, has taken the product unintentionally. Unconscious patients must not be given food or drink.

It should be ensured that there is always a pre-informed person that can call a doctor in case of emergency.

If you forget to take Ziclag tablets:

It is important to take your medicine every day as regular treatment works better.

However, if you forget to take a dose of Gliclazide 30 mg prolonged release tablets take the next dose at the usual time. Do not take a double dose to make up for a forgotten dose.

If you stop taking Ziclagseg Tablets:

As the treatment for diabetes is usually life long, you should discuss with your doctor before stopping this medicinal product. Stopping could cause high blood sugar (hyperglycaemia) which increases the risk of developing complications of diabetes.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, Ziclagseg Tablets can cause side-effects, although not everybody gets them. The most commonly observed side effect is low blood sugar (hypoglycaemia). For symptoms and signs see Section "Warnings and precautions" in "Section 2 What you need to know before you take".

If left untreated these symptoms could progress to drowsiness, loss of consciousness or possibly coma. If an episode of low blood sugar is severe or prolonged, even if it is temporarily controlled by eating sugar, you should seek immediate medical attention.

Blood disorders

Decrease in the number of cells in the blood has been reported (e.g. platelets, red and white blood cells).

This may cause: paleness, prolonged bleeding, bruising, sore throat, fever

These symptoms usually vanish when the treatment is discontinued.

Liver disorders

There have been isolated reports of abnormal liver function, which can cause yellow skin and eyes. If you get this, see your doctor immediately. The symptoms generally disappear if the medicine is stopped. Your doctor will decide whether to stop your treatment.

Skin disorders

Skin reactions have been reported such as:

- rash, redness, itching, hives,
- angioedema (rapid swelling of tissues such as eyelids, face, lips, mouth, tongue or throat that may result in breathing difficulty). The rash may progress to widespread blistering or peeling of the skin.
- exceptionally, signs of severe hypersensitivity reactions (DRESS): initially as flu-like symptoms and a rash on the face then an extended rash with a high temperature.

Digestive disorders

- Stomach pain or discomfort, feeling or being sick, indigestion, diarrhoea, constipation.

These effects are reduced when Gliclazide 30 mg prolonged release tablet is taken with a meal as recommended, See Section 3 "How to take".

Eye disorders

Your vision may be affected for a short time especially at the start of treatment. This effect is due to changes in blood sugar levels.

As for other sulfonylureas, the following adverse events have been observed: cases of severe changes in the number of blood cells and allergic inflammation of the wall of blood vessels, reduction in blood sodium (hyponatraemia), symptoms of liver impairment (e.g. jaundice) which in most cases disappeared after withdrawal of the sulfonylureas but may lead to life-threatening liver failure in isolated cases.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

You can also report side effects directly via the Yellow Card Scheme, website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Ziclag Tablets

Keep out of the sight and reach of children.

Do not use this medicine after the expiry date, which is stated on the carton and blister. The expiry date refers to the last day of that month.

Do not store above 25°C.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Ziclag Tablets contain:

The active substance is gliclazide.

Each prolonged release tablet contains 30mg of the active substance gliclazide.

The other ingredients are:

calcium hydrogen phosphate dihydrate, povidone K30, hypromellose and magnesium stearate.

What Ziclag Tablets look like and contents of the pack:

Ziclag 30mg prolonged release tablets are white to off-white capsule shaped, biconvex tablets debossed with “30” on one side and plain on the other side.

Ziclag 30mg prolonged release tablets are available in blister packs containing 10, 20, 28, 30, 56, 60, 90 and 120 tablets* packaged in cartons.

* Not all pack sizes may be marketed

Marketing Authorisation Holder and Manufacturer

Lupin Healthcare (UK) Limited

The Urban Building, 2nd floor

3-9 Albert Street, Slough, Berkshire

SL1 2BE, United Kingdom

This leaflet was last revised in September 2019